

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
6 December 2001 (06.12.2001)

PCT

(10) International Publication Number
WO 01/92235 A1

(51) International Patent Classification⁷: **C07D 243/24**,
223/18, 223/16, 223/12, 243/14, 243/12, A61K 31/55,
31/5513, A61P 25/28

(74) Agent: **LARSEN, Scott, K.**; Dupont Pharmaceuticals
Company, Legal Patent Records Center, 1007 Market
Street, Wilmington, DE 19898 (US).

(21) International Application Number: **PCT/US01/17865**

(81) Designated States (*national*): AT, AU, BR, CA, CH, CN,
CZ, DE, DK, EE, ES, FI, GB, HU, IL, IN, JP, KR, LT, LU,
LV, MX, NO, NZ, PL, PT, RO, RU, SE, SG, SI, SK, UA,
VN, ZA.

(22) International Filing Date: **1 June 2001 (01.06.2001)**

(25) Filing Language: **English**

(84) Designated States (*regional*): Eurasian patent (AM, AZ,
BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE,
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,
NL, PT, SE, TR).

(26) Publication Language: **English**

(30) Priority Data:
60/208,536 **1 June 2000 (01.06.2000) US**

Published:


- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

(71) Applicant: **DUPONT PHARMACEUTICALS COM-
PANY [US/US]**; Chestnut Run Plaza, 974 Centre Road,
Wilmington, DE 19805 (US).

(72) Inventor: **OLSON, Richard, E.**; 7 Pelham Road, Wilm-
ington, DE 19803 (US).

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: **LACTAMS SUBSTITUTED BY CYCLIC SUCCINATES AS INHIBITORS OF A β PROTEIN PRODUCTION**



(I)

(57) Abstract: This invention relates to novel
lactams having formula (I) to their pharmaceutical
compositions and to their methods of use. These
novel compounds inhibit the processing of amyloid
precursor protein and, more specifically, inhibit
the production of A β -peptide, thereby acting to

prevent the formation of neurological deposits of amyloid protein. More particularly, the present invention relates to the treatment
of neurological disorders related to β -amyloid production such as Alzheimer's disease and Down's Syndrome.

AQ

LACTAMS SUBSTITUTED BY CYCLIC SUCCINATES AS INHIBITORS OF
A β PROTEIN PRODUCTION

5

FIELD OF THE INVENTION

This invention relates to novel lactams substituted by cyclic succinates having drug and bio-affecting properties, their pharmaceutical compositions and methods of use. These novel compounds inhibit the processing of amyloid precursor protein and, more specifically, inhibit the production of A β -peptide, thereby acting to prevent the formation of neurological deposits of amyloid protein. More particularly, the present invention relates to the treatment of neurological disorders related to β -amyloid production such as Alzheimer's disease and Down's Syndrome.

BACKGROUND OF THE INVENTION

Alzheimer's disease (AD) is a degenerative brain disorder characterized clinically by progressive loss of memory, temporal and local orientation, cognition, reasoning, judgment and emotional stability. AD is a common cause of progressive dementia in humans and is one of the major causes of death in the United States. AD has been observed in all races and ethnic groups worldwide, and is a major present and future health problem. No treatment that effectively prevents AD or reverses the clinical symptoms and underlying pathophysiology is currently available (for review see Dennis J. Selkoe; Cell Biology of the amyloid (beta)-protein precursor and the mechanism of Alzheimer's disease, Annu Rev Cell Biol, 1994, 10: 373-403).

Histopathological examination of brain tissue derived upon autopsy or from neurosurgical specimens in effected individuals revealed the occurrence of amyloid plaques and neurofibrillar tangles in the cerebral cortex of such patients. Similar alterations were observed in patients with Trisomy 21 (Down's syndrome), and hereditary cerebral hemorrhage with amyloidosis of the Dutch-type.

Neurofibrillar tangles are nonmembrane-bound bundles of abnormal proteinaceous filaments and biochemical and immunochemical studies led to the conclusion that their principle protein subunit is an altered phosphorylated form of the tau protein (reviewed in Selkoe, 1994).

Biochemical and immunological studies revealed that the dominant proteinaceous component of the amyloid plaque is an approximately 4.2 kilodalton (kD) protein of about 39 to 43 amino acids. This protein was designated A β , β -amyloid peptide, and sometimes β /A4; referred to herein as A β . In addition to its deposition in amyloid plaques, A β is also found in the walls of meningeal and parenchymal arterioles, small arteries, capillaries, and sometimes, venules. A β was first purified and a partial amino acid reported in 1984 (Glenner and Wong, Biochem. Biophys. Res. Commun. 120: 885-890). The isolation and sequence data for the first 28 amino acids are described in U.S. Pat. No 4,666,829.

Compelling evidence accumulated during the last decade revealed that A β is an internal polypeptide derived from a type 1 integral membrane protein, termed β amyloid precursor protein (APP). β APP is normally produced by many cells both in vivo and in cultured cells, derived from various animals and humans. A β is derived from cleavage of β APP by as yet unknown enzyme (protease) system(s), collectively termed secretases.

The existence of at least four proteolytic activities has been postulated. They include β secretase(s), generating the N-terminus of A β , α secretase(s) cleaving around the 16/17 peptide bond in A β , and γ secretases, generating C-terminal A β fragments ending at position 38, 39, 40, 42, and 43 or generating C-terminal extended precursors which are subsequently truncated to the above polypeptides.

Several lines of evidence suggest that abnormal accumulation of A β plays a key role in the pathogenesis of AD. Firstly, A β is the major protein found in amyloid plaques. Secondly, A β is neurotoxic and may be causally


related to neuronal death observed in AD patients.

Thirdly, missense DNA mutations at position 717 in the 770 isoform of β APP can be found in effected members but not unaffected members of several families with a genetically determined (familiar) form of AD. In addition, several other β APP mutations have been described in familiar forms of AD. Fourthly, similar neuropathological changes have been observed in transgenic animals overexpressing mutant forms of human β APP. Fifthly, individuals with Down's syndrome have an increased gene dosage of β APP and develop early-onset AD. Taken together, these observations strongly suggest that A β depositions may be causally related to the AD.

It is hypothesized that inhibiting the production of A β will prevent and reduce neurological degeneration, by controlling the formation of amyloid plaques, reducing neurotoxicity and, generally, mediating the pathology associated with A β production. One method of treatment methods would therefore be based on drugs that inhibit the formation of A β in vivo.


Methods of treatment could target the formation of A β through the enzymes involved in the proteolytic processing of β amyloid precursor protein. Compounds that inhibit β or γ secretase activity, either directly or indirectly, could control the production of A β . Advantageously, compounds that specifically target γ secretases, could control the production of A β . Such inhibition of β or γ secretases could thereby reduce production of A β , which, thereby, could reduce or prevent the neurological disorders associated with A β protein.

PCT publication number WO 96/29313 discloses the general formula:




covering metalloprotease inhibiting compounds useful for the treatment of diseases associated with excess and/or unwanted matrix metalloprotease activity, particularly collagenase and or stromelysin activity.

5 Compounds of general formula:




are disclosed in PCT publication number WO 95/22966 relating to matrix metalloprotease inhibitors. The compounds of the invention are useful for the treatment of conditions associated with the destruction of cartilage, including corneal ulceration, osteoporosis, periodontitis and cancer.

15 European Patent Application number EP 0652009A1 relates to the general formula:



and discloses compounds that are protease inhibitors that inhibit A β production.


US Patent Number 5703129 discloses the general formula:



25 which covers 5-amino-6-cyclohexyl-4-hydroxy-hexanamide derivatives that inhibit A β production and are useful in the treatment of Alzheimer's disease.

Copending, commonly assigned U.S. patent application Serial Number 09/370089 filed August 7, 1999 (equivalent to

international application PCT US99/17717) discloses lactams of general formula:



5

wherein the lactam ring B is substituted by succinamide and a carbocyclic, aryl, or heteroaryl group. These compounds inhibit the processing of amyloid precursor protein and, more specifically, inhibit the production of A β -peptide, thereby acting to prevent the formation of neurological deposits of amyloid protein.

None of the above references teaches or suggests the compounds of the present invention which are described in detail below.

15

SUMMARY OF THE INVENTION


One object of the present invention is to provide novel compounds which are useful as inhibitors of the production of A β protein or pharmaceutically acceptable salts or prodrugs thereof.

It is another object of the present invention to provide pharmaceutical compositions comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of at least one of the compounds of the present invention or a pharmaceutically acceptable salt or prodrug form thereof.

It is another object of the present invention to provide a method for treating degenerative neurological disorders comprising administering to a host in need of such treatment a therapeutically effective amount of at least one of the compounds of the present invention or a pharmaceutically acceptable salt or prodrug form thereof.

These and other objects, which will become apparent during the following detailed description, have been

achieved by the inventors' discovery that compounds of Formula (I):




(I)

5 or pharmaceutically acceptable salt or prodrug forms thereof, wherein R^3 and R^{3a} are combined to form a carbocyclic or heterocyclic ring, R^5 , R^{5a} , R^6 , Q , B , W , X , Y , and Z are defined below, are effective inhibitors of the
10 production of $A\beta$.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

Thus, in a first embodiment, the present invention provides a novel compound of Formula (I):



(I)

or a pharmaceutically acceptable salt or prodrug thereof,
20 wherein:

Q is $-NR^1R^2$;

25 R^1 is H, C_1 - C_4 alkyl, phenyl, benzyl, C_3 - C_6 cycloalkyl, or (C_3 - C_6 cycloalkyl)methyl-;

R^2 is H, C_1 - C_4 alkyl, OH, C_1 - C_4 alkoxy, phenyl, benzyl, C_3 - C_6 cycloalkyl, or (C_3 - C_6 cycloalkyl)methyl-;

30 R^3 and R^{3a} may be combined to form a 3-8 membered carbocyclic moiety;

wherein said 3-8 membered carbocyclic moiety is saturated or partially unsaturated;

wherein said 3-8 membered carbocyclic moiety may optionally contain a heteroatom selected from -O-,
5 -S-, -S(=O)-, -S(=O)₂-, -N=, -NH-, and -N(R²⁰)-, and
wherein said 3-8 membered carbocyclic moiety is substituted with 0-4 R⁴;

10 additionally, two R⁴ substituents on adjacent atoms may be combined to form a benzo fused radical; wherein said benzo fused radical is substituted with 0-4 R²³;

15 additionally, two R⁴ substituents on adjacent atoms may be combined to form a 5 to 6 membered heteroaryl fused radical, wherein said 5 to 6 membered heteroaryl fused radical comprises 1 or 2 heteroatoms selected from N, O, and S; wherein said 5 to 6 membered heteroaryl fused radical is substituted with 0-3 R²³;

20 additionally, two R⁴ substituents on the same or adjacent carbon atoms may be combined to form a C₃-C₆ carbocycle substituted with 0-3 R²³;

25 alternatively, R³ is H; C₁-C₆ alkyl substituted with 0-3 R⁴; C₂-C₆ alkenyl substituted with 0-3 R⁴; or C₂-C₆ alkynyl substituted with 0-3 R⁴; and R^{3a} is H, C₁-C₆ alkyl, or C₂-C₆ alkenyl;

30 R⁴, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, NR¹⁵R¹⁶, OR^{14a}, C₁-C₄ alkyl, C₂-C₆ alkenyl, alkynyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-,
35 C₃-C₆ carbocycle, aryl, and a 5 to 6 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur;

- R⁵ is H, C₁-C₆ alkoxy;
C₁-C₆ alkyl substituted with 0-3 R^{5b};
C₂-C₆ alkenyl substituted with 0-3 R^{5b};
C₂-C₆ alkynyl substituted with 0-3 R^{5b};
5 C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};
C₆-C₁₀ aryl substituted with 0-3 R^{5c}; or
5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered heterocycle
10 is substituted with 0-3 R^{5c};
- R^{5a} is H, C₁-C₄ alkyl, or C₂-C₄ alkenyl;
- R^{5b}, at each occurrence, is independently selected from:
15 H, C₁-C₆ alkyl, CF₃, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂,
NR¹⁵R¹⁶;
C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};
C₆-C₁₀ aryl substituted with 0-3 R^{5c}; or
5 to 10 membered heterocycle containing 1 to 4
20 heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered heterocycle
is substituted with 0-3 R^{5c};
- R^{5c}, at each occurrence, is independently selected from H,
25 OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
S(=O)CH₃, S(=O)₂CH₃,
C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,
C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;
- 30 alternatively, R⁵ and R^{5a} may be combined to form a 3-8
membered carbocyclic moiety;
wherein said 3-8 membered carbocyclic moiety is
saturated or partially unsaturated;
wherein said 3-8 membered carbocyclic moiety may
35 optionally contain a heteroatom selected from -O-,
-S-, -S(=O)-, -S(=O)₂-, -N=, -NH-, and -N(R²⁰)-, and
wherein said 3-8 membered carbocyclic moiety is
substituted with 0-4 R^{5c};

provided at least:

- 1) R^3 and R^{3a} are combined to form a 3-8 membered carbocyclic moiety; or
- 2) R^5 and R^{5a} are combined to form a 3-8 membered carbocyclic moiety; or
- 3) R^3 and R^{3a} are combined to form a 3-8 membered carbocyclic moiety and R^5 and R^{5a} are combined to form a 3-8 membered carbocyclic moiety;

R^6 is H;

C_1 - C_6 alkyl substituted with 0-3 R^{6a} ;
 C_3 - C_{10} carbocycle substituted with 0-3 R^{6b} ; or
 C_6 - C_{10} aryl substituted with 0-3 R^{6b} ;

R^{6a} , at each occurrence, is independently selected from H, C_1 - C_6 alkyl, OR^{14} , Cl, F, Br, I, =O, CN, NO_2 , $NR^{15}R^{16}$, aryl or CF_3 ;

R^{6b} , at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO_2 , $NR^{15}R^{16}$, CF_3 , C_1 - C_6 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl, and C_1 - C_4 haloalkoxy;

Ring B is a 6, 7, or 8 membered lactam,

wherein the lactam is saturated, partially saturated or unsaturated;

wherein each additional lactam carbon is substituted with 0-2 R^{11} ; and,

optionally, the lactam contains an additional

heteroatom selected from -O-, -S-, -S(=O)-, -S(=O)₂-, -N=, -NH-, and -N(R^{10})-;

additionally, two R^{11} substituents on adjacent atoms may be combined to form a benzo fused radical; wherein said benzo fused radical is substituted with 0-4 R^{13} ;

additionally, two R^{11} substituents on adjacent atoms may be combined to form a 5 to 6 membered heteroaryl fused

radical, wherein said 5 to 6 membered heteroaryl fused radical comprises 1 or 2 heteroatoms selected from N, O, and S; wherein said 5 to 6 membered heteroaryl fused radical is substituted with 0-3 R¹³;

5

additionally, two R¹¹ substituents on the same or adjacent carbon atoms may be combined to form a C₃-C₆ carbocycle substituted with 0-3 R¹³;

10 R¹⁰ is H, C(=O)R¹⁷, C(=O)OR¹⁷, C(=O)NR¹⁸R¹⁹, S(=O)₂NR¹⁸R¹⁹, S(=O)₂R¹⁷;

C₁-C₆ alkyl optionally substituted with 0-3 R^{10a};

C₆-C₁₀ aryl substituted with 0-4 R^{10b};

C₃-C₁₀ carbocycle substituted with 0-3 R^{10b}; or

15 5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{10b};

20 R^{10a}, at each occurrence, is independently selected from H, C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶, CF₃, aryl substituted with 0-4 R^{10b}; or a 5-6 membered heterocycle substituted with 0-4 R^{10b};

25 R^{10b}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

30 R¹¹, at each occurrence, is independently selected from H, C₁-C₄ alkoxy, Cl, F, Br, I, =O, CN, NO₂, NR¹⁸R¹⁹, C(=O)R¹⁷, C(=O)OR¹⁷, C(=O)NR¹⁸R¹⁹, S(=O)₂NR¹⁸R¹⁹, CF₃; C₁-C₆ alkyl optionally substituted with 0-3 R^{11a};

C₆-C₁₀ aryl substituted with 0-3 R^{11b};

35 C₃-C₁₀ carbocycle substituted with 0-3 R^{11b}; or

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{11b};

5 R^{11a}, at each occurrence, is independently selected from
H, C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂,
NR¹⁵R¹⁶, CF₃;
phenyl substituted with 0-3 R^{11b};
C₃-C₆ cycloalkyl substituted with 0-3 R^{11b}; and
5 to 6 membered heterocycle containing 1 to 4
10 heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 6 membered heterocycle
is substituted with 0-3 R^{11b};

15 R^{11b}, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
S(=O)CH₃, S(=O)₂CH₃,
C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,
C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

20 W is a bond or -(CR⁸R^{8a})_p-;

p is 0, 1, 2, 3, or 4;

25 R⁸ and R^{8a}, at each occurrence, are independently selected
from H, F, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl
and C₃-C₈ cycloalkyl;

X is a bond;

30 C₆-C₁₀ aryl substituted with 0-3 R^{Xb};
C₃-C₁₀ carbocycle substituted with 0-3 R^{Xb}; or
5 to 10 membered heterocycle substituted with 0-2 R^{Xb};

35 R^{Xb}, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄
haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ halothioalkoxy;

Y is a bond or -(CR⁹R^{9a})_t-V-(CR⁹R^{9a})_u-;

t is 0, 1, or 2;

u is 0, 1, or 2;

5

R⁹ and R^{9a}, at each occurrence, are independently selected from H, F, C₁-C₆ alkyl or C₃-C₈ cycloalkyl;

10 V is a bond, -C(=O)-, -O-, -S-, -S(=O)-, -S(=O)₂-, -N(R¹⁹)-, -C(=O)NR^{19b}-, -NR^{19b}C(=O)-, -NR^{19b}S(=O)₂-, -S(=O)₂NR^{19b}-, -NR^{19b}S(=O)-, -S(=O)NR^{19b}-, -C(=O)O-, or -OC(=O)-;

Z is H;

15 C₁-C₈ alkyl substituted with 0-3 R^{12a};
C₂-C₆ alkenyl substituted with 0-3 R^{12a};
C₂-C₆ alkynyl substituted with 0-3 R^{12a};
C₆-C₁₀ aryl substituted with 0-4 R^{12b};
C₃-C₁₀ carbocycle substituted with 0-4 R^{12b}; or
20 5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{12b};

25 R^{12a}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, -C(=O)NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, C₁-C₄ haloalkyl-S-,
30 C₆-C₁₀ aryl substituted with 0-4 R^{12b};
C₃-C₁₀ carbocycle substituted with 0-4 R^{12b}; or
5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle
35 is substituted with 0-3 R^{12b};

R^{12b}, at each occurrence, is independently selected from

H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₃-C₆ cycloalkyl, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, C₁-C₄ haloalkyl-S, and aryl substituted with 0-3 R^{12c};

R^{12c}, at each occurrence, is independently selected from H, methyl, ethyl, propyl, methoxy, ethoxy, amino, hydroxy, Cl, F, Br, I, CF₃, SCH₃, S(O)CH₃, SO₂CH₃, -N(CH₃)₂, N(CH₃)H, CN, NO₂, OCF₃, C(=O)CH₃, CO₂H, CO₂CH₃, and C₁-C₃ haloalkyl;

R¹³, at each occurrence, is independently selected from H, OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, and CF₃;

R¹⁴ is H, phenyl, benzyl, C₁-C₆ alkyl, C₂-C₆ alkoxyalkyl, or C₃-C₆ cycloalkyl;

R^{14a} is H, phenyl, benzyl, or C₁-C₄ alkyl;

R¹⁵, at each occurrence, is independently selected from H, C₁-C₆ alkyl, benzyl, phenethyl, (C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;

R¹⁶, at each occurrence, is independently selected from H, C₁-C₆ alkyl, benzyl, phenethyl, (C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;

R¹⁷ is H, C₁-C₆ alkyl, C₂-C₆ alkoxyalkyl, aryl substituted by 0-4 R^{17a}, or -CH₂-aryl substituted by 0-4 R^{17a};

R^{17a} is H, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, butoxy, -OH, F, Cl, Br, I, CF₃, OCF₃, SCH₃, S(O)CH₃, SO₂CH₃, -NH₂, -N(CH₃)₂, or C₁-C₄ haloalkyl;

R¹⁸, at each occurrence, is independently selected from

H, C₁-C₆ alkyl, phenyl, benzyl, phenethyl,
(C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;

5 R¹⁹, at each occurrence, is independently selected from
H, OH, C₁-C₆ alkyl, phenyl, benzyl, phenethyl,
(C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;

10 additionally, R¹⁸ and R¹⁹, when substituents on the same
atom, may be combined to form a 3 to 6 membered
heterocyclic ring;

R^{19b}, at each occurrence, is independently is H or C₁-C₄
alkyl;

15 R²⁰ is H, C(=O)R¹⁷, C(=O)OR¹⁷, C(=O)NR¹⁸R¹⁹,
S(=O)₂NR¹⁸R¹⁹, S(=O)₂R¹⁷;
C₁-C₆ alkyl optionally substituted with 0-3 R^{20a}; or
C₆-C₁₀ aryl substituted with 0-4 R^{20b};


20 R^{20a}, at each occurrence, is independently selected from H,
C₁-C₄ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶,
CF₃, or aryl substituted with 0-4 R^{20b};

25 R^{20b}, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
S(=O)CH₃, S(=O)₂CH₃, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄
haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

30 R²³, at each occurrence, is independently selected from
H, OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN,
NO₂, NR¹⁵R¹⁶, and CF₃.

[2] In a preferred embodiment the present invention
provides for a compound of Formula (II):

35



(II)

or a pharmaceutically acceptable salt or prodrug thereof,
 5 wherein:

R³ and R^{3a} may be combined to form a 3-8 membered
 carbocyclic moiety;
 wherein said 3-8 membered carbocyclic moiety is
 10 saturated or partially unsaturated; and

R⁵ and R^{5a} may be combined to form a 3-8 membered
 carbocyclic moiety;
 wherein said 3-8 membered carbocyclic moiety is
 15 saturated or partially unsaturated;

provided at least:

- 1) R³ and R^{3a} are combined to form a 3-8 membered
 carbocyclic moiety; or
- 20 2) R⁵ and R^{5a} are combined to form a 3-8 membered
 carbocyclic moiety; or
- 3) R³ and R^{3a} are combined to form a 3-8 membered
 carbocyclic moiety and R⁵ and R^{5a} are combined to form
 a 3-8 membered carbocyclic moiety.

25 [3] In a preferred embodiment the present invention
 provides for a compound of Formula (II) wherein:

R³ and R^{3a} may be combined to form a 3-8 membered
 30 carbocyclic moiety selected from cyclopropyl,
 cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl,
 and cyclohexenyl;

alternatively, R^3 and R^{3a} are independently selected from the group H, C_1 - C_6 alkyl, C_2 - C_6 alkenyl, and C_2 - C_6 alkynyl;

5 R^5 is H;

C_1 - C_4 alkyl substituted with 0-1 R^{5b} ;

C_2 - C_4 alkenyl substituted with 0-1 R^{5b} ;

C_2 - C_4 alkynyl substituted with 0-1 R^{5b} ;

10 R^{5a} is H, C_1 - C_4 alkyl, or C_2 - C_4 alkenyl;

R^{5b} is selected from:

H, methyl, ethyl, propyl, butyl, CF_3 , Cl, F, $NR^{15}R^{16}$,

cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl;

15

alternatively, R^5 and R^{5a} may be combined to form a 3-8 membered carbocyclic moiety selected from cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, and cyclohexenyl;

20

provided at least:

1) R^3 and R^{3a} are combined to form a 3-8 membered carbocyclic moiety; or

2) R^5 and R^{5a} are combined to form a 3-8 membered carbocyclic moiety; or

25

3) R^3 and R^{3a} are combined to form a 3-8 membered carbocyclic moiety and R^5 and R^{5a} are combined to form a 3-8 membered carbocyclic moiety;

30 Ring B is a 7 membered lactam,

wherein the lactam is saturated, partially saturated or unsaturated;

wherein each additional lactam carbon is substituted with 0-2 R^{11} ; and,

35

optionally, the lactam contains an additional heteroatom selected from -O-, -S-, -S(=O)-, -S(=O)₂-, -N=, -NH-, and -N(R^{10})-;


additionally, two R^{11} substituents on adjacent atoms may be combined to form a benzo fused radical; wherein said benzo fused radical is substituted with 0-4 R^{13} ;

5 additionally, two R^{11} substituents on adjacent atoms may be combined to form a 5 to 6 membered heteroaryl fused radical, wherein said 5 to 6 membered heteroaryl fused radical comprises 1 or 2 heteroatoms selected from N, O, and S; wherein said 5 to 6 membered heteroaryl
10 fused radical is substituted with 0-3 R^{13} ;

additionally, two R^{11} substituents on the same or adjacent carbon atoms may be combined to form a C_3 - C_6 carbocycle substituted with 0-3 R^{13} .

15

[4] In a preferred embodiment the present invention provides for a compound of Formula (I):



20

(I)

or a pharmaceutically acceptable salt or prodrug thereof, wherein:

25 Q is $-NR^1R^2$;

R^1 is H, methyl, ethyl, propyl, butyl, phenyl, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclobutylmethyl,
30 cyclopentylmethyl, or cyclohexylmethyl;

R^2 is H, methyl, ethyl, propyl, butyl, OH, methoxy, ethoxy, propoxy, phenyl, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl,

cyclobutylmethyl, cyclopentylmethyl, or
cyclohexylmethyl;

5 R^3 and R^{3a} are combined to form a 3-8 membered carbocyclic moiety;
 wherein said 3-8 membered carbocyclic moiety is
 saturated or partially unsaturated;
 wherein said 3-8 membered carbocyclic moiety may
 optionally contain a heteroatom selected from -O-,
10 -S-, -S(=O)-, -S(=O)₂-, -N=, -NH-, and -N(R^{20})-, and
 wherein said 3-8 membered carbocyclic moiety is
 substituted with 0-4 R^4 ;

15 additionally, two R^4 substituents on adjacent atoms may be
 combined to form a benzo fused radical; wherein said
 benzo fused radical is substituted with 0-4 R^{23} ;

20 additionally, two R^4 substituents on adjacent atoms may be
 combined to form a 5 to 6 membered heteroaryl fused
 radical, wherein said 5 to 6 membered heteroaryl fused
 radical comprises 1 or 2 heteroatoms selected from N,
 O, and S; wherein said 5 to 6 membered heteroaryl
 fused radical is substituted with 0-3 R^{23} ;

25 additionally, two R^4 substituents on the same or adjacent
 carbon atoms may be combined to form a C₃-C₆
 carbocycle substituted with 0-3 R^{23} ;

30 R^4 , at each occurrence, is independently selected from H,
 OH, Cl, F, Br, I, CN, NO₂, CF₃, acetyl, SCH₃, S(=O)CH₃,
 S(=O)₂CH₃, NR¹⁵R¹⁶, OR^{14a}, C₁-C₄ alkyl, C₂-C₆ alkenyl,
 alkynyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,
 C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-,
 C₃-C₆ carbocycle, aryl, and a
35 5 to 6 membered heterocycle containing 1 to 4
 heteroatoms selected from nitrogen, oxygen, and
 sulphur;

R⁵ is H, C₁-C₆ alkoxy;

C₁-C₆ alkyl substituted with 0-3 R^{5b};

C₂-C₆ alkenyl substituted with 0-3 R^{5b};

C₂-C₆ alkynyl substituted with 0-3 R^{5b};

5 C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};

C₆-C₁₀ aryl substituted with 0-3 R^{5c}; or

5 to 10 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 10 membered heterocycle

10 is substituted with 0-3 R^{5c};

R^{5a} is H, C₁-C₄ alkyl, or C₂-C₄ alkenyl;

R^{5b}, at each occurrence, is independently selected from:

15 H, C₁-C₆ alkyl, CF₃, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂,
NR¹⁵R¹⁶;

C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};

C₆-C₁₀ aryl substituted with 0-3 R^{5c}; or

5 to 10 membered heterocycle containing 1 to 4

20 heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 10 membered heterocycle

is substituted with 0-3 R^{5c};

R^{5c}, at each occurrence, is independently selected from H,

25 OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
S(=O)CH₃, S(=O)₂CH₃,

C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,

C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

30 R⁶ is H;

C₁-C₆ alkyl substituted with 0-3 R^{6a};

C₃-C₁₀ carbocycle substituted with 0-3 R^{6b}; or

C₆-C₁₀ aryl substituted with 0-3 R^{6b};

35 R^{6a}, at each occurrence, is independently selected from H,

C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶,
aryl or CF₃;

R^{6b}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, and C₁-C₄ haloalkoxy;

- 5 Ring B is a 7 membered lactam,
wherein the lactam is saturated, partially saturated
or unsaturated;
wherein each additional lactam carbon is substituted
with 0-2 R¹¹; and,
10 optionally, the lactam contains an additional
heteroatom selected from -O-, -S-, -S(=O)-, -S(=O)₂-
, -N=, -NH-, and -N(R¹⁰)-;

- additionally, two R¹¹ substituents on adjacent atoms may be
15 combined to form a benzo fused radical; wherein said
benzo fused radical is substituted with 0-4 R¹³;

- additionally, two R¹¹ substituents on adjacent atoms may be
combined to form a 5 to 6 membered heteroaryl fused
20 radical, wherein said 5 to 6 membered heteroaryl fused
radical comprises 1 or 2 heteroatoms selected from N,
O, and S; wherein said 5 to 6 membered heteroaryl
fused radical is substituted with 0-3 R¹³;

- 25 additionally, two R¹¹ substituents on the same or adjacent
carbon atoms may be combined to form a C₃-C₆
carbocycle substituted with 0-3 R¹³;

- R¹⁰ is H, C(=O)R¹⁷, C(=O)OR¹⁷, C(=O)NR¹⁸R¹⁹,
30 S(=O)₂NR¹⁸R¹⁹, S(=O)₂R¹⁷;
C₁-C₆ alkyl optionally substituted with 0-3 R^{10a};
C₆-C₁₀ aryl substituted with 0-4 R^{10b};
C₃-C₁₀ carbocycle substituted with 0-3 R^{10b}; or
5 to 10 membered heterocycle containing 1 to 4
35 heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered heterocycle
is substituted with 0-3 R^{10b};

R^{10a}, at each occurrence, is independently selected from H, C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶, CF₃, aryl substituted with 0-4 R^{10b}; or a 5-6 membered heterocycle substituted with 0-4 R^{10b};

5

R^{10b}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

10

R¹¹, at each occurrence, is independently selected from H, C₁-C₄ alkoxy, Cl, F, Br, I, =O, CN, NO₂, NR¹⁸R¹⁹, C(=O)R¹⁷, C(=O)OR¹⁷, C(=O)NR¹⁸R¹⁹, S(=O)₂NR¹⁸R¹⁹, CF₃; C₁-C₆ alkyl optionally substituted with 0-3 R^{11a};

15

C₆-C₁₀ aryl substituted with 0-3 R^{11b};

C₃-C₁₀ carbocycle substituted with 0-3 R^{11b}; or

5 to 10 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 10 membered heterocycle

20

is substituted with 0-3 R^{11b};

R^{11a}, at each occurrence, is independently selected from H, C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶, CF₃;

25

phenyl substituted with 0-3 R^{11b};

C₃-C₆ cycloalkyl substituted with 0-3 R^{11b}; and

5 to 6 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 6 membered heterocycle

30

is substituted with 0-3 R^{11b};

R^{11b}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃,

35

C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,

C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

W is a bond or -(CR⁸R^{8a})_p-;

p is 0, 1, 2, 3, or 4;

5 R^8 and R^{8a} , at each occurrence, are independently selected from H, F, C_1 - C_4 alkyl, C_2 - C_4 alkenyl, C_2 - C_4 alkynyl and C_3 - C_8 cycloalkyl;

X is a bond;
 C_6 - C_{10} aryl substituted with 0-3 R^{Xb} ;
10 C_3 - C_{10} carbocycle substituted with 0-3 R^{Xb} ; or
5 to 10 membered heterocycle substituted with 0-2 R^{Xb} ;

R^{Xb} , at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO_2 , $NR^{15}R^{16}$, CF_3 , acetyl, SCH_3 ,
15 $S(=O)CH_3$, $S(=O)_2CH_3$, C_1 - C_6 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl, C_1 - C_4 haloalkoxy, and C_1 - C_4 halothioalkoxy;

Y is a bond or $-(CR^9R^{9a})_t-V-(CR^9R^{9a})_u-$;

20 t is 0, 1, or 2;

u is 0, 1, or 2;

R^9 and R^{9a} , at each occurrence, are independently selected
25 from H, F, C_1 - C_6 alkyl or C_3 - C_8 cycloalkyl;

V is a bond, $-C(=O)-$, $-O-$, $-S-$, $-S(=O)-$, $-S(=O)_2-$, $-N(R^{19})-$,
30 $-C(=O)NR^{19b}-$, $-NR^{19b}C(=O)-$, $-NR^{19b}S(=O)_2-$, $-S(=O)_2NR^{19b}-$, $-NR^{19b}S(=O)-$, $-S(=O)NR^{19b}-$, $-C(=O)O-$, or $-OC(=O)-$;

Z is H;
 C_1 - C_8 alkyl substituted with 0-3 R^{12a} ;
 C_2 - C_6 alkenyl substituted with 0-3 R^{12a} ;
35 C_2 - C_6 alkynyl substituted with 0-3 R^{12a} ;
 C_6 - C_{10} aryl substituted with 0-4 R^{12b} ;
 C_3 - C_{10} carbocycle substituted with 0-4 R^{12b} ; or

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{12b};

5

R^{12a}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, -C(=O)NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, C₁-C₄ haloalkyl-S-, C₆-C₁₀ aryl substituted with 0-4 R^{12b}; C₃-C₁₀ carbocycle substituted with 0-4 R^{12b}; or 5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{12b};

10

15

R^{12b}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₃-C₆ cycloalkyl, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, C₁-C₄ haloalkyl-S, and aryl substituted with 0-3 R^{12c};

20

R^{12c}, at each occurrence, is independently selected from H, methyl, ethyl, propyl, methoxy, ethoxy, amino, hydroxy, Cl, F, Br, I, CF₃, SCH₃, S(O)CH₃, SO₂CH₃, -N(CH₃)₂, N(CH₃)H, CN, NO₂, OCF₃, C(=O)CH₃, CO₂H, CO₂CH₃, and C₁-C₃ haloalkyl;

30

R¹³, at each occurrence, is independently selected from H, OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, and CF₃;

R¹⁴ is H, phenyl, benzyl, C₁-C₆ alkyl, C₂-C₆ alkoxyalkyl, or C₃-C₆ cycloalkyl;


R^{14a} is H, phenyl, benzyl, or C₁-C₄ alkyl;

- R¹⁵, at each occurrence, is independently selected from H, C₁-C₆ alkyl, benzyl, phenethyl, (C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;
- 5 R¹⁶, at each occurrence, is independently selected from H, C₁-C₆ alkyl, benzyl, phenethyl, (C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;
- 10 R¹⁷ is H, C₁-C₆ alkyl, C₂-C₆ alkoxyalkyl, aryl substituted by 0-4 R^{17a}, or -CH₂-aryl substituted by 0-4 R^{17a};
- 15 R^{17a} is H, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, butoxy, -OH, F, Cl, Br, I, CF₃, OCF₃, SCH₃, S(O)CH₃, SO₂CH₃, -NH₂, -N(CH₃)₂, or C₁-C₄ haloalkyl;
- R¹⁸, at each occurrence, is independently selected from H, C₁-C₆ alkyl, phenyl, benzyl, phenethyl, (C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;
- 20 R¹⁹, at each occurrence, is independently selected from H, OH, C₁-C₆ alkyl, phenyl, benzyl, phenethyl, (C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;
- 25 additionally, R¹⁸ and R¹⁹, when substituents on the same atom, may be combined to form a 3 to 6 membered heterocyclic ring;
- 30 R^{19b}, at each occurrence, is independently is H or C₁-C₄ alkyl;
- R²⁰ is H, C(=O)R¹⁷, C(=O)OR¹⁷, C(=O)NR¹⁸R¹⁹, S(=O)₂NR¹⁸R¹⁹, S(=O)₂R¹⁷;
- 35 C₁-C₆ alkyl optionally substituted with 0-3 R^{20a}; or C₆-C₁₀ aryl substituted with 0-4 R^{20b};

R^{20a} , at each occurrence, is independently selected from H, C_1 - C_4 alkyl, OR^{14} , Cl, F, Br, I, =O, CN, NO_2 , $NR^{15}R^{16}$, CF_3 , or aryl substituted with 0-4 R^{20b} ;

- 5 R^{20b} , at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO_2 , $NR^{15}R^{16}$, CF_3 , acetyl, SCH_3 , $S(=O)CH_3$, $S(=O)_2CH_3$, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl, C_1 - C_4 haloalkoxy, and C_1 - C_4 haloalkyl-S-;
- 10 R^{23} , at each occurrence, is independently selected from H, OH, C_1 - C_6 alkyl, C_1 - C_4 alkoxy, Cl, F, Br, I, CN, NO_2 , $NR^{15}R^{16}$, and CF_3 .

[5] In another preferred embodiment the present
15 invention provides for a compound of Formula (Ia):



20 or a pharmaceutically acceptable salt or prodrug thereof,
wherein:

R^3 and R^{3a} are combined to form a 3-8 membered carbocyclic moiety;

- 25 wherein said 3-8 membered carbocyclic moiety is saturated or partially unsaturated;
wherein said 3-8 membered carbocyclic moiety is substituted with 0-4 R^4 ;

30 additionally, two R^4 substituents on adjacent atoms may be combined to form a benzo fused radical; wherein said benzo fused radical is substituted with 0-4 R^{23} ;

35 additionally, two R^4 substituents on adjacent atoms may be combined to form a 5 to 6 membered heteroaryl fused

radical, wherein said 5 to 6 membered heteroaryl fused radical comprises 1 or 2 heteroatoms selected from N, O, and S; wherein said 5 to 6 membered heteroaryl fused radical is substituted with 0-3 R²³;

5

additionally, two R⁴ substituents on the same or adjacent carbon atoms may be combined to form a C₃-C₆ carbocycle substituted with 0-3 R²³;

10 R⁴, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, NR¹⁵R¹⁶, OR^{14a}, C₁-C₄ alkyl, C₂-C₆ alkenyl, alkynyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-,
15 C₃-C₆ carbocycle, aryl, and a 5 to 6 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur; and

20 R⁵ is H, C₁-C₄ alkoxy; C₁-C₆ alkyl substituted with 0-3 R^{5b}; C₂-C₆ alkenyl substituted with 0-3 R^{5b}; C₂-C₆ alkynyl substituted with 0-3 R^{5b}; C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};
25 C₆-C₁₀ aryl substituted with 0-3 R^{5c}; or 5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{5c};

30


R^{5b}, at each occurrence, is independently selected from: H, C₁-C₆ alkyl, CF₃, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶;
C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};
35 C₆-C₁₀ aryl substituted with 0-3 R^{5c}; or 5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{5c} ;

R^{5c} , at each occurrence, is independently selected from H,
 5 OH, Cl, F, Br, I, CN, NO_2 , $NR^{15}R^{16}$, CF_3 , acetyl, SCH_3 ,
 $S(=O)CH_3$, $S(=O)_2CH_3$,
 C_1 - C_6 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl,
 C_1 - C_4 haloalkoxy, and C_1 - C_4 haloalkyl-S-;

10 R^6 is H, methyl, or ethyl;

Ring B is selected from:



20 R^{10} is H, $C(=O)R^{17}$, $C(=O)OR^{17}$, $C(=O)NR^{18}R^{19}$,
 $S(=O)_2NR^{18}R^{19}$, $S(=O)_2R^{17}$;
 C_1 - C_6 alkyl optionally substituted with 0-3 R^{10a} ;
 C_6 - C_{10} aryl substituted with 0-4 R^{10b} ;
 C_3 - C_{10} carbocycle substituted with 0-3 R^{10b} ; or

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{10b};

5

R^{10a}, at each occurrence, is independently selected from H, C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶, CF₃, aryl substituted with 0-4 R^{10b}; or a 5-6 membered heterocycle substituted with 0-4 R^{10b};

10

R^{10b}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

15

R¹¹, at each occurrence, is independently selected from H, C₁-C₄ alkoxy, Cl, F, Br, I, =O, CN, NO₂, NR¹⁸R¹⁹, C(=O)R¹⁷, C(=O)OR¹⁷, C(=O)NR¹⁸R¹⁹, S(=O)₂NR¹⁸R¹⁹, CF₃; C₁-C₆ alkyl optionally substituted with 0-3 R^{11a};

20

C₆-C₁₀ aryl substituted with 0-3 R^{11b};

C₃-C₁₀ carbocycle substituted with 0-3 R^{11b}; or

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{11b};

25

R^{11a}, at each occurrence, is independently selected from H, C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶, CF₃;

30

phenyl substituted with 0-3 R^{11b};

C₃-C₆ cycloalkyl substituted with 0-3 R^{11b}; and

5 to 6 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R^{11b};

35

R^{11b}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

W is a bond or -(CH₂)_p-;

p is 1 or 2;

X is a bond;

phenyl substituted with 0-2 R^{Xb};
C₃-C₆ carbocycle substituted with 0-2 R^{Xb}; or
5 to 6 membered heterocycle substituted with 0-2 R^{Xb};

R^{Xb}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₄ alkyl, C₁-C₃ alkoxy, C₁-C₃ haloalkyl, C₁-C₃ haloalkoxy, and C₁-C₃ halothioalkoxy;

Y is a bond, -C(=O)-, -O-, -S-, -S(=O)-, -S(=O)₂-, -N(R¹⁹)-, -C(=O)NR^{19b}-, -NR^{19b}C(=O)-, -NR^{19b}S(=O)₂-, -S(=O)₂NR^{19b}-, -NR^{19b}S(=O)-, -S(=O)NR^{19b}-, -C(=O)O-, or -OC(=O)-;

Z is H;

C₁-C₈ alkyl substituted with 0-3 R^{12a};
C₂-C₆ alkenyl substituted with 0-3 R^{12a};
C₂-C₆ alkynyl substituted with 0-3 R^{12a};
C₆-C₁₀ aryl substituted with 0-4 R^{12b};
C₃-C₁₀ carbocycle substituted with 0-4 R^{12b}; or
5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{12b};

R^{12a}, at each occurrence, is independently selected from

H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, -C(=O)NR¹⁵R¹⁶,
CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃,
C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,
C₁-C₄ haloalkoxy, C₁-C₄ haloalkyl-S-,
5 C₆-C₁₀ aryl substituted with 0-4 R^{12b};
C₃-C₁₀ carbocycle substituted with 0-4 R^{12b}; or
5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered heterocycle
10 is substituted with 0-3 R^{12b};

R^{12b}, at each occurrence, is independently selected from
H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl,
SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₃-C₆ cycloalkyl,
15 C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,
C₁-C₄ haloalkoxy, C₁-C₄ haloalkyl-S, and
aryl substituted with 0-3 R^{12c};

R^{12c}, at each occurrence, is independently selected from
20 H, methyl, ethyl, propyl, methoxy, ethoxy, amino,
hydroxy, Cl, F, Br, I, CF₃, SCH₃, S(O)CH₃, SO₂CH₃,
-N(CH₃)₂, N(CH₃)H, CN, NO₂, OCF₃, C(=O)CH₃, CO₂H,
CO₂CH₃, and C₁-C₃ haloalkyl;

25 R¹³, at each occurrence, is independently selected from
H, OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN,
NO₂, NR¹⁵R¹⁶, and CF₃;

R¹⁴ is H, phenyl, benzyl, C₁-C₆ alkyl, C₂-C₆ alkoxyalkyl, or
30 C₃-C₆ cycloalkyl;

R^{14a} is H, phenyl, benzyl, or C₁-C₄ alkyl;

R¹⁵, at each occurrence, is independently selected from H,
35 C₁-C₆ alkyl, benzyl, phenethyl, (C₁-C₆ alkyl)-C(=O)-,
and (C₁-C₆ alkyl)-S(=O)₂-;

R¹⁶, at each occurrence, is independently selected from

H, C₁-C₆ alkyl, benzyl, phenethyl,
(C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;

R¹⁷ is H, C₁-C₆ alkyl, C₂-C₆ alkoxyalkyl,
5 aryl substituted by 0-4 R^{17a}, or
-CH₂-aryl substituted by 0-4 R^{17a};

R^{17a} is H, methyl, ethyl, propyl, butyl, methoxy, ethoxy,
propoxy, butoxy, -OH, F, Cl, Br, I, CF₃, OCF₃, SCH₃,
10 S(O)CH₃, SO₂CH₃, -NH₂, -N(CH₃)₂, or C₁-C₄ haloalkyl;

R¹⁸, at each occurrence, is independently selected from
H, C₁-C₆ alkyl, phenyl, benzyl, phenethyl,
(C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;
15


R¹⁹, at each occurrence, is independently selected from
H, OH, methyl, ethyl, propyl, butyl, phenyl, benzyl,
phenethyl;

20 additionally, R¹⁸ and R¹⁹, when substituents on the same
atom, may be combined to form a 3 to 6 membered
heterocyclic ring selected from pyrrolyl, imidazolyl,
imidazolidinyl, pyrrolidinyl, piperidinyl,
piperazinyl, and morpholinyl;

25 R^{19b}, at each occurrence, is independently is H or C₁-C₄
alkyl; and

R²³, at each occurrence, is independently selected from
30 H, OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN,
NO₂, NR¹⁵R¹⁶, and CF₃.

[6] In another preferred embodiment the present
invention provides for a compound of Formula (Ib):
35



or a pharmaceutically acceptable salt or prodrug thereof,
 5 wherein:

R³ and R^{3a} are combined to form a 3-8 membered carbocyclic moiety;

wherein said 3-8 membered carbocyclic moiety is

10 saturated or partially unsaturated;

wherein said 3-8 membered carbocyclic moiety is
 substituted with 0-4 R⁴;

additionally, two R⁴ substituents on adjacent atoms may be
 15 combined to form a benzo fused radical; wherein said
 benzo fused radical is substituted with 0-4 R²³;

additionally, two R⁴ substituents on adjacent atoms may be
 combined to form a 5 to 6 membered heteroaryl fused
 20 radical, wherein said 5 to 6 membered heteroaryl fused
 radical comprises 1 or 2 heteroatoms selected from N,
 O, and S; wherein said 5 to 6 membered heteroaryl
 fused radical is substituted with 0-3 R²³;

25 additionally, two R⁴ substituents on the same or adjacent
 carbon atoms may be combined to form a C₃-C₆
 carbocycle substituted with 0-3 R²³;

R⁴, at each occurrence, is independently selected from H,
 30 OH, Cl, F, Br, I, CN, NO₂, CF₃, acetyl, SCH₃, S(=O)CH₃,
 S(=O)₂CH₃, NR¹⁵R¹⁶, OR^{14a}, C₁-C₄ alkyl, C₂-C₆ alkenyl,
 alkynyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,
 C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-,
 C₃-C₆ carbocycle, aryl, and a

5 to 6 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur; and

5 R^5 is H;

C_1 - C_6 alkyl substituted with 0-3 R^{5b} ;

C_2 - C_6 alkenyl substituted with 0-3 R^{5b} ;

C_2 - C_6 alkynyl substituted with 0-3 R^{5b} ;

C_3 - C_{10} carbocycle substituted with 0-3 R^{5c} ;

10 C_6 - C_{10} aryl substituted with 0-3 R^{5c} ; or

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{5c} ;

15

R^{5b} , at each occurrence, is independently selected from:

H, C_1 - C_6 alkyl, CF_3 , OR^{14} , Cl, F, Br, I, =O, CN, NO_2 , $NR^{15}R^{16}$;

C_3 - C_{10} carbocycle substituted with 0-3 R^{5c} ;

20 C_6 - C_{10} aryl substituted with 0-3 R^{5c} ; or

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{5c} ;

25


R^{5c} , at each occurrence, is independently selected from H,


OH, Cl, F, Br, I, CN, NO_2 , $NR^{15}R^{16}$, CF_3 , acetyl, SCH_3 , $S(=O)CH_3$, $S(=O)_2CH_3$,

C_1 - C_6 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl,

30 C_1 - C_4 haloalkoxy, and C_1 - C_4 haloalkyl-S-;

Ring B is selected from:





- 5 R¹¹, at each occurrence, is independently selected from
 H, C₁-C₄ alkoxy, Cl, F, Br, I, =O, CN, NO₂, NR¹⁸R¹⁹,
 C(=O)R¹⁷, C(=O)OR¹⁷, C(=O)NR¹⁸R¹⁹, S(=O)₂NR¹⁸R¹⁹, CF₃;
 C₁-C₆ alkyl optionally substituted with 0-3 R^{11a};
 C₆-C₁₀ aryl substituted with 0-3 R^{11b};
 C₃-C₁₀ carbocycle substituted with 0-3 R^{11b}; or
 10 5 to 10 membered heterocycle containing 1 to 4
 heteroatoms selected from nitrogen, oxygen, and
 sulphur, wherein said 5 to 10 membered heterocycle
 is substituted with 0-3 R^{11b};
- 15 R^{11a}, at each occurrence, is independently selected from
 H, C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂,
 NR¹⁵R¹⁶, CF₃;
 phenyl substituted with 0-3 R^{11b};
 C₃-C₆ cycloalkyl substituted with 0-3 R^{11b}; and
 20 5 to 6 membered heterocycle containing 1 to 4
 heteroatoms selected from nitrogen, oxygen, and
 sulphur, wherein said 5 to 6 membered heterocycle
 is substituted with 0-3 R^{11b};
- 25 R^{11b}, at each occurrence, is independently selected from H,
 OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
 S(=O)CH₃, S(=O)₂CH₃,
 C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,
 C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;
 30
- W is a bond;
 X is a bond;
 Y is a bond;

Z is H;

C₁-C₈ alkyl substituted with 0-3 R^{12a};

C₂-C₆ alkenyl substituted with 0-3 R^{12a};

C₂-C₆ alkynyl substituted with 0-3 R^{12a};

5 C₆-C₁₀ aryl substituted with 0-4 R^{12b};

C₃-C₁₀ carbocycle substituted with 0-4 R^{12b}; or

5 to 10 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 10 membered heterocycle

10 is substituted with 0-3 R^{12b};

R^{12a}, at each occurrence, is independently selected from

H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, -C(=O)NR¹⁵R¹⁶,

CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃,

15 C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,

C₁-C₄ haloalkoxy, C₁-C₄ haloalkyl-S-,

C₆-C₁₀ aryl substituted with 0-4 R^{12b};

C₃-C₁₀ carbocycle substituted with 0-4 R^{12b}; or

5 to 10 membered heterocycle containing 1 to 4

20 heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 10 membered heterocycle

is substituted with 0-3 R^{12b};

R^{12b}, at each occurrence, is independently selected from

25 H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl,

SCH₃, S(=O)CH₃, S(=O)₂CH₃,

C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,

C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

30 R¹³, at each occurrence, is independently selected from

H, OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN,

NO₂, NR¹⁵R¹⁶, and CF₃;

R¹⁴ is H, phenyl, benzyl, C₁-C₆ alkyl, C₂-C₆ alkoxyalkyl, or

35 C₃-C₆ cycloalkyl;

R^{14a} is H, phenyl, benzyl, or C₁-C₄ alkyl;

- R¹⁵, at each occurrence, is independently selected from H, C₁-C₆ alkyl, benzyl, phenethyl, (C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;
- 5 R¹⁶, at each occurrence, is independently selected from H, C₁-C₆ alkyl, benzyl, phenethyl, (C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;
- 10 R¹⁷ is H, C₁-C₆ alkyl, C₂-C₆ alkoxyalkyl, aryl substituted by 0-4 R^{17a}, or -CH₂-aryl substituted by 0-4 R^{17a};
- 15 R^{17a} is H, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, butoxy, -OH, F, Cl, Br, I, CF₃, OCF₃, SCH₃, S(O)CH₃, SO₂CH₃, -NH₂, -N(CH₃)₂, or C₁-C₄ haloalkyl;
- 20 R¹⁸, at each occurrence, is independently selected from H, C₁-C₆ alkyl, phenyl, benzyl, phenethyl, (C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;
- R¹⁹, at each occurrence, is independently selected from H, OH, methyl, ethyl, propyl, butyl, phenyl, benzyl, phenethyl; and
- 25 R²³, at each occurrence, is independently selected from H, OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, and CF₃.

- [7] In another preferred embodiment the present invention provides for a compound of Formula (Ib):
- 30 R³ and R^{3a} are combined to form a 3-8 membered carbocyclic moiety;
- wherein said 3-8 membered carbocyclic moiety is saturated or partially unsaturated;
- 35 wherein said 3-8 membered carbocyclic moiety is substituted with 0-3 R⁴;

R⁴, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, CF₃, acetyl, SCH₃, S(=O)CH₃,
S(=O)₂CH₃, NR¹⁵R¹⁶, OR^{14a}, C₁-C₄ alkyl, C₂-C₄ alkenyl,
C₂-C₄ alkynyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,
5 C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

R⁵ is H;

C₁-C₆ alkyl substituted with 0-3 R^{5b};
C₂-C₆ alkenyl substituted with 0-3 R^{5b};
10 C₂-C₆ alkynyl substituted with 0-3 R^{5b};
C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};
C₆-C₁₀ aryl substituted with 0-3 R^{5c}; or
5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
15 sulphur, wherein said 5 to 10 membered heterocycle
is substituted with 0-3 R^{5c};


R^{5b}, at each occurrence, is independently selected from:

H, C₁-C₆ alkyl, CF₃, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂,
20 NR¹⁵R¹⁶;
C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};
C₆-C₁₀ aryl substituted with 0-3 R^{5c}; or
5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
25 sulphur, wherein said 5 to 10 membered heterocycle
is substituted with 0-3 R^{5c};

R^{5c}, at each occurrence, is independently selected from H,

OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
30 S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄
haloalkyl, and C₁-C₄ haloalkoxy;

Ring B is selected from:



- 5 R¹¹, at each occurrence, is independently selected from
H, =O, NR¹⁸R¹⁹, CF₃;
C₁-C₄ alkyl optionally substituted with 0-1 R^{11a};
phenyl substituted with 0-3 R^{11b};
C₃-C₇ carbocycle substituted with 0-3 R^{11b}; and
- 10 5 to 7 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 7 membered heterocycle is
substituted with 0-3 R^{11b}; wherein said 5 to 7
membered heterocycle is selected from pyridinyl,
15 pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,
pyrrolyl, piperazinyl, piperidinyl, homopiperidinyl,
pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and
tetrazolyl;
- 20 R^{11a}, at each occurrence, is independently selected from H,
C₁-C₄ alkyl, OR¹⁴, F, Cl, =O, NR¹⁵R¹⁶, CF₃, or phenyl
substituted with 0-3 R^{11b};
- 25 R^{11b}, at each occurrence, is independently selected from H,
OH, Cl, F, NR¹⁵R¹⁶, CF₃, methyl, ethyl, propyl, butyl,
methoxy, ethoxy, propoxy, C₁-C₂ haloalkyl, and C₁-C₂
haloalkoxy;

W is a bond;

30 X is a bond;

Y is a bond;

Z is H;

- C₁-C₈ alkyl substituted with 0-3 R^{12a};
5 C₂-C₆ alkenyl substituted with 0-3 R^{12a}; or
C₂-C₆ alkynyl substituted with 0-3 R^{12a};

- R^{12a}, at each occurrence, is independently selected from
H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, -C(=O)NR¹⁵R¹⁶,
10 CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃,
C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,
C₁-C₄ haloalkoxy, C₁-C₄ haloalkyl-S-,
C₆-C₁₀ aryl substituted with 0-4 R^{12b};
C₃-C₁₀ carbocycle substituted with 0-4 R^{12b}; or
15 5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered heterocycle
is substituted with 0-3 R^{12b}; and wherein said 5 to
10 membered heterocycle is selected from pyridinyl,
20 pyrimidinyl, triazinyl, furanyl, thienyl,
thiazolyl, pyrrolyl, pyrazolyl, imidazolyl,
oxazolyl, isoxazolyl, tetrazolyl, benzofuranyl,
benzothiofuranyl, indolyl, benzimidazolyl,
1H-indazolyl, oxazolidinyl, isoxazolidinyl,
25 benzotriazolyl, benzisoxazolyl, oxindolyl,
benzoxazolinyl, quinolinyl, and isoquinolinyl;

- R^{12b}, at each occurrence, is independently selected from
H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl,
30 SCH₃, S(=O)CH₃, S(=O)₂CH₃,
C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,
C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

- R¹³, at each occurrence, is independently selected from
35 H, OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN,
NO₂, NR¹⁵R¹⁶, and CF₃;

R¹⁴ is H, phenyl, benzyl, C₁-C₆ alkyl, C₂-C₆ alkoxyalkyl, or C₃-C₆ cycloalkyl;

R^{14a} is H, phenyl, benzyl, or C₁-C₄ alkyl;

5

R¹⁵, at each occurrence, is independently selected from H, C₁-C₆ alkyl, benzyl, phenethyl, (C₁-C₄ alkyl)-C(=O)-, and (C₁-C₄ alkyl)-S(=O)₂-;

10 R¹⁶, at each occurrence, is independently selected from H, C₁-C₆ alkyl, benzyl, phenethyl, (C₁-C₄ alkyl)-C(=O)-, and (C₁-C₄ alkyl)-S(=O)₂-;

15 R¹⁸, at each occurrence, is independently selected from H, C₁-C₆ alkyl, phenyl, benzyl, phenethyl, (C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-; and

20 R¹⁹, at each occurrence, is independently selected from H, OH, methyl, ethyl, propyl, butyl, phenyl, benzyl, and phenethyl.

[8] In another preferred embodiment the present invention provides for a compound of Formula (Ib):

25 R³ and R^{3a} are combined to form a 3-6 membered carbocyclic moiety;
wherein said 3-6 membered carbocyclic moiety is saturated or partially unsaturated;
wherein said 3-6 membered carbocyclic moiety is
30 substituted with 0-2 R⁴;

R⁴, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, CF₃, acetyl, SCH₃, methyl, ethyl, methoxy, ethoxy, allyl, -OCF₃, and -SCF₃;

35

R⁵ is H;
C₁-C₄ alkyl substituted with 0-3 R^{5b};
C₂-C₄ alkenyl substituted with 0-3 R^{5b}; or

C₂-C₄ alkynyl substituted with 0-3 R^{5b};

R^{5b}, at each occurrence, is independently selected from:

H, methyl, ethyl, propyl, butyl, CF₃, OR¹⁴, Cl, F, Br,

5 I, =O;

C₃-C₆ carbocycle substituted with 0-3 R^{5c};

phenyl substituted with 0-3 R^{5c}; or

5 to 6 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and


10 sulphur, wherein said 5 to 6 membered heterocycle
is substituted with 0-3 R^{5c};

R^{5c}, at each occurrence, is independently selected from H,

OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,

15 S(=O)CH₃, S(=O)₂CH₃, C₁-C₄ alkyl, C₁-C₃ alkoxy, C₁-C₂
haloalkyl, and C₁-C₂ haloalkoxy;

Ring B is selected from:



R¹¹, at each occurrence, is independently selected from

25 H, =O, NR¹⁸R¹⁹, CF₃;

C₁-C₄ alkyl optionally substituted with 0-1 R^{11a};

phenyl substituted with 0-3 R^{11b};

C₃-C₇ carbocycle substituted with 0-3 R^{11b}; and

5 to 7 membered heterocycle containing 1 to 4

30 heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 7 membered heterocycle is

substituted with 0-3 R^{11b}; wherein said 5 to 7
membered heterocycle is selected from pyridinyl,
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,
pyrrolyl, piperazinyl, piperidinyl, homopiperidinyl,
5 pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and
tetrazolyl;

R^{11a}, at each occurrence, is independently selected from H,
methyl, ethyl, propyl, butyl, methoxy, ethoxy,
10 propoxy, phenoxy, F, Cl, =O, NR¹⁵R¹⁶, CF₃, or phenyl
substituted with 0-3 R^{11b};

R^{11b}, at each occurrence, is independently selected from H,
OH, Cl, F, NR¹⁵R¹⁶, CF₃, methyl, ethyl, propyl, butyl,
15 methoxy, ethoxy, propoxy, C₁-C₂ haloalkyl, and C₁-C₂
haloalkoxy;

W is a bond;
X is a bond;
20 Y is a bond;

Z is H;
C₁-C₄ alkyl substituted with 0-3 R^{12a};
C₂-C₄ alkenyl substituted with 0-3 R^{12a}; or
25 C₂-C₄ alkynyl substituted with 0-3 R^{12a};

R^{12a}, at each occurrence, is independently selected from
H, OH, Cl, F, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃,
S(=O)₂CH₃, C₁-C₄ alkyl, C₁-C₃ alkoxy, C₁-C₂ haloalkyl,
30 and C₁-C₂ haloalkoxy;

R¹³, at each occurrence, is independently selected from
H, OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN,
NO₂, NR¹⁵R¹⁶, and CF₃;

35 R¹⁴ is H, phenyl, benzyl, C₁-C₄ alkyl, or C₂-C₄ alkoxyalkyl;

R¹⁵, at each occurrence, is independently selected from H, C₁-C₄ alkyl, and benzyl;

5 R¹⁶, at each occurrence, is independently selected from H, methyl, ethyl, propyl, butyl, benzyl, phenethyl, methyl-C(=O)-, ethyl-C(=O)-, methyl-S(=O)₂-, and ethyl-S(=O)₂-;

10 R¹⁸, at each occurrence, is independently selected from H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and phenethyl; and

15 R¹⁹, at each occurrence, is independently selected from H, methyl, ethyl, propyl, and butyl.

[9] In another preferred embodiment the present invention provides for a compound of Formula (Ib):

20 R³ and R^{3a} are combined to form a 3-6 membered carbocyclic moiety selected from cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, and cyclohexenyl; wherein said 3-6 membered carbocyclic moiety is substituted with 0-1 R⁴;

25 R⁴ is selected from H, OH, Cl, F, CN, CF₃, methyl, ethyl, methoxy, ethoxy, allyl, and -OCF₃;


30 R⁵ is C₁-C₄ alkyl substituted with 0-1 R^{5b};
C₂-C₄ alkenyl substituted with 0-1 R^{5b};
C₂-C₄ alkynyl substituted with 0-1 R^{5b};

35 R^{5b}, at each occurrence, is independently selected from:
H, methyl, ethyl, propyl, butyl, CF₃, OR¹⁴, =O;
C₃-C₆ carbocycle substituted with 0-2 R^{5c};
phenyl substituted with 0-3 R^{5c}; or
5 to 6 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle

is substituted with 0-3 R^{5c} ; wherein said 5 to 6 membered heterocycle is selected from pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, pyrrolyl, piperazinyl, piperidinyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

R^{5c} , at each occurrence, is independently selected from H, OH, Cl, F, $NR^{15}R^{16}$, CF_3 , acetyl, SCH_3 , $S(=O)CH_3$, $S(=O)_2CH_3$, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, C_1 - C_2 haloalkyl, and C_1 - C_2 haloalkoxy;

Ring B is selected from:



R^{11} , at each occurrence, is independently selected from H, =O, $NR^{18}R^{19}$; C_1 - C_4 alkyl optionally substituted with 0-1 R^{11a} ; phenyl substituted with 0-3 R^{11b} ; cyclohexyl substituted with 0-3 R^{11b} ; cycloheptyl substituted with 0-3 R^{11b} ; 5 to 7 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 7 membered heterocycle is substituted with 0-3 R^{11b} ; wherein said 5 to 7 membered heterocycle is selected from pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, pyrrolyl, piperazinyl, piperidinyl, homopiperidinyl,

pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and
tetrazolyl;

5 R^{11a}, at each occurrence, is independently selected from H,
methyl, ethyl, propyl, methoxy, ethoxy, propoxy,
phenoxy, F, Cl, =O, NR¹⁵R¹⁶, CF₃, or phenyl substituted
with 0-3 R^{11b};

10 R^{11b}, at each occurrence, is independently selected from H,
OH, Cl, F, NR¹⁵R¹⁶, CF₃, methyl, ethyl, propyl, butyl,
methoxy, ethoxy, propoxy, C₁-C₂ haloalkyl, and C₁-C₂
haloalkoxy;

W is a bond;

15 X is a bond;

Y is a bond;

Z is H;

20 C₁-C₄ alkyl substituted with 0-1 R^{12a};
C₂-C₄ alkenyl substituted with 0-1 R^{12a}; or
C₂-C₄ alkynyl substituted with 0-1 R^{12a};

25 R^{12a}, at each occurrence, is independently selected from
H, OH, Cl, F, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃,
S(=O)₂CH₃, methyl, ethyl, propyl, butyl, methoxy,
ethoxy, propoxy, C₁-C₂ haloalkyl, and C₁-C₂ haloalkoxy;

30 R¹³, at each occurrence, is independently selected from
H, OH, methyl, ethyl, propyl, butyl, methoxy, ethoxy,
Cl, F, Br, CN, NR¹⁵R¹⁶, and CF₃;

R¹⁴ is H, phenyl, benzyl, methyl, ethyl, propyl, or butyl;

35 R¹⁵, at each occurrence, is independently selected from H,
methyl, ethyl, propyl, and butyl;

R¹⁶, at each occurrence, is independently selected from

H, methyl, ethyl, propyl, butyl, benzyl, and phenethyl;

5 R^{18} , at each occurrence, is independently selected from H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and phenethyl; and

R^{19} , at each occurrence, is independently selected from H, methyl, ethyl, propyl, and butyl.

10

[10] In another preferred embodiment the present invention provides for a compound of Formula (Ib):

15 R^3 and R^{3a} are combined to form cyclobutyl, cyclopentyl, cyclopentenyl, or cyclohexyl;

R^5 is $-\text{CH}_3$, $-\text{CH}_2\text{CH}_3$, $-\text{CH}_2\text{CH}_2\text{CH}_3$, $-\text{CH}(\text{CH}_3)_2$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$,
 $-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3$, $-\text{CH}_2\text{CH}(\text{CH}_3)_2$, $-\text{CH}_2\text{C}(\text{CH}_3)_3$,
 $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $-\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_2\text{CH}_3$, $-\text{CH}_2\text{CH}(\text{CH}_3)\text{CH}_2\text{CH}_3$,
20 $-\text{CH}_2\text{CH}_2\text{CH}(\text{CH}_3)_2$, $-\text{CH}(\text{CH}_2\text{CH}_3)_2$, $-\text{CF}_3$, $-\text{CH}_2\text{CF}_3$, $-\text{CH}_2\text{CH}_2\text{CF}_3$,
 $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CF}_3$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2\text{CF}_3$, $-\text{CH}=\text{CH}_2$, $-\text{CH}_2\text{CH}=\text{CH}_2$,
 $-\text{CH}=\text{CHCH}_3$, $\text{cis-CH}_2\text{CH}=\text{CH}(\text{CH}_3)$, $\text{trans-CH}_2\text{CH}=\text{CH}(\text{CH}_3)$,
 $\text{trans-CH}_2\text{CH}=\text{CH}(\text{C}_6\text{H}_5)$, $-\text{CH}_2\text{CH}=\text{C}(\text{CH}_3)_2$, $\text{cis-CH}_2\text{CH}=\text{CHCH}_2\text{CH}_3$,
 $\text{trans-CH}_2\text{CH}=\text{CHCH}_2\text{CH}_3$, $\text{cis-CH}_2\text{CH}_2\text{CH}=\text{CH}(\text{CH}_3)$,
25 $\text{trans-CH}_2\text{CH}_2\text{CH}=\text{CH}(\text{CH}_3)$, $\text{trans-CH}_2\text{CH}=\text{CHCH}_2(\text{C}_6\text{H}_5)$,
 $-\text{C}\equiv\text{CH}$, $-\text{CH}_2\text{C}\equiv\text{CH}$, $-\text{CH}_2\text{C}\equiv\text{C}(\text{CH}_3)$, $-\text{CH}_2\text{C}\equiv\text{C}(\text{C}_6\text{H}_5)$,
 $-\text{CH}_2\text{CH}_2\text{C}\equiv\text{CH}$, $-\text{CH}_2\text{CH}_2\text{C}\equiv\text{C}(\text{CH}_3)$, $-\text{CH}_2\text{CH}_2\text{C}\equiv\text{C}(\text{C}_6\text{H}_5)$,
 $-\text{CH}_2\text{CH}_2\text{CH}_2\text{C}\equiv\text{CH}$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{C}\equiv\text{C}(\text{CH}_3)$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{C}\equiv\text{C}(\text{C}_6\text{H}_5)$,
cyclopropyl- CH_2- , cyclobutyl- CH_2- , cyclopentyl- CH_2- ,
30 cyclohexyl- CH_2- , (2- CH_3 -cyclopropyl) CH_2- ,
(3- CH_3 -cyclobutyl) CH_2- ,
cyclopropyl- CH_2CH_2- , cyclobutyl- CH_2CH_2- ,
cyclopentyl- CH_2CH_2- , cyclohexyl- CH_2CH_2- ,
(2- CH_3 -cyclopropyl) CH_2CH_2- , (3- CH_3 -cyclobutyl) CH_2CH_2- ,
35 phenyl- CH_2- , (2-F-phenyl) CH_2- , (3-F-phenyl) CH_2- ,
(4-F-phenyl) CH_2- , furanyl- CH_2- , thienyl- CH_2- ,
pyridyl- CH_2- , 1-imidazolyl- CH_2- , oxazolyl- CH_2- ,
isoxazolyl- CH_2- , phenyl- CH_2CH_2- , (2-F-phenyl) CH_2CH_2- ,

(3-F-phenyl)CH₂CH₂-, (4-F-phenyl)CH₂CH₂-,
furanyl-CH₂CH₂-, thienyl-CH₂CH₂-, pyridyl-CH₂CH₂-,
1-imidazolyl-CH₂CH₂-, oxazolyl-CH₂CH₂-,
isoxazolyl-CH₂CH₂-,

5

W is a bond;

X is a bond;

Y is a bond;

10 Z is methyl, ethyl, i-propyl, n-propyl, n-butyl, i-butyl,
s-butyl, t-butyl, or allyl;

R¹¹, at each occurrence, is independently selected from

15 H, =O, methyl, ethyl, phenyl, benzyl, phenethyl,
4-F-phenyl, (4-F-phenyl)CH₂-, (4-F-phenyl)CH₂CH₂-,
3-F-phenyl, (3-F-phenyl)CH₂-, (3-F-phenyl)CH₂CH₂-,
2-F-phenyl, (2-F-phenyl)CH₂-, (2-F-phenyl)CH₂CH₂-,
4-Cl-phenyl, (4-Cl-phenyl)CH₂-, (4-Cl-phenyl)CH₂CH₂-,
3-Cl-phenyl, (3-Cl-phenyl)CH₂-, (3-Cl-phenyl)CH₂CH₂-,
20 4-CH₃-phenyl, (4-CH₃-phenyl)CH₂-, (4-CH₃-phenyl)CH₂CH₂-,
3-CH₃-phenyl, (3-CH₃-phenyl)CH₂-, (3-CH₃-phenyl)CH₂CH₂-,
4-CF₃-phenyl, (4-CF₃-phenyl)CH₂-, (4-CF₃-phenyl)CH₂CH₂-,
pyrid-2-yl, pyrid-3-yl, pyrid-4-yl, cyclohexyl,
cycloheptyl, piperidinyl, or homopiperidinyl; and

25

R¹³, at each occurrence, is independently selected from
H, F, Cl, OH, -CH₃, -CH₂CH₃, -OCH₃, or -CF₃.

[11] In another preferred embodiment the present
30 invention provides for a compound of Formula (Ib):

R³ and R^{3a} are combined to form cyclobutyl, cyclopentyl,
cyclopentenyl, or cyclohexyl;

35 R⁵ is -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH(CH₃)₂, -CH₂CH₂CH₂CH₃,
-CH(CH₃)CH₂CH₃, -CH₂CH(CH₃)₂, -CH₂C(CH₃)₃,
-CH₂CH₂CH₂CH₂CH₃, -CH(CH₃)CH₂CH₂CH₃, -CH₂CH(CH₃)CH₂CH₃,

-CH₂CH₂CH(CH₃)₂, -CH(CH₂CH₃)₂, cyclopropyl-CH₂-,
cyclobutyl-CH₂-, (2-CH₃-cyclopropyl)CH₂-, or
(3-CH₃-cyclobutyl)CH₂-;

5 W is a bond;

X is a bond;

Y is a bond;


10 Z is methyl, ethyl, i-propyl, n-propyl, n-butyl, i-butyl,
s-butyl, t-butyl, or allyl;

R¹¹, at each occurrence, is independently selected from
H, =O, methyl, ethyl, phenyl, benzyl, phenethyl,
4-F-phenyl, (4-F-phenyl)CH₂-, (4-F-phenyl)CH₂CH₂-,
15 3-F-phenyl, (3-F-phenyl)CH₂-, (3-F-phenyl)CH₂CH₂-,
2-F-phenyl, (2-F-phenyl)CH₂-, (2-F-phenyl)CH₂CH₂-,
4-Cl-phenyl, (4-Cl-phenyl)CH₂-, (4-Cl-phenyl)CH₂CH₂-,
3-Cl-phenyl, (3-Cl-phenyl)CH₂-, (3-Cl-phenyl)CH₂CH₂-,
4-CH₃-phenyl, (4-CH₃-phenyl)CH₂-, (4-CH₃-phenyl)CH₂CH₂-,
20 3-CH₃-phenyl, (3-CH₃-phenyl)CH₂-, (3-CH₃-phenyl)CH₂CH₂-,
4-CF₃-phenyl, (4-CF₃-phenyl)CH₂-, (4-CF₃-phenyl)CH₂CH₂-,
pyrid-2-yl, pyrid-3-yl, pyrid-4-yl, cyclohexyl,
cycloheptyl, piperidinyl, or homopiperidinyl; and

25 R¹³, at each occurrence, is independently selected from
H, F, Cl, OH, -CH₃, -CH₂CH₃, -OCH₃, or -CF₃.

[12] In another preferred embodiment the present
invention provides for a compound of Formula (Ib):

30



or a pharmaceutically acceptable salt or prodrug thereof,
35 wherein:

R³ and R^{3a} are combined to form a 3-8 membered carbocyclic moiety;

5 wherein said 3-8 membered carbocyclic moiety is saturated or partially unsaturated;
wherein said 3-8 membered carbocyclic moiety is substituted with 0-3 R⁴;

10 R⁴, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, NR¹⁵R¹⁶, OR^{14a}, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

15 R⁵ is H;
C₁-C₆ alkyl substituted with 0-3 R^{5b};
C₂-C₆ alkenyl substituted with 0-3 R^{5b};
C₂-C₆ alkynyl substituted with 0-3 R^{5b};
C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};
20 C₆-C₁₀ aryl substituted with 0-3 R^{5c}; or
5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{5c};


25 R^{5b}, at each occurrence, is independently selected from:
H, C₁-C₆ alkyl, CF₃, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶;
C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};
30 C₆-C₁₀ aryl substituted with 0-3 R^{5c}; or
5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{5c};

35 R^{5c}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,

S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, and C₁-C₄ haloalkoxy;

Ring B is selected from:

5



- 10 R¹¹, at each occurrence, is independently selected from
H, =O, NR¹⁸R¹⁹, CF₃;
C₁-C₄ alkyl optionally substituted with 0-1 R^{11a};
phenyl substituted with 0-3 R^{11b};
C₃-C₇ carbocycle substituted with 0-3 R^{11b}; and
15 5 to 7 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 7 membered heterocycle is
substituted with 0-3 R^{11b}; wherein said 5 to 7
membered heterocycle is selected from pyridinyl,
20 pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,
pyrrolyl, piperazinyl, piperidinyl, homopiperidinyl,
pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and
tetrazolyl;

- 25 R^{11a}, at each occurrence, is independently selected from H,
C₁-C₄ alkyl, OR¹⁴, F, Cl, =O, NR¹⁵R¹⁶, CF₃, or phenyl
substituted with 0-3 R^{11b};

- 30 R^{11b}, at each occurrence, is independently selected from H,
OH, Cl, F, NR¹⁵R¹⁶, CF₃, methyl, ethyl, propyl, butyl,

methoxy, ethoxy, propoxy, C₁-C₂ haloalkyl, and C₁-C₂ haloalkoxy;

W is a bond, -CH₂-, -CH₂CH₂-;

5

X is a bond;

phenyl substituted with 0-2 R^{Xb};

C₃-C₆ cycloalkyl substituted with 0-2 R^{Xb}; or

5 to 6 membered heterocycle substituted with 0-2 R^{Xb};

10

R^{Xb}, at each occurrence, is independently selected from H, OH, Cl, F, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₄ alkyl, C₁-C₃ alkoxy, C₁-C₂ haloalkyl, and C₁-C₂ haloalkoxy;

15

Y is a bond, -C(=O)-, -O-, -S-, -S(=O)-, -S(=O)₂-, -N(R¹⁹)-, -C(=O)NR^{19b}-, -NR^{19b}C(=O)-, -NR^{19b}S(=O)₂-, -S(=O)₂NR^{19b}-, -NR^{19b}S(=O)-, -S(=O)NR^{19b}-, -C(=O)O-, or -OC(=O)-;

20

Z is C₁-C₃ alkyl substituted with 1-2 R^{12a};

C₆-C₁₀ aryl substituted with 0-4 R^{12b};

C₃-C₁₀ carbocycle substituted with 0-3 R^{12b}; or

5 to 10 membered heterocycle containing 1 to 4

25

heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{12b};

30

R^{12a}, at each occurrence, is independently selected from

C₆-C₁₀ aryl substituted with 0-4 R^{12b};

C₃-C₁₀ carbocycle substituted with 0-4 R^{12b}; and

5 to 10 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle

35

is substituted with 0-3 R^{12b};

R^{12b}, at each occurrence, is independently selected from

H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

5

R¹³, at each occurrence, is independently selected from H, OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, and CF₃;

10 R¹⁴ is H, phenyl, benzyl, C₁-C₆ alkyl, C₂-C₆ alkoxyalkyl, or C₃-C₆ cycloalkyl;

R^{14a} is H, phenyl, benzyl, or C₁-C₄ alkyl;

15 R¹⁵, at each occurrence, is independently selected from H, C₁-C₆ alkyl, benzyl, phenethyl, (C₁-C₄ alkyl)-C(=O)-, and (C₁-C₄ alkyl)-S(=O)₂-;

20 R¹⁶, at each occurrence, is independently selected from H, OH, C₁-C₆ alkyl, benzyl, phenethyl, (C₁-C₄ alkyl)-C(=O)-, and (C₁-C₄ alkyl)-S(=O)₂-;

25 R¹⁸, at each occurrence, is independently selected from H, C₁-C₆ alkyl, phenyl, benzyl, phenethyl, (C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;

R¹⁹, at each occurrence, is independently selected from H, OH, methyl, ethyl, propyl, butyl, phenyl, benzyl, and phenethyl; and

30

R^{19b}, at each occurrence, is independently selected from H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and phenethyl.

35 [13] In another preferred embodiment the present invention provides for a compound of Formula (Ib) wherein:

R^3 and R^{3a} are combined to form a 3-6 membered carbocyclic moiety;

wherein said 3-6 membered carbocyclic moiety is saturated or partially unsaturated;

5 wherein said 3-6 membered carbocyclic moiety is substituted with 0-2 R^4 ;

10 R^4 , at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO_2 , CF_3 , acetyl, SCH_3 , methyl, ethyl, methoxy, ethoxy, allyl, $-OCF_3$, and $-SCF_3$;

R^5 is H;

C_1 - C_4 alkyl substituted with 0-3 R^{5b} ;

C_2 - C_4 alkenyl substituted with 0-3 R^{5b} ; or

15 C_2 - C_4 alkynyl substituted with 0-3 R^{5b} ;

R^{5b} , at each occurrence, is independently selected from:

H, methyl, ethyl, propyl, butyl, CF_3 , OR^{14} , Cl, F, Br, I, $=O$;

20 C_3 - C_6 carbocycle substituted with 0-3 R^{5c} ;

phenyl substituted with 0-3 R^{5c} ; or

5 to 6 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 6 membered heterocycle

25 is substituted with 0-3 R^{5c} ;


R^{5c} , at each occurrence, is independently selected from H,


OH, Cl, F, Br, I, CN, NO_2 , $NR^{15}R^{16}$, CF_3 , acetyl, SCH_3 ,

$S(=O)CH_3$, $S(=O)_2CH_3$, C_1 - C_4 alkyl, C_1 - C_3 alkoxy, C_1 - C_2

30 haloalkyl, and C_1 - C_2 haloalkoxy;

Ring B is selected from:





R^{11} , at each occurrence, is independently selected from
 5 H, =O, $NR^{18}R^{19}$, CF_3 ;

C_1 - C_4 alkyl optionally substituted with 0-1 R^{11a} ;

phenyl substituted with 0-3 R^{11b} ;

C_3 - C_7 carbocycle substituted with 0-3 R^{11b} ; and

5 to 7 membered heterocycle containing 1 to 4

10 heteroatoms selected from nitrogen, oxygen, and
 sulphur, wherein said 5 to 7 membered heterocycle is
 substituted with 0-3 R^{11b} ; wherein said 5 to 7
 membered heterocycle is selected from pyridinyl,
 pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,
 15 pyrrolyl, piperazinyl, piperidinyl, homopiperidinyl,
 pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and
 tetrazolyl;

R^{11a} , at each occurrence, is independently selected from H,
 20 methyl, ethyl, propyl, butyl, methoxy, ethoxy,
 propoxy, phenoxy, F, Cl, =O, $NR^{15}R^{16}$, CF_3 , or phenyl
 substituted with 0-3 R^{11b} ;

R^{11b} , at each occurrence, is independently selected from H,
 25 OH, Cl, F, $NR^{15}R^{16}$, CF_3 , methyl, ethyl, propyl, butyl,
 methoxy, ethoxy, propoxy, C_1 - C_2 haloalkyl, and C_1 - C_2
 haloalkoxy;

W is a bond, $-CH_2-$, $-CH_2CH_2-$;

30 X is a bond;

phenyl substituted with 0-1 R^{Xb} ;

C_3 - C_6 cycloalkyl substituted with 0-1 R^{Xb} ; or

5 to 6 membered heterocycle substituted with 0-1 R^{Xb} ;

R^{Xb} is selected from H, OH, Cl, F, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, methyl, ethyl, propyl, methoxy, ethoxy, propoxy, and -OCF₃;

5

Y is a bond, -C(=O)-, -O-, -S-, -S(=O)-, -S(=O)₂-, -NH-, -N(CH₃)-, or -N(CH₂CH₃)-;

Z is C₁-C₂ alkyl substituted with 1-2 R^{12a};

10

C₆-C₁₀ aryl substituted with 0-4 R^{12b};

C₃-C₁₀ carbocycle substituted with 0-3 R^{12b}; or

5 to 10 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 10 membered heterocycle

15

is substituted with 0-3 R^{12b};

R^{12a}, at each occurrence, is independently selected from

C₆-C₁₀ aryl substituted with 0-4 R^{12b};

C₃-C₁₀ carbocycle substituted with 0-4 R^{12b}; and

20

5 to 10 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 10 membered heterocycle

is substituted with 0-3 R^{12b};

25

R^{12b}, at each occurrence, is independently selected from

H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl,

SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy,

C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and

C₁-C₄ haloalkyl-S-;

30

R¹³, at each occurrence, is independently selected from

H, OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN,

NO₂, NR¹⁵R¹⁶, and CF₃;

35

R¹⁴ is H, phenyl, benzyl, C₁-C₄ alkyl, or C₂-C₄ alkoxyalkyl;

R¹⁵, at each occurrence, is independently selected from H,

C₁-C₄ alkyl, and benzyl;

R¹⁶, at each occurrence, is independently selected from
H, OH, methyl, ethyl, propyl, butyl, benzyl,
phenethyl, methyl-C(=O)-, ethyl-C(=O)-,
5 methyl-S(=O)₂-, and ethyl-S(=O)₂-;

R¹⁸, at each occurrence, is independently selected from
H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and
phenethyl; and

10 R¹⁹, at each occurrence, is independently selected from
H, methyl, ethyl, propyl, and butyl.

[14] In another preferred embodiment the present
15 invention provides for a compound of Formula (Ib) wherein:

R³ and R^{3a} are combined to form a 3-6 membered carbocyclic
moiety selected from cyclopropyl, cyclobutyl,
cyclopentyl, cyclopentenyl, cyclohexyl, and
20 cyclohexenyl; wherein said 3-6 membered carbocyclic
moiety is substituted with 0-1 R⁴;

R⁴ is selected from H, OH, Cl, F, CN, CF₃, methyl, ethyl,
methoxy, ethoxy, allyl, and -OCF₃;

25 R⁵ is C₁-C₄ alkyl substituted with 0-1 R^{5b};
C₂-C₄ alkenyl substituted with 0-1 R^{5b};
C₂-C₄ alkynyl substituted with 0-1 R^{5b};

30 R^{5b}, at each occurrence, is independently selected from:
H, methyl, ethyl, propyl, butyl, CF₃, OR¹⁴, =O;
C₃-C₆ carbocycle substituted with 0-2 R^{5c};
phenyl substituted with 0-3 R^{5c}; or
5 to 6 membered heterocycle containing 1 to 4
35 heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 6 membered heterocycle
is substituted with 0-3 R^{5c}; wherein said 5 to 6
membered heterocycle is selected from pyridinyl,


pyrimidinyl, triazinyl, furanyl, thienyl,
thiazolyl, pyrrolyl, piperazinyl, piperidinyl,
pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and
tetrazolyl;

5

R^{5c} , at each occurrence, is independently selected from H,
OH, Cl, F, $NR^{15}R^{16}$, CF_3 , acetyl, SCH_3 , $S(=O)CH_3$,
 $S(=O)_2CH_3$, methyl, ethyl, propyl, butyl, methoxy,
ethoxy, propoxy, C_1 - C_2 haloalkyl, and C_1 - C_2 haloalkoxy;

10

Ring B is selected from:



15

R^{11} , at each occurrence, is independently selected from
H, =O, $NR^{18}R^{19}$;

C_1 - C_4 alkyl optionally substituted with 0-1 R^{11a} ;

20

phenyl substituted with 0-3 R^{11b} ;

cyclohexyl substituted with 0-3 R^{11b} ;

cycloheptyl substituted with 0-3 R^{11b} ;

5 to 7 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and

25

sulphur, wherein said 5 to 7 membered heterocycle is
substituted with 0-3 R^{11b} ; wherein said 5 to 7

membered heterocycle is selected from pyridinyl,

pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,

pyrrolyl, piperazinyl, piperidinyl, homopiperidinyl,

30

pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and
tetrazolyl;

- 5 R^{11a} , at each occurrence, is independently selected from H, methyl, ethyl, propyl, methoxy, ethoxy, propoxy, phenoxy, F, Cl, =O, $NR^{15}R^{16}$, CF_3 , or phenyl substituted with 0-3 R^{11b} ;
- 10 R^{11b} , at each occurrence, is independently selected from H, OH, Cl, F, $NR^{15}R^{16}$, CF_3 , methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, C_1 - C_2 haloalkyl, and C_1 - C_2 haloalkoxy;
- W is a bond or $-CH_2-$;
- 15 X is a bond;
 phenyl substituted with 0-1 R^{Xb} ;
 C_3 - C_6 cycloalkyl substituted with 0-1 R^{Xb} ; or
 5 to 6 membered heterocycle substituted with 0-1 R^{Xb} ;
- 20 R^{Xb} is selected from H, OH, Cl, F, $NR^{15}R^{16}$, CF_3 , acetyl, methyl, ethyl, methoxy, ethoxy, and $-OCF_3$;
- Y is a bond, $-C(=O)-$, $-O-$, $-S-$, $-S(=O)-$, $-S(=O)_2-$, $-NH-$, $-N(CH_3)-$, or $-N(CH_2CH_3)-$;
- 25 Z is C_1 - C_2 alkyl substituted with 1-2 R^{12a} ;
 C_6 - C_{10} aryl substituted with 0-4 R^{12b} ;
 C_3 - C_{10} carbocycle substituted with 0-3 R^{12b} ; or
 5 to 10 membered heterocycle containing 1 to 4
 heteroatoms selected from nitrogen, oxygen, and
30 sulphur, wherein said 5 to 10 membered heterocycle
 is substituted with 0-3 R^{12b} ;
- 35 R^{12a} , at each occurrence, is independently selected from
 C_6 - C_{10} aryl substituted with 0-4 R^{12b} ;
 C_3 - C_{10} carbocycle substituted with 0-4 R^{12b} ; and
 5 to 10 membered heterocycle containing 1 to 4
 heteroatoms selected from nitrogen, oxygen, and
 sulphur, wherein said 5 to 10 membered heterocycle

is substituted with 0-3 R^{12b}; and wherein said 5 to 10 membered heterocycle is selected from pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, pyrrolyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, tetrazolyl, benzofuranyl, benzothiofuranyl, indolyl, benzimidazolyl, 1H-indazolyl, oxazolidinyl, isoxazolidinyl, benzotriazolyl, benzisoxazolyl, oxindolyl, benzoxazolinyl, quinolinyl, and isoquinolinyl;

R^{12b}, at each occurrence, is independently selected from H, OH, Cl, F, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, and -OCF₃;

R¹³, at each occurrence, is independently selected from H, OH, methyl, ethyl, propyl, butyl, methoxy, ethoxy, Cl, F, Br, CN, NR¹⁵R¹⁶, and CF₃;

R¹⁴ is H, phenyl, benzyl, methyl, ethyl, propyl, or butyl;

R¹⁵, at each occurrence, is independently selected from H, methyl, ethyl, propyl, and butyl;

R¹⁶, at each occurrence, is independently selected from H, OH, methyl, ethyl, propyl, butyl, benzyl, and phenethyl;

R¹⁸, at each occurrence, is independently selected from H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and phenethyl; and

R¹⁹, at each occurrence, is independently selected from H, methyl, ethyl, propyl, and butyl.

[15] In another preferred embodiment the present invention provides for a compound of Formula (Ib) wherein:


R³ and R^{3a} are combined to form cyclobutyl, cyclopentyl, cyclopentenyl, or cyclohexyl;

R⁵ is -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH(CH₃)₂, -CH₂CH₂CH₂CH₃,
 5 -CH(CH₃)CH₂CH₃, -CH₂CH(CH₃)₂, -CH₂C(CH₃)₃,
 -CH₂CH₂CH₂CH₂CH₃, -CH(CH₃)CH₂CH₂CH₃, -CH₂CH(CH₃)CH₂CH₃,
 -CH₂CH₂CH(CH₃)₂, -CH(CH₂CH₃)₂, cyclopropyl-CH₂-,
 cyclobutyl-CH₂-, (2-CH₃-cyclopropyl)CH₂-, or
 (3-CH₃-cyclobutyl)CH₂-;

10

W is a bond or -CH₂-;

X is a bond;



Y is a bond, -C(=O)-, -O-, -S-, -S(=O)-, -S(=O)₂-, -NH-, or
 -N(CH₃)-,

20

Z is phenyl, 2-F-phenyl, 3-F-phenyl, 4-F-phenyl, 2-Cl-phenyl, 3-Cl-phenyl, 4-Cl-phenyl, 2,3-diF-phenyl, 2,4-diF-phenyl, 2,5-diF-phenyl, 2,6-diF-phenyl, 3,4-diF-phenyl, 3,5-diF-phenyl, 2,3-diCl-phenyl, 2,4-diCl-phenyl, 2,5-diCl-phenyl, 2,6-diCl-phenyl, 3,4-diCl-phenyl, 3,5-diCl-phenyl, 3-F-4-Cl-phenyl, 3-F-5-Cl-phenyl, 3-Cl-4-F-phenyl, 2-MeO-phenyl, 3-MeO-phenyl, 4-MeO-phenyl, 2-Me-phenyl, 3-Me-phenyl, 4-Me-phenyl, 2-MeS-phenyl, 3-MeS-phenyl, 4-MeS-phenyl, 2-CF₃O-phenyl, 3-CF₃O-phenyl, 4-CF₃O-phenyl, furanyl, thienyl, pyridyl, 2-Me-pyridyl, 3-Me-pyridyl, 4-Me-pyridyl, 1-imidazolyl, oxazolyl, isoxazolyl, 1-benzimidazolyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, morpholino, N-piperinyl, phenyl-CH₂-, (2-F-phenyl)CH₂-, (3-F-phenyl)CH₂-,
 25
 30
 35

- (4-F-phenyl)CH₂-, (2-Cl-phenyl)CH₂-, (3-Cl-phenyl)CH₂,
(4-Cl-phenyl)CH₂-, (2,3-diF-phenyl)CH₂-,
(2,4-diF-phenyl)CH₂-, (2,5-diF-phenyl)CH₂-,
(2,6-diF-phenyl)CH₂-, (3,4-diF-phenyl)CH₂-,
5 (3,5-diF-phenyl)CH₂-, (2,3-diCl-phenyl)CH₂-,
(2,4-diCl-phenyl)CH₂-, (2,5-diCl-phenyl)CH₂-,
(2,6-diCl-phenyl)CH₂-, (3,4-diCl-phenyl)CH₂-,
(3,5-diCl-phenyl)CH₂-, (3-F-4-Cl-phenyl)CH₂-,
(3-F-5-Cl-phenyl)CH₂-, (3-Cl-4-F-phenyl)CH₂-,
10 (2-MeO-phenyl)CH₂-, (3-MeO-phenyl)CH₂-,
(4-MeO-phenyl)CH₂-, (2-Me-phenyl)CH₂-,
(3-Me-phenyl)CH₂-, (4-Me-phenyl)CH₂-,
(2-MeS-phenyl)CH₂-, (3-MeS-phenyl)CH₂-,
4-MeS-phenyl)CH₂-, (2-CF₃O-phenyl)CH₂-,
15 (3-CF₃O-phenyl)CH₂-, (4-CF₃O-phenyl)CH₂-,
(furanyl)CH₂-, (thienyl)CH₂-, (pyridyl)CH₂-,
(2-Me-pyridyl)CH₂-, (3-Me-pyridyl)CH₂-,
(4-Me-pyridyl)CH₂-, (1-imidazolyl)CH₂-,
(oxazolyl)CH₂-, (isoxazolyl)CH₂-,
20 (1-benzimidazolyl)CH₂-, (cyclopropyl)CH₂-,
(cyclobutyl)CH₂-, (cyclopentyl)CH₂-,
(cyclohexyl)CH₂-, (morpholino)CH₂-,
(N-pipridinyl)CH₂-, phenyl-CH₂CH₂-,
(phenyl)₂CHCH₂-, (2-F-phenyl)CH₂CH₂-,
25 (3-F-phenyl)CH₂CH₂-, (4-F-phenyl)CH₂CH₂-,
(2-Cl-phenyl)CH₂CH₂-, (3-Cl-phenyl)CH₂CH₂-,
(4-Cl-phenyl)CH₂CH₂-, (2,3-diF-phenyl)CH₂CH₂-,
(2,4-diF-phenyl)CH₂CH₂-, (2,5-diF-phenyl)CH₂CH₂-,
(2,6-diF-phenyl)CH₂CH₂-, (3,4-diF-phenyl)CH₂CH₂-,
30 (3,5-diF-phenyl)CH₂CH₂-, (2,3-diCl-phenyl)CH₂CH₂-,
(2,4-diCl-phenyl)CH₂CH₂-, (2,5-diCl-phenyl)CH₂CH₂-,
(2,6-diCl-phenyl)CH₂CH₂-, (3,4-diCl-phenyl)CH₂CH₂-,
(3,5-diCl-phenyl)CH₂CH₂-, (3-F-4-Cl-phenyl)CH₂CH₂-,
(3-F-5-Cl-phenyl)CH₂CH₂-, (3-Cl-4-F-phenyl)CH₂CH₂-,
35 (2-MeO-phenyl)CH₂CH₂-, (3-MeO-phenyl)CH₂CH₂-,
(4-MeO-phenyl)CH₂CH₂-, (2-Me-phenyl)CH₂CH₂-,
(3-Me-phenyl)CH₂CH₂-, (4-Me-phenyl)CH₂CH₂-,
(2-MeS-phenyl)CH₂CH₂-, (3-MeS-phenyl)CH₂CH₂-,


(4-MeS-phenyl)CH₂CH₂-, (2-CF₃O-phenyl)CH₂CH₂-,
 (3-CF₃O-phenyl)CH₂CH₂-, (4-CF₃O-phenyl)CH₂CH₂-,
 (furanyl)CH₂CH₂-, (thienyl)CH₂CH₂-, (pyridyl)CH₂CH₂-,
 (2-Me-pyridyl)CH₂CH₂-, (3-Me-pyridyl)CH₂CH₂-,
 5 (4-Me-pyridyl)CH₂CH₂-, (imidazolyl)CH₂CH₂-,
 (oxazolyl)CH₂CH₂-, (isoxazolyl)CH₂CH₂-,
 (benzimidazolyl)CH₂CH₂-, (cyclopropyl)CH₂CH₂-,
 (cyclobutyl)CH₂CH₂-, (cyclopentyl)CH₂CH₂-,
 (cyclohexyl)CH₂CH₂-, (morpholino)CH₂CH₂-, or
 10 (N-pipridinyl)CH₂CH₂-;

R¹¹, at each occurrence, is independently selected from
 H, =O, methyl, ethyl, phenyl, benzyl, phenethyl,
 4-F-phenyl, (4-F-phenyl)CH₂-, (4-F-phenyl)CH₂CH₂-,
 15 3-F-phenyl, (3-F-phenyl)CH₂-, (3-F-phenyl)CH₂CH₂-,
 2-F-phenyl, (2-F-phenyl)CH₂-, (2-F-phenyl)CH₂CH₂-,
 4-Cl-phenyl, (4-Cl-phenyl)CH₂-, (4-Cl-phenyl)CH₂CH₂-,
 3-Cl-phenyl, (3-Cl-phenyl)CH₂-, (3-Cl-phenyl)CH₂CH₂-,
 4-CH₃-phenyl, (4-CH₃-phenyl)CH₂-, (4-CH₃-phenyl)CH₂CH₂-,
 20 3-CH₃-phenyl, (3-CH₃-phenyl)CH₂-, (3-CH₃-phenyl)CH₂CH₂-,
 4-CF₃-phenyl, (4-CF₃-phenyl)CH₂-, (4-CF₃-phenyl)CH₂CH₂-,
 pyrid-2-yl, pyrid-3-yl, pyrid-4-yl, cyclohexyl,
 cycloheptyl, piperidinyl, or homopiperidinyl; and

25 R¹³, at each occurrence, is independently selected from
 H, F, Cl, OH, -CH₃, -CH₂CH₃, -OCH₃, or -CF₃.


[16] In another preferred embodiment the present
 invention provides for a compound of Formula (Ic):

30



or a pharmaceutically acceptable salt or prodrug thereof.

[17] In another preferred embodiment the present invention provides for a compound of Formula (Id):




5

(Id)

or a pharmaceutically acceptable salt or prodrug thereof.

[18] In another preferred embodiment the present invention provides for a compound of Formula (Ie):

10




(Ie)

or a pharmaceutically acceptable salt or prodrug thereof.

15


[19] In another preferred embodiment the present invention provides for a compound of Formula (If):



(If)

20 or a pharmaceutically acceptable salt or prodrug thereof.

[20] In another preferred embodiment the present invention provides for a compound of Formula (Ig):



25

(Ig)

or a pharmaceutically acceptable salt or prodrug thereof.

[21] In another preferred embodiment the present
5 invention provides for a compound of Formula (I) selected
from:

1-[3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-
benzodiazepin-3-ylcarbamoyl]-butyl]-cyclopent-3-
10 enecarboxylic amide;

1-[3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-
benzodiazepin-3-ylcarbamoyl]-butyl]-cyclopentanecarboxylic
amide;

15 1-[3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-(4-
chlorophenyl)-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-
cyclohexanecarboxylic amide;

20 1-[3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-
benzodiazepin-3-ylcarbamoyl]-butyl]-cyclohexanecarboxylic
amide;

25 1-[3-Methyl-1-(5-methyl-6-oxo-6,7-dihydro-5H-dibenzo[b,d]-
azepin-7-ylcarbamoyl)-butyl]-cyclohexanecarboxylic acid
amide;

30 1-[3-Methyl-1-(5-methyl-6-oxo-6,7-dihydro-5H-dibenzo[b,d]-
azepin-7-ylcarbamoyl)-butyl]-cyclohexanecarboxylic acid
amide;

35 1-(1-{1-[3-(2-Fluoro-phenoxy)-benzyl]-2-oxo-2,3,4,5-
tetrahydro-1H-benzo[b]azepin-3-ylcarbamoyl}-3-methyl-
butyl)-cyclopent-3-enecarboxylic acid amide;

1-{3-Methyl-1-[2-oxo-1-(3-phenylamino-benzyl)-2,3,4,5-
tetrahydro-1H-benzo[b]azepin-3-ylcarbamoyl]-butyl}-
cyclopent-3-enecarboxylic acid amide;

1-{3-Methyl-1-[2-oxo-1-(3-phenylamino-benzyl)-2,3,4,5-tetrahydro-1H-benzo[b]azepin-3-ylcarbamoyl]-butyl}-cyclopentanecarboxylic acid amide;

5

1-[2-Cyclopropyl-1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-ylcarbamoyl]-ethyl]-cyclopent-3-enecarboxylic amide;

10 1-[2-Cyclopropyl-1-[1,3-dihydro-1-methyl-2-oxo-5-(2-trifluoromethylphenyl)-2H-1,4-benzodiazepin-3-ylcarbamoyl]-ethyl]-cyclopent-3-enecarboxylic amide;

15 1-[2-Cyclopropyl-1-(5-methyl-6-oxo-6,7-dihydro-5H-dibenzo[b,d]azepin-7-ylcarbamoyl)-ethyl]-cyclopent-3-enecarboxylic acid amide;

20 1-{3-Methyl-1-[2-oxo-1-(3-o-tolylamino-benzyl)-azepan-3-ylcarbamoyl]-butyl}-cyclopent-3-enecarboxylic acid amide;

1-[3-Methyl-1-(5-methyl-6-oxo-6,7-dihydro-5H-dibenzo[b,d]azepin-7-ylcarbamoyl)-butyl]-cyclopent-3-enecarboxylic acid amide;

25 1-[3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-(4-chlorophenyl)-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclopent-3-enecarboxylic amide;

30 1-[3-Methyl-1-(5-methyl-6-oxo-6,7-dihydro-5H-dibenzo[b,d]azepin-7-ylcarbamoyl)-butyl]-cyclopentanecarboxylic acid amide;

35 1-[3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-(4-chlorophenyl)-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclopentanecarboxylic amide;

- 1-[3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-(4-trifluoromethylphenyl)-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclopent-3-enecarboxylic amide;
- 5 1-[3-methyl-1-[1,3-dihydro-1-(i-propyl)-2-oxo-5-(2-fluorophenyl)-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclopent-3-enecarboxylic amide;
- 10 1-[3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-(4-trifluoromethylphenyl)-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclopentanecarboxylic amide;
- 15 1-[3-methyl-1-[1,3-dihydro-1-(2-cyclopropylethyl)-2-oxo-5-(2-fluorophenyl)-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclopent-3-enecarboxylic amide;
- 20 1-[3-methyl-1-[1,3-dihydro-1-(2-methylpropyl)-2-oxo-5-(2-fluorophenyl)-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclobutanecarboxylic amide;
- 25 1-[3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclobutanecarboxylic amide; and
- 30 1-[3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-cycloheptyl-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclopent-3-enecarboxylic amide.


In a more preferred embodiment of the present invention, Q is NH₂.

35


It is appreciated that certain features of the invention, which are, for clarity, described herein in the context of separate embodiments, may also be provided in

combination in a single embodiment. As such, it is understood that any and all embodiments of the present invention may be taken in conjunction with any other embodiment to describe additional embodiments of the present invention. Conversely, various features of the invention which are for brevity, described herein in the context of a single embodiment, may also be provided separately or in any subcombination. As such, it is understood that any elements of an embodiment are meant to be combined with any and all other elements from any of the embodiments to describe additional embodiments.


In a preferred embodiment Ring B is selected from:



In another preferred embodiment Ring B is selected from:




5 In another preferred embodiment Ring B is singly:



10

In another preferred embodiment Ring B is singly:



15

Also included in the present invention are compounds as set forth in the embodiments above wherein R^3 and R^{3a} are combined to form a 3-8 membered carbocyclic moiety; and R^5 and R^{5a} are not combined to form a 3-8 membered carbocyclic moiety.

Also included in the present invention are compounds as set forth in the embodiments above wherein R^3 and R^{3a} are not combined to form a 3-8 membered carbocyclic moiety; and R^5 and R^{5a} are combined to form a 3-8 membered carbocyclic moiety;

Also included in the present invention are compounds as set forth in the embodiments above wherein R^3 and R^{3a} are combined to form a 3-8 membered carbocyclic moiety; and R^5 and R^{5a} are also combined to form a 3-8 membered carbocyclic moiety.

Also included in the present invention are compounds as set forth in the embodiments above wherein R^3 and R^{3a} are combined to form a 3-8 membered carbocyclic moiety which is saturated or partially unsaturated.

In another preferred embodiment R^3 and R^{3a} are combined to form a 3-8 membered carbocyclic moiety selected from, cyclobutyl, cyclopentyl, cyclopentenyl, and cyclohexyl.

In another preferred embodiment R^3 and R^{3a} are combined to form a cyclobutyl moiety.

In another preferred embodiment R^3 and R^{3a} are combined to form a cyclopentyl moiety.

In another preferred embodiment R^3 and R^{3a} are combined to form a cyclopentenyl moiety.

In another preferred embodiment R^3 and R^{3a} are combined to form a cyclohexyl moiety.

5 Also included in the present invention are compounds as set forth in the embodiments above wherein R^5 and R^{5a} are combined to form a 3-8 membered carbocyclic moiety which is saturated or partially unsaturated.

10 In another preferred embodiment R^5 and R^{5a} are combined to form a 3-8 membered carbocyclic moiety selected from, cyclobutyl, cyclopentyl, cyclopentenyl, and cyclohexyl.

15 In another preferred embodiment R^5 and R^{5a} are combined to form a cyclobutyl moiety.

In another preferred embodiment R^5 and R^{5a} are combined to form a cyclopentyl moiety.

20 In another preferred embodiment R^5 and R^{5a} are combined to form a cyclopentenyl moiety.

In another preferred embodiment R^5 and R^{5a} are combined to form a cyclohexyl moiety.

25 Also included in the present invention are compounds as set forth in the embodiments above wherein R^3 and R^{3a} are combined to form a 3-8 membered carbocyclic moiety; R^5 is C_1-C_6 alkyl, C_2-C_6 alkenyl, C_2-C_6 alkynyl, or $(C_3-C_6$ cycloalkyl) C_1-C_3 alkyl-; and R^{5a} is H.

30 In another preferred embodiment R^3 and R^{3a} are combined to form a 3-8 membered carbocyclic moiety; R^5 is C_1-C_6 alkyl, C_2-C_6 alkenyl, or $(C_3-C_6$ cycloalkyl) C_1-C_3 alkyl-; and R^{5a} is H.

35 In three more preferred embodiments R^3 and R^{3a} are combined to form a 3-8 membered carbocyclic moiety; R^5 is C_1-C_6

alkyl or C₂-C₆ alkenyl or (C₃-C₆ cycloalkyl)C₁-C₃ alkyl-;
and R^{5a} is H.

5 In another preferred embodiment R³ and R^{3a} are combined to form a 3-8 membered carbocyclic moiety; R⁵ is C₁-C₄ alkyl, C₂-C₄ alkenyl, or (C₃-C₆ cycloalkyl)C₁-C₂ alkyl-; and R^{5a} is H.

10 In another preferred embodiment R³ and R^{3a} are combined to form a 3-8 membered carbocyclic moiety; R⁵ is methyl, ethyl, propyl, butyl, allyl, cyclopropylmethyl, cyclobutylmethyl, or cyclohexylmethyl; and R^{5a} is H.

15 In another preferred embodiment R³ and R^{3a} are combined to form a 3-8 membered carbocyclic moiety; R⁵ is ethyl, propyl, butyl, allyl, or cyclopropylmethyl; and R^{5a} is H.

20 In another preferred embodiment R³ and R^{3a} are combined to form a 3-8 membered carbocyclic moiety; R⁵ is n-propyl, i-propyl, n-butyl, i-butyl, s-butyl, allyl, or cyclopropylmethyl; and R^{5a} is H.

25 In another preferred embodiment R³ and R^{3a} are combined to form a 3-8 membered carbocyclic moiety; R⁵ is n-butyl, i-butyl, or cyclopropylmethyl; and R^{5a} is H.

30 In more preferred embodiments R³ and R^{3a} are combined to form a 3-8 membered carbocyclic moiety; R⁵ is methyl or ethyl or propyl or butyl or allyl or cyclopropylmethyl or cyclobutylmethyl or cyclohexylmethyl; and R^{5a} is H.

35 Also included in the present invention are compounds as set forth in the embodiments above wherein R⁵ is C₁-C₈ alkyl, C₂-C₈ alkenyl, C₂-C₈ alkynyl, or (C₃-C₆ cycloalkyl)C₁-C₄ alkyl, (NR¹⁵R¹⁶)C₁-C₄ alkyl.

In another preferred embodiment R⁵ is C₂-C₈ alkyl, C₂-C₈ alkenyl, or C₂-C₈ alkynyl.

In another preferred embodiment R^5 is C_3 - C_8 alkyl, C_3 - C_8 alkenyl, or C_3 - C_8 alkynyl.

- 5 In another preferred embodiment R^5 is C_4 - C_8 alkyl, C_4 - C_8 alkenyl, C_4 - C_8 alkynyl.

In another preferred embodiment R^5 is C_2 - C_8 alkyl.

In another preferred embodiment R^5 is C_3 - C_8 alkyl.

- 10 In another preferred embodiment R^5 is C_4 - C_8 alkyl.

In another preferred embodiment R^5 is

(C_3 - C_6 cycloalkyl) C_1 - C_4 alkyl.

In another preferred embodiment R^5 is $(NR^{15}R^{16})C_1$ - C_4 alkyl.

- 15 In another preferred embodiment R^5 is $-CH_3$, $-CH_2CH_3$,
 $-CH_2CH_2CH_3$, $-CH_2CH(CH_3)_2$, $-CH_2CH_2CH_2CH_3$, $-CH_2CH_2CH_2CH_2CH_3$,
 $-CH_2CH_2CH(CH_3)_2$, $-CH_2CH_2CH_2CH_2CH_2CH_3$, $-CH_2CH_2CH_2CH(CH_3)_2$,
 $-CH_2CH_2CH_2CH_2CH(CH_3)_2$, $-CH_2NH_2$, $-CH_2N(CH_3)_2$, $-$
 $CH_2N(CH_2CH_3)_2$, $-CH_2CH_2NH_2$, $-CH_2CH_2N(CH_3)_2$, $-$
20 $CH_2CH_2N(CH_2CH_3)_2$,
 $-CH_2$ -cyclopropyl, $-CH_2$ -cyclobutyl, $-CH_2$ -cyclopentyl, or
 $-CH_2$ -cyclohexyl.

In another preferred embodiment R^5 is $-CH_3$, $-CH_2CH_3$,

- 25 $-CH_2CH_2CH_3$, $-CH_2CH(CH_3)_2$, $-CH_2CH_2CH_2CH_3$, $-CH_2CH_2CH_2CH_2CH_3$,
 $-CH_2CH_2CH(CH_3)_2$, $-CH_2CH_2CH_2CH_2CH_2CH_3$, $-CH_2CH_2CH_2CH(CH_3)_2$, or
 $-CH_2CH_2CH_2CH_2CH(CH_3)_2$.

In another preferred embodiment R^5 is $-CH_2NH_2$, $-CH_2N(CH_3)_2$,

- 30 $-CH_2N(CH_2CH_3)_2$, $-CH_2CH_2NH_2$, $-CH_2CH_2N(CH_3)_2$, or
 $-CH_2CH_2N(CH_2CH_3)_2$.

In another preferred embodiment R^5 is $-CH_2$ -cyclopropyl,

$-CH_2$ -cyclobutyl, $-CH_2$ -cyclopentyl, or $-CH_2$ -cyclohexyl.

35

Also included in the present invention are compounds as set forth in the embodiments above wherein R^6 is H.

Also included in the present invention are compounds as set forth in the embodiments above wherein R¹¹ is

H, NR¹⁸R¹⁹;

C₁-C₄ alkyl optionally substituted with 0-1 R^{11a};

5 phenyl substituted with 0-3 R^{11b};

C₃-C₇ cycloalkyl substituted with 0-3 R^{11b}; or

pyridinyl substituted with 0-3 R^{11b};

wherein R^{11a} is phenyl substituted with 0-3 R^{11b};

10 wherein R^{11b}, at each occurrence, is independently selected from H, OH, Cl, F, CF₃, methyl, ethyl, propyl, butyl, methoxy, ethoxy, and propoxy.

In another preferred embodiment R¹¹ is independently

15 selected from

H, methyl, ethyl, cyclopentyl, cyclohexyl, cycloheptyl, phenyl, benzyl, phenethyl,

4-F-phenyl, (4-F-phenyl)CH₂-, (4-F-phenyl)CH₂CH₂-,

3-F-phenyl, (3-F-phenyl)CH₂-, (3-F-phenyl)CH₂CH₂-,

20 2-F-phenyl, (2-F-phenyl)CH₂-, (2-F-phenyl)CH₂CH₂-,

4-Cl-phenyl, (4-Cl-phenyl)CH₂-, (4-Cl-phenyl)CH₂CH₂-,

3-Cl-phenyl, (3-Cl-phenyl)CH₂-, (3-Cl-phenyl)CH₂CH₂-,

4-CH₃-phenyl, (4-CH₃-phenyl)CH₂-, (4-CH₃-phenyl)CH₂CH₂-,

3-CH₃-phenyl, (3-CH₃-phenyl)CH₂-, (3-CH₃-phenyl)CH₂CH₂-,

25 4-CF₃-phenyl, (4-CF₃-phenyl)CH₂-, (4-CF₃-phenyl)CH₂CH₂-,

pyrid-2-yl, 4-F-pyrid-2-yl, 4-Cl-pyrid-2-yl,

4-CH₃-pyrid-2-yl, 4-CF₃-pyrid-2-yl, pyrid-3-yl,

4-F-pyrid-3-yl, 4-Cl-pyrid-3-yl, 4-CH₃-pyrid-3-yl,

4-CF₃-pyrid-3-yl, and pyrid-4-yl.

30

In another preferred embodiment R¹¹ is independently selected from

H, methyl, ethyl, cyclopentyl, cyclohexyl, cycloheptyl, phenyl, 4-F-phenyl, 3-F-phenyl,

35 2-F-phenyl, 4-Cl-phenyl, 3-Cl-phenyl, 4-CH₃-phenyl,

3-CH₃-phenyl, 4-CF₃-phenyl, pyrid-2-yl, 4-F-pyrid-2-yl,

4-Cl-pyrid-2-yl, 4-CH₃-pyrid-2-yl, and 4-CF₃-pyrid-2-yl.

In another preferred embodiment R^{11} is independently selected from phenyl, 4-F-phenyl, 3-F-phenyl, 2-F-phenyl, 4-Cl-phenyl, 3-Cl-phenyl, 4-CH₃-phenyl, 3-CH₃-phenyl, and 4-CF₃-phenyl.

5

In another preferred embodiment R^{11} is independently selected from cyclopentyl, cyclohexyl, and cycloheptyl.

10 In another preferred embodiment R^{11} is independently selected from pyrid-2-yl, 4-F-pyrid-2-yl, 4-Cl-pyrid-2-yl, 4-CH₃-pyrid-2-yl, and 4-CF₃-pyrid-2-yl.

15 Also included in the present invention are compounds as set forth in the embodiments above wherein W may be selected from a bond, -CH₂-, -CH₂CH₂-, or -CH(CH₃)-.

In another preferred embodiment W is a bond or -(CH₂)_p-.
In another preferred embodiment W is a bond, -CH₂-, or -CH₂CH₂-.

20 In another preferred embodiment W is a bond or -CH₂-.
In another preferred embodiment W is -CH₂-.
In another preferred embodiment W is a bond.

25 Also included in the present invention are compounds as set forth in the embodiments above wherein the integer p may be selected from 0, 1, 2, or 3.

In another preferred embodiment the integer p is 0, 1 or 2.
In another preferred embodiment the integer p is 0 or 1.
30 In another preferred embodiment the integer p is 0.

Also included in the present invention are compounds as set forth in the embodiments above wherein X is a bond, C₆-C₁₀ aryl, C₃-C₁₀ carbocycle or 5 to 10 membered heterocycle.


35

In another preferred embodiment X is a bond, phenyl, C₃-C₆ carbocycle, or 5 to 6 membered heterocycle.


In another preferred embodiment X is a bond, phenyl, C₃-C₆ cycoalkyl, or 5 to 6 membered heterocycle.

In another preferred embodiment X is a bond;

5



10 In another preferred embodiment X is a bond;



15 In another preferred embodiment X is a bond or phen-1,3-diyl.

In another preferred embodiment X is phen-1,3-diyl.

In another preferred embodiment X is a bond.

20 Also included in the present invention are compounds as set forth in the embodiments above wherein Y is a bond, -C(=O)-, -O-, -S-, -S(=O)-, -S(=O)₂-, -N(R¹⁹)-, -C(=O)NR^{19b}-, -NR^{19b}C(=O)-, -NR^{19b}S(=O)₂-, -S(=O)₂NR^{19b}-, -NR^{19b}S(=O)-, -S(=O)NR^{19b}-, -C(=O)O-, or -OC(=O)-.

25 In another preferred embodiment Y is a bond, -C(=O)-, -O-, -S-, -S(=O)-, -S(=O)₂-, -NH-, -N(CH₃)-, -C(=O)NH-, -NHC(=O)-, -NHS(=O)₂-, -S(=O)₂NH-, -NHS(=O)-, -S(=O)NH-, -C(=O)O-, or -OC(=O)-.

30 In another preferred embodiment Y is a bond, -C(=O)-, -O-, -S-, -S(=O)-, -S(=O)₂-, -NH-, -N(CH₃)-, or -N(CH₂CH₃)-.

In another preferred embodiment Y is a bond, $-C(=O)-$, $-O-$, $-S-$, $-S(=O)-$, $-S(=O)_2-$, $-NH-$, or $-N(CH_3)-$.

5 In another preferred embodiment Y is a bond, $-C(=O)-$, $-O-$, $-NH-$, or $-N(CH_3)-$.

In another preferred embodiment Y is $-O-$.

In another preferred embodiment Y is $-NH-$.

In another preferred embodiment Y is $-N(CH_3)-$.

In another preferred embodiment Y is a bond.

10

Also included in the present invention are compounds as set forth in the embodiments above wherein Z is

C_1-C_4 alkyl, C_2-C_4 alkenyl, C_2-C_4 alkynyl,

C_1-C_2 alkyl substituted with 1-2 R^{12a} ;

15 phenyl substituted with 0-4 R^{12b} ;

C_3-C_6 carbocycle substituted with 0-3 R^{12b} ; or

5 to 6 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 6 membered heterocycle

20

is substituted with 0-3 R^{12b} ;

wherein R^{12a} is phenyl substituted with 0-4 R^{12b} ;

C_3-C_6 carbocycle substituted with 0-4 R^{12b} ; or

5 to 6 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and

25

sulphur, wherein said 5 to 6 membered heterocycle

is substituted with 0-3 R^{12b} ; and

wherein R^{12b} , at each occurrence, is independently

selected from H, OH, Cl, F, $NR^{15}R^{16}$, CF_3 , acetyl,

SCH_3 , $S(=O)CH_3$, $S(=O)_2CH_3$, methyl, ethyl, propyl,

30

butyl, methoxy, ethoxy, propoxy, and $-OCF_3$;

In another preferred embodiment Z is

C_1-C_2 alkyl substituted with 1-2 R^{12a} ; or

phenyl substituted with 0-4 R^{12b} ;

35

wherein R^{12a} is phenyl substituted with 0-4 R^{12b} ;

wherein R^{12b} , at each occurrence, is independently

selected from H, OH, Cl, F, $NR^{15}R^{16}$, CF_3 , acetyl,

SCH₃, S(=O)CH₃, S(=O)₂CH₃, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, and -OCF₃;

In another preferred embodiment Z is C₁-C₄ alkyl, C₂-C₄ alkenyl, or C₂-C₄ alkynyl.

In another preferred embodiment Z is phenyl, 2-F-phenyl, 3-F-phenyl, 4-F-phenyl, 2-Cl-phenyl, 3-Cl-phenyl, 4-Cl-phenyl, 2,3-diF-phenyl, 2,4-diF-phenyl, 2,5-diF-phenyl, 2,6-diF-phenyl, 3,4-diF-phenyl, 3,5-diF-phenyl, 2,3-diCl-phenyl, 2,4-diCl-phenyl, 2,5-diCl-phenyl, 2,6-diCl-phenyl, 3,4-diCl-phenyl, 3,5-diCl-phenyl, 3-F-4-Cl-phenyl, 3-F-5-Cl-phenyl, 3-Cl-4-F-phenyl, 2-MeO-phenyl, 3-MeO-phenyl, 4-MeO-phenyl, 2-Me-phenyl, 3-Me-phenyl, 4-Me-phenyl, 2-MeS-phenyl, 3-MeS-phenyl, 4-MeS-phenyl, 2-CF₃O-phenyl, 3-CF₃O-phenyl, 4-CF₃O-phenyl, furanyl, thienyl, pyridyl, 2-Me-pyridyl, 3-Me-pyridyl, 4-Me-pyridyl, 1-imidazolyl, oxazolyl, isoxazolyl, 1-benzimidazolyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, morpholino, N-piperinyl, phenyl-CH₂-, (2-F-phenyl)CH₂-, (3-F-phenyl)CH₂-, (4-F-phenyl)CH₂-, (2-Cl-phenyl)CH₂-, (3-Cl-phenyl)CH₂-, (4-Cl-phenyl)CH₂-, (2,3-diF-phenyl)CH₂-, (2,4-diF-phenyl)CH₂-, (2,5-diF-phenyl)CH₂-, (2,6-diF-phenyl)CH₂-, (3,4-diF-phenyl)CH₂-, (3,5-diF-phenyl)CH₂-, (2,3-diCl-phenyl)CH₂-, (2,4-diCl-phenyl)CH₂-, (2,5-diCl-phenyl)CH₂-, (2,6-diCl-phenyl)CH₂-, (3,4-diCl-phenyl)CH₂-, (3,5-diCl-phenyl)CH₂-, (3-F-4-Cl-phenyl)CH₂-, (3-F-5-Cl-phenyl)CH₂-, (3-Cl-4-F-phenyl)CH₂-, (2-MeO-phenyl)CH₂-, (3-MeO-phenyl)CH₂-, (4-MeO-phenyl)CH₂-, (2-Me-phenyl)CH₂-, (3-Me-phenyl)CH₂-, (4-Me-phenyl)CH₂-, (2-MeS-phenyl)CH₂-, (3-MeS-phenyl)CH₂-, (4-MeS-phenyl)CH₂-, (2-CF₃O-phenyl)CH₂-, (3-CF₃O-phenyl)CH₂-, (4-CF₃O-phenyl)CH₂-, (furanyl)CH₂-, (thienyl)CH₂-, (pyridyl)CH₂-, (2-Me-pyridyl)CH₂-, (3-Me-pyridyl)CH₂-,

- (4-Me-pyridyl)CH₂-, (1-imidazolyl)CH₂-,
 (oxazolyl)CH₂-, (isoxazolyl)CH₂-,
 (1-benzimidazolyl)CH₂-, (cyclopropyl)CH₂-,
 (cyclobutyl)CH₂-, (cyclopentyl)CH₂-, (cyclohexyl)CH₂-,
 5 (morpholino)CH₂-, (N-pipridinyl)CH₂-, or (phenyl)₂CH-.

- In another preferred embodiment Z is phenyl, 2-F-phenyl,
 3-F-phenyl, 4-F-phenyl, 2-Cl-phenyl, 3-Cl-phenyl,
 4-Cl-phenyl, 2,3-diF-phenyl, 2,4-diF-phenyl,
 10 2,5-diF-phenyl, 2,6-diF-phenyl, 3,4-diF-phenyl,
 3,5-diF-phenyl, 2,3-diCl-phenyl, 2,4-diCl-phenyl,
 2,5-diCl-phenyl, 2,6-diCl-phenyl, 3,4-diCl-phenyl,
 3,5-diCl-phenyl, 3-F-4-Cl-phenyl, 3-F-5-Cl-phenyl,
 3-Cl-4-F-phenyl, 2-MeO-phenyl, 3-MeO-phenyl, 4-MeO-phenyl,
 15 2-Me-phenyl, 3-Me-phenyl, 4-Me-phenyl, 2-MeS-phenyl,
 3-MeS-phenyl, 4-MeS-phenyl, 2-CF₃O-phenyl, 3-CF₃O-phenyl,
 4-CF₃O-phenyl, or 4-phenyl-phenyl.

- In another preferred embodiment Z is phenyl-CH₂-,
 20 (2-F-phenyl)CH₂-, (3-F-phenyl)CH₂-, (4-F-phenyl)CH₂-,
 (2-Cl-phenyl)CH₂-, (3-Cl-phenyl)CH₂-,
 (4-Cl-phenyl)CH₂-, (2,3-diF-phenyl)CH₂-,
 (2,4-diF-phenyl)CH₂-, (2,5-diF-phenyl)CH₂-,
 (2,6-diF-phenyl)CH₂-, (3,4-diF-phenyl)CH₂-,
 25 (3,5-diF-phenyl)CH₂-, (2,3-diCl-phenyl)CH₂-,
 (2,4-diCl-phenyl)CH₂-, (2,5-diCl-phenyl)CH₂-,
 (2,6-diCl-phenyl)CH₂-, (3,4-diCl-phenyl)CH₂-,
 (3,5-diCl-phenyl)CH₂-, (3-F-4-Cl-phenyl)CH₂-,
 (3-F-5-Cl-phenyl)CH₂-, (3-Cl-4-F-phenyl)CH₂-,
 30 (2-MeO-phenyl)CH₂-, (3-MeO-phenyl)CH₂-,
 (4-MeO-phenyl)CH₂-, (2-Me-phenyl)CH₂-,
 (3-Me-phenyl)CH₂-, (4-Me-phenyl)CH₂-,
 (2-MeS-phenyl)CH₂-, (3-MeS-phenyl)CH₂-,
 4-MeS-phenyl)CH₂-, (2-CF₃O-phenyl)CH₂-,
 35 (3-CF₃O-phenyl)CH₂-, (4-CF₃O-phenyl)CH₂-,
 or (phenyl)₂CH-.

Also included in the present invention are compounds as set forth in the embodiments above wherein R^{13} , at each occurrence, is independently selected from H, F, Cl, OH, $-CH_3$, $-CH_2CH_3$, $-OCH_3$, and $-CF_3$.

5

Also included in the present invention are compounds as set forth in the embodiments above wherein R^{14} is H, phenyl, benzyl, methyl, ethyl, propyl, or butyl;

10 Also included in the present invention are compounds as set forth in the embodiments above wherein R^{15} , at each occurrence, is independently selected from H, methyl, ethyl, propyl, and butyl.

15 Also included in the present invention are compounds as set forth in the embodiments above wherein R^{16} , at each occurrence, is independently selected from H, OH, methyl, ethyl, propyl, butyl, benzyl, and phenethyl.

20 Also included in the present invention are compounds as set forth in the embodiments above wherein R^{18} , at each occurrence, is independently selected from H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and phenethyl.

25 Also included in the present invention are compounds as set forth in the embodiments above wherein R^{19} , at each occurrence, is independently selected from H, methyl, ethyl, propyl, and butyl.

30 In a second embodiment, the present invention provides a pharmaceutical composition comprising a compound of Formula (I) and a pharmaceutically acceptable carrier.

35 In a third embodiment, the present invention provides a method for the treatment of neurological disorders associated with β -amyloid production comprising administering to a host in need of such treatment a

therapeutically effective amount of a compound of Formula (I).

5 In a preferred embodiment the neurological disorder associated with β -amyloid production is Alzheimer's Disease.

10 In a fourth embodiment, the present invention provides a method for inhibiting γ -secretase activity for the treatment of a physiological disorder associated with inhibiting γ -secretase activity comprising administering to a host in need of such inhibition a therapeutically effective amount of a compound of Formula (I) that inhibits γ -secretase activity.

15 Thus, the present invention provides a method for inhibiting γ -secretase activity comprising administering to a host in need of such inhibition a therapeutically effective amount of a compound of Formula (I) that inhibits γ -secretase activity.

20 In a preferred embodiment the physiological disorder associated with inhibiting γ -secretase activity is Alzheimer's Disease.

25 In a fifth embodiment, the present invention provides a compound of Formula (I) for use in therapy.

30 In a preferred embodiment the present invention provides a compound of Formula (I) for use in therapy of Alzheimer's Disease.

35 In a sixth embodiment, the present invention provides for the use of a compound of Formula (I) for the manufacture of a medicament for the treatment of Alzheimer's Disease.

DEFINITIONS

As used herein, the term "A β " denotes the protein designated A β , β -amyloid peptide, and sometimes β /A4, in the art. A β is an approximately 4.2 kilodalton (kD) protein of about 39 to 43 amino acids found in amyloid plaques, the walls of meningeal and parenchymal arterioles, small arteries, capillaries, and sometimes, venules. The isolation and sequence data for the first 28 amino acids are described in U.S. Pat. No 4,666,829. The 43 amino acid sequence is:

```

1
Asp   Ala   Glu   Phe   Arg   His   Asp   Ser   Gly   Tyr   (
11
Glu   Val   His   His   Gln   Lys   Leu   Val   Phe   Phe
21
Ala   Glu   Asp   Val   Gly   Ser   Asn   Lys   Gly   Ala
31
Ile   Ile   Gly   Leu   Met   Val   Gly   Gly   Val   Val
41
Ile   Ala   Thr

```

The term "APP", as used herein, refers to the protein known in the art as β amyloid precursor protein. This protein is the precursor for A β and through the activity of "secretase" enzymes, as used herein, it is processed into A β . Differing secretase enzymes, known in the art, have been designated β secretase, generating the N-terminus of A β , α secretase cleaving around the 16/17 peptide bond in A β , and " γ secretases", as used herein, generating C-terminal A β fragments ending at position 38, 39, 40, 42, and 43 or generating C-terminal extended precursors which are subsequently truncated to the above polypeptides.

The compounds herein described may have asymmetric centers. Compounds of the present invention containing an asymmetrically substituted atom may be isolated in optically active or racemic forms. It is well known in the

art how to prepare optically active forms, such as by resolution of racemic forms or by synthesis from optically active starting materials. Many geometric isomers of olefins, C=N double bonds, and the like can also be present in the compounds described herein, and all such stable isomers are contemplated in the present invention. Cis and trans geometric isomers of the compounds of the present invention are described and may be isolated as a mixture of isomers or as separated isomeric forms. All chiral, diastereomeric, racemic forms and all geometric isomeric forms of a structure are intended, unless the specific stereochemistry or isomeric form is specifically indicated.

The term "substituted," as used herein, means that any one or more hydrogens on the designated atom is replaced with a selection from the indicated group, provided that the designated atom's normal valency is not exceeded, and that the substitution results in a stable compound. When a substituent is keto (i.e. =O), then 2 hydrogens on the atom are replaced.

When any variable (e.g. R^{5b}) occurs more than one time in any constituent or formula for a compound, its definition at each occurrence is independent of its definition at every other occurrence. Thus, for example, if a group is shown to be substituted with 0-2 R^{5b} , then said group may optionally be substituted with up to two R^{5b} groups and R^{5b} at each occurrence is selected independently from the definition of R^{5b} . Also, combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

When a bond to a substituent is shown to cross a bond connecting two atoms in a ring, then such substituent may be bonded to any atom on the ring. When a substituent is listed without indicating the atom via which such substituent is bonded to the rest of the compound of a given formula, then such substituent may be bonded via any atom in such substituent. Combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

As used herein, "alkyl" or "alkylene" is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms; for example, "C₁-C₆ alkyl" denotes alkyl

5 having 1, 2, 3, 4, 5, or 6 carbon atoms. Examples of alkyl include, but are not limited to, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, sec-butyl, t-butyl, pentyl, and hexyl. Preferred "alkyl" group, unless otherwise specified, is "C₁-C₄ alkyl". Additionally, unless
10 otherwise specified, "propyl" denotes n-propyl or i-propyl; "butyl" denotes n-butyl, i-butyl, sec-butyl, or t-butyl.

As used herein, "alkenyl" or "alkenylene" is intended to include hydrocarbon chains of either a straight or branched configuration and one or more unsaturated
15 carbon-carbon bonds which may occur in any stable point along the chain. Examples of "C₂-C₆ alkenyl" include, but are not limited to, ethenyl, 1-propenyl, 2-propenyl, 1-butenyl, 2-butenyl, 3-butenyl, 3-methyl-2-butenyl, 2-pentenyl, 3-pentenyl, hexenyl, and the like.

20 As used herein, "alkynyl" or "alkynylene" is intended to include hydrocarbon chains of either a straight or branched configuration and one or more carbon-carbon triple bonds which may occur in any stable point along the chain, such as ethynyl, 1-propynyl, 2-propynyl, 1-butylnyl, 2-
25 butynyl, 3-butylnyl, and the like.

"Alkoxy" or "alkyloxy" represents an alkyl group as defined above with the indicated number of carbon atoms attached through an oxygen bridge. Examples of alkoxy include, but are not limited to, methoxy, ethoxy,
30 n-propoxy, i-propoxy, n-butoxy, s-butoxy, t-butoxy, n-pentoxy, and s-pentoxy. Preferred alkoxy groups are methoxy, ethoxy, n-propoxy, i-propoxy, n-butoxy, s-butoxy, t-butoxy. Similarly, "alkylthio" or "thioalkoxy" is represents an alkyl group as defined above with the
35 indicated number of carbon atoms attached through a sulphur bridge.

"Halo" or "halogen" as used herein refers to fluoro, chloro, bromo, and iodo. Unless otherwise specified,

preferred halo is fluoro and chloro. "Counterion" is used to represent a small, negatively charged species such as chloride, bromide, hydroxide, acetate, sulfate, and the like.

5 "Haloalkyl" is intended to include both branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms, substituted with 1 or more halogen (for example $-C_vF_w$ where $v = 1$ to 3 and $w = 1$ to $(2v+1)$). Examples of haloalkyl include, but
10 are not limited to, trifluoromethyl, trichloromethyl, pentafluoroethyl, pentachloroethyl, 2,2,2-trifluoroethyl, 2,2-difluoroethyl, heptafluoropropyl, and heptachloropropyl. "Haloalkoxy" is intended to mean a haloalkyl group as defined above with the indicated number
15 of carbon atoms attached through an oxygen bridge; for example trifluoromethoxy, pentafluoroethoxy, 2,2,2-trifluoroethoxy, and the like. "Halothioalkoxy" is intended to mean a haloalkyl group as defined above with the indicated number of carbon atoms attached through a
20 sulphur bridge.

"Cycloalkyl" is intended to include saturated ring groups, having the specified number of carbon atoms. For example, "C₃-C₆ cycloalkyl" denotes such as cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl.

25 As used herein, "carbocycle" is intended to mean any stable 3- to 7-membered monocyclic or bicyclic or 7- to 13-membered bicyclic or tricyclic, any of which may be saturated, partially unsaturated, or aromatic. Examples of such carbocycles include, but are not limited to,
30 cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, adamantyl, cyclooctyl, [3.3.0]bicyclooctane, [4.3.0]bicyclononane, [4.4.0]bicyclodecane (decalin), [2.2.2]bicyclooctane, fluorenyl, phenyl, naphthyl, indanyl, adamantyl, or tetrahydronaphthyl (tetralin). Preferred
35 "carbocycle" are cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl.

As used herein, "carbocyclic moiety" is intended to mean any stable 3- to 8-membered monocyclic ring of carbon

atoms, any of which may be saturated or partially unsaturated. Additionally, the 3 to 8 membered monocyclic ring of carbon atoms may be contain a heteroatom selected from oxygen, sulphur, or nitrogen, wherein a carbon atom of the ring has been substituted for the heteroatom. Examples of such carbocycles include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclopent-3-enyl, cyclohex-3-enyl, tetrahydrofurnayl, pyranyl, pyrrolidinyl, and piperidinyl.

Preferred examples of a "carbocyclic moiety" are cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl.

As used herein, the term "heterocycle" or "heterocyclic ring" is intended to mean a stable 5- to 7-membered monocyclic or bicyclic or 7- to 14-membered bicyclic heterocyclic ring which is saturated partially unsaturated or unsaturated (aromatic), and which consists of carbon atoms and 1, 2, 3 or 4 heteroatoms independently selected from the group consisting of N, O and S and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring. The nitrogen and sulfur heteroatoms may optionally be oxidized. The heterocyclic ring may be attached to its pendant group at any heteroatom or carbon atom which results in a stable structure. The heterocyclic rings described herein may be substituted on carbon or on a nitrogen atom if the resulting compound is stable. If specifically noted, a nitrogen in the heterocycle may optionally be quaternized. It is preferred that when the total number of S and O atoms in the heterocycle exceeds 1, then these heteroatoms are not adjacent to one another. It is preferred that the total number of S and O atoms in the heterocycle is not more than 1.

Examples of heterocycles include, but are not limited to, 1H-indazole, 2-pyrrolidonyl, 2H,6H-1,5,2-dithiazinyl, 2H-pyrrolyl, 3H-indolyl, 4-piperidonyl, 4aH-carbazole, 4H-quinolizinyl, 6H-1,2,5-thiadiazinyl, acridinyl, azocinyl, benzimidazolyl, benzofuranyl, benzothiofuranyl, benzothiophenyl, benzoxazolyl, benzthiazolyl,

benztriazolyl, benztetrazolyl, benzisoxazolyl,
benzisothiazolyl, benzimidazolyl, carbazolyl,
4aH-carbazolyl, b-carbolinyl, chromanyl, chromenyl,
cinnolinyl, decahydroquinolinyl, 2H,6H-1,5,2-dithiazinyl,
5 dihydrofuro[2,3-b]tetrahydrofuran, furanyl, furazanyl,
imidazolidinyl, imidazolyl, imidazolyl, 1H-indazolyl,
indolenyl, indolinyl, indoliziny, indolyl,
isobenzofuranyl, isochromanyl, isoindazolyl, isoindolinyl,
isoindolyl, isoquinolinyl, isothiazolyl, isoxazolyl,
10 morpholinyl, naphthyridinyl, octahydroisoquinolinyl,
oxadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl,
1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, oxazolidinyl,
oxazolyl, oxazolidinylperimidinyl, phenanthridinyl,
phenanthrolinyl, phenarsazinyl, phenazinyl, phenothiazinyl,
15 phenoxathiinyl, phenoxazinyl, phthalazinyl, piperazinyl,
piperidinyl, pteridinyl, piperidonyl, 4-piperidonyl,
pteridinyl, purinyl, pyranal, pyrazinyl, pyrazolidinyl,
pyrazolinyl, pyrazolyl, pyridazinyl, pyridooxazole,
pyridoimidazole, pyridothiazole, pyridinyl, pyridyl,
20 pyrimidinyl, pyrrolidinyl, pyrrolinyl, pyrrolyl,
quinazolinyl, quinolinyl, 4H-quinoliziny, quinoxalinyl,
quinuclidinyl, carbolinyl, tetrahydrofuranyl,
tetrahydroisoquinolinyl, tetrahydroquinolinyl,
6H-1,2,5-thiadiazinyl, 1,2,3-thiadiazolyl,
25 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl,
thianthrenyl, thiazolyl, thienyl, thienothiazolyl,
thienooxazolyl, thienoimidazolyl, thiophenyl, triazinyl,
1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,5-triazolyl,
1,3,4-triazolyl, xanthenyl. Preferred 5 to 10 membered
30 heterocycles include, but are not limited to, pyridinyl,
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,
pyrrolyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl,
tetrazolyl, benzofuranyl, benzothiofuranyl, indolyl,
benzimidazolyl, 1H-indazolyl, oxazolidinyl, isoxazolidinyl,
35 benzotriazolyl, benzisoxazolyl, oxindolyl, benzoxazolinyl,
quinolinyl, and isoquinolinyl. Preferred 5 to 6 membered
heterocycles include, but are not limited to, pyridinyl,
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,


pyrrolyl, piperazinyll, piperidinyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, tetrazolyl; more preferred 5 to 6 membered heterocycles include, but are not limited to,

pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, piperazinyll, piperidinyl, pyrazolyl, imidazolyl, and tetrazolyl. Also included are fused ring and spiro compounds containing, for example, the above heterocycles.

As used herein, the term "aryl", "C₆-C₁₀ aryl" or aromatic residue, is intended to mean an aromatic moiety containing the specified number of carbon atoms; for example phenyl, pyridinyl or naphthyl. Preferred "aryl" is phenyl. Unless otherwise specified, "aryl" may be unsubstituted or substituted with 0 to 3 groups selected from H, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, butoxy, amino, hydroxy, Cl, F, Br, I, CF₃, SCH₃, S(O)CH₃, SO₂CH₃, -N(CH₃)₂, N(CH₃)H, CN, NO₂, OCF₃, C(=O)CH₃, CO₂H, CO₂CH₃, or C₁-C₄ haloalkyl.

As used herein, the term "heteroaryl fused radical" is intended to denote a 5 or 6 membered aromatic ring comprising carbon atoms and one or two heteroatoms selected from nitrogen, sulphur and oxygen. The 5 or 6 membered ring is fused to two adjacent atoms of a second ring wherein the second ring is a "carbocyclic moiety" or ring B as defined above. Examples of a "heteroaryl fused radical" are furanyl, imidazolyl, isoxazolyl, oxazolyl, pyrrolyl, thiophenyl, thiazolyl, isothiozalyll, pyridyl, pyridazinyl, pyrimidinyl, and pyrazinyl.




The phrase "additional lactam carbons", as used herein, is intended to denote the number of optional carbon atoms in the lactam ring B of Formula (I). Formula (I''):




(I'')


represents the lactam ring B of Formula (I). Additional lactam carbons are carbons in lactam ring B other than the

carbons numbered 2 and 3 in the backbone of the formula. The additional lactam carbons may be optionally replaced by a heteroatom selected from oxygen, nitrogen and sulfur. Lactam ring B contains 1, 2, 3, 4, 5, 6 or 7 optional
 5 carbons, wherein one optional carbon may optionally be replaced by a heteroatom, such that the total number of members of lactam ring B, including atoms numbered 1, 2 and 3 in the backbone, does not exceed 10. It is preferred that the total number of atoms of lactam ring B is 6, 7 or
 10 8; it is more preferred that the total number of atoms of lactam ring B is seven. It is further understood that lactam ring B may optionally be unsaturated or partially unsaturated (i.e. two adjacent atoms in the ring form a double bond) wherein the backbone of lactam ring B may
 15 contain one, two or three double bonds. Examples of lactam ring B include:







B12




B13



B14




B15




B16

but are not intended to limit the invention. Preferred examples of lactam ring B are B1, B2, B5, B6, B8, B9, B13, and B16; more preferred examples of lactam ring B are B1, B6, B8, B9, and B13. Preferred examples of substituent R¹⁰ or R¹¹ on lactam B are hydrogen, methyl, ethyl, phenyl, benzyl, phenethyl, 4-fluorophenyl, 4-chlorophenyl, 4-methylphenyl, 4-CF₃-phenyl, (4-fluorophenyl)methyl, (4-chlorophenyl)methyl, (4-methylphenyl)methyl, (4-CF₃-phenyl)methyl, (4-fluorophenyl)ethyl, (4-chlorophenyl)ethyl, (4-methylphenyl)ethyl, (4-CF₃-phenyl)ethyl, and 2-, 3-, and 4-pyridinyl. More preferred examples of substituent R¹⁰ or R¹¹ on lactam B are methyl, ethyl, phenyl, 4-fluorophenyl, 4-chlorophenyl, 4-CF₃-phenyl, (4-fluorophenyl)methyl, (4-chlorophenyl)methyl, (4-CF₃-phenyl)methyl, and 2-, 3-, and 4-pyridinyl. Preferred examples of R¹³ on lactam B are F, Cl, OH, methyl, ethyl, methoxy, and trifluoromethyl.

The compounds herein described may have asymmetric centers. One enantiomer of a compound of Formula (I) may display superior biological activity over the opposite enantiomer. For example carbon 3 of lactam ring B Formula (I") may exist in either an S or R configuration. Thus, an R or S configuration at carbon 3 in Formula (I") is considered part of the invention. An example of such configuration includes, the S isomer:



and the R isomer:



5

but is not intended to be limited to this example of ring B. When required, separation of the racemic material can be achieved by methods known in the art. Additionally, the carbon atoms to which R³ and R⁵ are attached may describe chiral carbons which may display superior biological activity over the opposite enantiomer. For example, where and R⁵ is not H, then the configuration of the two centers may be described as (2R,3R), (2R,3S), (2S,3R), or (2S,3S). All configurations are considered part of the invention.

The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

As used herein, "pharmaceutically acceptable salts" refer to derivatives of the disclosed compounds wherein the parent compound is modified by making acid or base salts thereof. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like. The pharmaceutically acceptable salts include the

conventional non-toxic salts or the quaternary ammonium salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include those derived from
5 inorganic acids such as hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, nitric and the like; and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pantoic, maleic, hydroxymaleic,
10 phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic, and the like.

The pharmaceutically acceptable salts of the present
15 invention can be synthesized from the parent compound which contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in
20 water or in an organic solvent, or in a mixture of the two; generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in *Remington's Pharmaceutical Sciences*, 17th ed., Mack Publishing Company, Easton, PA,
25 1985, p. 1418, the disclosure of which is hereby incorporated by reference.

"Prodrugs" are intended to include any covalently bonded carriers which release the active parent drug according to Formula (I) *in vivo* when such prodrug is
30 administered to a mammalian subject. Prodrugs of a compound of Formula (I) are prepared by modifying functional groups present in the compound in such a way that the modifications are cleaved, either in routine manipulation or *in vivo*, to the parent compound. Prodrugs
35 include compounds of Formula (I) wherein a hydroxy, amino, or sulfhydryl group is bonded to any group that, when the prodrug or compound of Formula (I) is administered to a mammalian subject, cleaves to form a free hydroxyl, free

amino, or free sulfhydryl group, respectively. Examples of prodrugs include, but are not limited to, acetate, formate and benzoate derivatives of alcohol and amine functional groups in the compounds of Formula (I), and the like.

5 "Stable compound" and "stable structure" are meant to indicate a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic agent.

10

SYNTHESIS


The compounds of the present invention can be prepared in a number of ways well known to one skilled in the art of organic synthesis. The compounds of the present invention
15 can be synthesized using the methods described below, together with synthetic methods known in the art of synthetic organic chemistry, or variations thereon as appreciated by those skilled in the art. Preferred methods include, but are not limited to, those described below.

20 All references cited herein are hereby incorporated in their entirety herein by reference.

The novel compounds of this invention may be prepared using the reactions and techniques described in this section. The reactions are performed in solvents
25 appropriate to the reagents and materials employed and are suitable for the transformations being effected. Also, in the description of the synthetic methods described below, it is to be understood that all proposed reaction conditions, including choice of solvent, reaction
30 atmosphere, reaction temperature, duration of the experiment and workup procedures, are chosen to be the conditions standard for that reaction, which should be readily recognized by one skilled in the art. It is
understood by one skilled in the art of organic synthesis
35 that the functionality present on various portions of the molecule must be compatible with the reagents and reactions proposed. Such restrictions to the substituents which are compatible with the reaction conditions will be readily

apparent to one skilled in the art and alternate methods must then be used.

Methods for the synthesis of succinylamino lactams are known in the art and are disclosed in a number of references including PCT publication number WO 96/29313, which is hereby incorporated by reference.





Compounds of the present invention can be prepared by an amino acid coupling procedure. An example of the synthetic method employed to prepare compounds of the present invention is illustrated in the amino acid coupling of succinate derivative XI and lactam XII to give a compound of Formula II as shown in Scheme 1. It is understood that ring C represents variables R^3 and R^{3a} of Formula (I). The cyclic succinate derivative XI can be prepared from a mono-ester succinate derivative X which can be prepared from the corresponding diester or acid.

Examples of succinates wherein ring C is a carbocyclic or heterocyclic system are well known in the literature. For example, a dimethyl succinate having a 3-membered cyclopropyl ring C can be formed by a thermal or photolytic decomposition of a methyl 3(carbomethoxymethyl)-1-pyrazoline-3-carboxylate. See Bull. Soc. Chim. Fr. (1971), (6), 2290-5. A succinic acid derivative wherein ring C is a 4-membered cyclobutyl group can be formed by the method published in US 3,828,025. A succinic acid derivative

wherein ring C is a 5-membered cyclopentyl group can be formed using the methods described in Le Moal, H. et al., Bull. Soc. Chim. Fr., 1964, 579-584; Borenstein, M.R., et al., Heterocycles, 22, 1984, 2433-2438. Other examples of derivatives of succinate X wherein ring C is a five-membered cyclopentyl group or a 6-membered cyclohexyl group have been employed as matrix metalloproteinase inhibitors. See Bioorg. Med. Chem. Lett. (1998), 8(12), 1443-1448; Robinson, R.P., et al., Bioorg. Med. Chem. Lett. (1996), 6(14), 1719-1724. For the preparation of a succinic acid wherein ring C is an oxygen containing 3-membered oxirane see Kirshenbaum, K.S., Sharpless, K. B., J. Org. Chem. (1985), 50(11), 1979-82. For examples of succinate derivatives wherein ring C is a 5- or 6-membered heterocycle ring see Olivero, S., Dunach, E., Eur. J. Org. Chem. (1999), (8), 1885-1891; Eckardt et al. Helv. Chim. Acta, 55, 1972, 2432, 2433, 2434, 2438; Sandoz Ltd., NL 6409801 1963, Chem. Abstract., 63, 1965, 8324d; and Rice, L.M., et al., J. Med. Chem., 6, 1963, 388-402. It is understood that these references are only illustrative of the availability of some carbocyclic and heterocyclic succinates, however numerous references are known in literature which provide preparations of other substituted carbocyclic and heterocyclic succinates and their derivatives.

Scheme 2 illustrates one method for the introduction of a substitution on a carbon adjacent to the cyclic group in succinate IX via a deprotonation followed by standard alkylation procedures known to one skilled in the art. Treatment of IX with a base followed by addition of an R⁵-LG, wherein LG is a leaving group such as a halide, mesylate, triflate or a tosylate, and subsequent deprotection of the benzyl group by hydrogenation employing, for example, H₂ and Pd/C, would give the desired succinate X.


Scheme 2

Scheme 2a


An example of a general method whereby diesters of structure IX can be obtained from cyclic esters XXX is shown in Scheme 2a. Deprotonation of such esters with, for example LDA or lithium hexamethyldisilazide, followed by reaction of the resulting ester enolate with allyl bromide provides allyl esters XXXI, which may be oxidized using ruthenium peroxide in the presence of sodium periodate to give free acids XXXII. If desired, esterification may be carried out using e.g. benzyl bromide in the presence of potassium carbonate.

Succinate acids such as XXXV with defined stereochemistry may be prepared from cyclic mono-acids XXXIII by use of a chiral auxiliary such as an oxazolidinone, as shown in Scheme 2b. Thus, acid XXXIII may be converted to the oxazolidinone XXXIV and subjected to the Evans stereospecific alkylation sequence to provide, after removal of the auxiliary, acid esters XXXV. Suitable alkylating agents include alkyl, allyl, propenyl or benzyl iodides or triflates, as is known to those skilled in the art. Use of the appropriate stereochemistry in the chiral auxiliary can provide substituted cyclic succinates of either absolute configuration.

Scheme 2b



Scheme 3



5


An example of a class of compounds compound which can be prepared according to the general methods described above is shown in Scheme 3. In step 1, cyclopentyl succinate X-a is converted to cyclopentyl succinamide XI-a. The subsequent amino acid coupling between succinamide XI-a and a benzodiazepine 11 under standard coupling conditions known to one skilled in the art would give the product III.

In the preparation of a compound of Formula I, in some cases it may be desirable to perform the coupling of an amino lactam and a succinic acid derivative prior to the amidation of the succinic acid compound. In Scheme 4 the coupling of a benzodiazepine 11 and the succinic acid derivative 10 is performed first to give the coupled product 12. Subsequent conversion of the ester function to


20

an amide can be done by a deprotection step followed by an amino acid coupling step using standard coupling conditions to give a compound 13.

5

Scheme 4Scheme 4a


10



A variation of the coupling procedure of Scheme 4 useful in the synthesis of compounds of the present invention is illustrated in Scheme 4a, and involves


conversion of the acid ester 10 to the corresponding HOBt ester using standard procedures, such as EDC/HOBt. In a separate step, the activated ester XL is allowed to react with a lactam amine XLI in a suitable solvent, such as DMF, preferably with warming of the reaction mixture to 40-100 °C. As will be recognized by those skilled in the art, a variety of procedures for the synthesis of amides from carboxylic acids are known (see, for example, Peptide Synthesis Protocols, ed. by M. W. Pennington and B. M. Dunn, Methods in Molecular Biology, Vol. 35, Humana Press, 1994; and Comprehensive Organic Functional Group Transformations, ed. by A. R. Katritzky, O. Meth-Cohn, C. W. Rees, Volume 5, pp 274-281 [P. D. Bailey, I. D. Collier and K. M. Morgan, Amides], Pergamon, 1995), and the skilled practitioner will adjust the methods, reagents and conditions to the example at hand.

The methods illustrated above may be modified to prepare cyclic succinoyl lactams where R5 and R5a comprise a cyclic group as shown in Scheme 4b. Thus the free acid of succinate esters prepared using the methods of Schemes 2, 2a or 2b may be suitably protected and the ester deprotected. Coupling of the free acid with an aminolactam may be carried out as described, and the remaining ester group converted to the desired amide.



Compounds of the present invention in which R3 and R3a, and R5 and R5a comprise two cyclic groups may be prepared from the corresponding cyclic succinates as shown in Scheme 4c. Bis(cyclic) succinates useful in the preparation of intermediates represented by xx are available using the methods of Overberger et al. (J. Org. Chem. 1955, 20, 1717-1720) and Belletire et al. (Tet. Lett. 1984, 25, 5969-5972).

10


Scheme 4c

Methods for the synthesis of lactams XI of Scheme 1 as contemplated by the present invention in lactam ring B of Formula (I), including amino benzodiazepinones, dibenzo azepinones and other related heterocycles, are known in the art and are disclosed in a number of references including PCT publication number WO 98/28268, WO 99/66934, and WO 00/07995, which are hereby incorporated by reference. Additional references include Bock, et al, J. Org. Chem., 1987, 52, 3232-3239; Sherrill et al, J. Org. Chem., 1995, 60, 730-734; and Walsh, D. A., Synthesis, September 1980, p.677; and Brown, et al., Tetrahedron Letters, 1971, 8, 667-670.

An example of an L- α -amino- β -thio- ϵ -caprolactam, as shown in Scheme 5, where ring B is the amino lactam of XIII and J is a sulfur atom has been reported in the literature. See S. A. Ahmed et al, FEBS Letters, (1984), vol. 174, pages 76-9. One skilled in the art can extend this

methodology to the synthesis of β -amino and oxygen containing rings by analogy. The sulfur-containing molecules can also be oxidized to the sulfoxide and sulfone by methods known to one skilled in the art.


5



An approach to preparing representative compounds of Formula (I) is illustrated for caprolactam 20 in Scheme 6. The lactam nitrogen of intermediate 15 can be alkylated by generating the anion with bases, such as LDA, lithium bis(trimethylsilyl)amide or sodium hydride, in solvents such as THF, with or without cosolvents such as DMPU or HMPA and reacting this with a variety of groups containing leaving groups (LG) for example, bromide, iodide, mesylate or tosylate. Alkylating agents such as α -bromo amides, ketones and acids, if not commercially available, can be prepared by a number of literature methods including halogenation of amino acids by diazotization. Other suitable alkylating agents such as alkyl, allylic and benzylic halides can be formed from a variety of precursors such as free-radical addition of halides or activation of alcohols, and other chemistries known to one skilled in the art. For discussion of these types of reactions, see Carey, F.A. and Sundberg, R. J., Advanced Organic Chemistry, Part A, New York: Plenum Press, 1990, pages 304-305, 342-347 and 695-698.

The *N*-Boc protecting group of caprolactam **16** can be removed by any number of methods well known in the literature, for example TFA in methylene chloride, to give the intermediate **17**. The amine **17** can be coupled to an
 5 appropriately substituted carboxylic acid X-a, acid

Scheme 6




chloride or other activated acid derivative by methods well described in the literature for making amide bonds, for
 15 example, TBTU in DMF with a base such as NMM, to give the elaborated caprolactam **18**. Optionally, caprolactam **18** can be alkylated using standard bases, such as LDA, NaH, or NaHMDS, to deprotonate the amide hydrogen followed by addition of an alkylating agent with an appropriate leaving
 20 group, such as halide, mesylate, or triflate in an appropriate solvent to provide an *N*-R⁶ alkylated product of caprolactam **18**. The *t*-butyl carboxyl protecting group of the *N*-R⁶ alkylated product of caprolactam **18** can be removed, for example, by treatment with TFA in methylene
 25 chloride to give a carboxylic acid **19**.

The final product 20 can be prepared by treating an activated carboxylic acid derivative of 19 with an appropriately substituted amine HNR^1R^2 . For instance, activation of the carboxylic acid with HATU (O-(7-azabenzotriazol-1-yl)-1,1,3,3,-tetramethyluronium hexafluorophosphate) or PyBOP (benzotriazole-1-yl-oxy-tris-pyrrolidino-phosphonium hexafluorophosphate) or other coupling agents known to those skilled in the art allows condensation with ammonia to form primary amides.

Similarly, condensation of the activated acid with hydroxylamine hydrochloride provides the hydroxamic acid, or reaction with a primary or secondary amine provides the substituted amine derivative. For additional acylation reactions see, for example, Carey, F.A. and Sundberg, R. J., Advanced Organic Chemistry, Part A, New York: Plenum Press, 1990, pages 475-479.

A different synthetic route to compounds of the present invention is shown in Scheme 7, step 1-5. In the method of Scheme 7, cyclization of two succinate substituents is carried out after coupling of the aminolactam and a suitable succinic acid derivative. In step 1, the amino lactam XII is coupled to an appropriately substituted succinate derivative XIV (n is 1, 2 or 3) or acid chloride by methods well described in the literature for making amide bonds, for example, TBTU in DMF with a base, for example, NMM to give the elaborated compound XV.


Scheme 7

The protecting group of the carboxyl group can be removed using standard deprotection methods to provide compound XVI. A subsequent second amino acid coupling with an amine NHR^1R^2 provides compound XVII which undergoes a ring-closing-metathesis to form a compound of formula IV in the presence of Ruthenium complexes using the method described by Grubbs and coworkers, J. Am. Chem. Soc., 114, 7324 (1992). Reduction of the cycloalkene of IV to a

compound of formula V can be done using hydrogen or a hydrogen transfer reagent with Palladium as a catalyst or other reduction methods well known in the art.

The compound XIV of Scheme 7 can be prepared by a number of known procedures. See D. A. Evans et al, *Org. Synth.* **86**, p83 (1990) and P. Becket, M. J. Crimmin, M. H. Davis, Z. Spavold, Synlett, (1993), 137-138. The preparation of diastereomerically pure succinate XXI is outlined in Scheme 8 where acylation of an oxazolidinone with an acylating agent such as an acid chloride $\text{ClCOCH}_2\text{R}^5$ provides structure XVIII. Alkylation of XVIII with a $\text{BrCH}_2\text{CO}_2\text{t-Bu}$ provides XIX followed by cleavage of the chiral auxiliary to give carboxylic acid XX. Subsequent alkylations of XX provides a variety of disubstituted succinate XXI which can be further alkylated to give a compound of formula XIV.

Scheme 8




20

An example of diastereomerically pure succinate derivative XIV employing a method of Scheme 8 in its preparation is outlined in Scheme 9, adapted from P. Becket, M. J. Crimmin, M. H. Davis, Z. Spavold, Synlett, (1993), 137-138 incorporated herein by reference. This reference provides the synthesis shown below to obtain compound 24. Additional methods useful for the

25


preparation of succinate derivatives are known by those skilled in the art. Such references include McClure and Axt, Bioorganic & Medicinal Chemistry Letters, 8 (1998) 143-146; Jacobson and Reddy, Tetrahedron Letters, Vol 37, No. 46, 8263-8266 (1996); Pratt et al., SYNLETT, May 1998, p. 531; WO 97/18207; and WO 98/51665. The synthetic disclosures of WO97/18207 and WO 98/51665 are hereby incorporated by reference. A further alkylation of disubstituted succinates such as XXI and 24 provides intermediates such as 25 useful as substrates, after esterification, for cyclization reactions known to one skilled in the art, such as ring closing metathesis (RCM) reactions using Grubbs' catalyst as illustrated in Scheme 9. It will be appreciated by those skilled in the art that the analogous preparation of other cyclization substrates and the use of alternative ring forming methodologies will provide access to carbo- and heterocyclic analogs of intermediates 26 and 27. Such strategies include oxidative olefin cleavage using, for example, ozonolysis, followed by reduction to the corresponding diols, mono-activation with tosyl chloride, and cyclization to oxygen containing heterocycles using base. Treatment of dialdehydes with primary amines in the presence of sodium borohydride may be used to provide analogous nitrogen heterocyclic intermediates useful in the preparation of nitrogen heterocycles of Formula (I). The use of cycloalkylidene succinates such as 27 in the preparation of compounds of Formula (I) is illustrated in Schemes 7 and 10.



5


The preparation of compounds 32 and 33 using the methods of Scheme 7 is shown in Scheme 10. The succinate product 29 is obtained from an amino acid coupling of a diallyl succinate 28 with a benzodiazepine 11 using a standard coupling procedure (HATU, DIEA, DMF). The carboxyl protecting group BOC is removed in TFA/CH₂Cl₂ to give a carboxylic acid 30. A second amino acid coupling of carboxylic acid 30 with ammonia in the presence of HATU and DIEA in DMF provides a diallyl succinate 31. A ring-closing-metathesis using a catalytic amount of Cl₂Ru(PCy₃)₂(CHC₆H₅) as the metal carbene compound gives the cyclized product 32. Compound 33 is obtained from compound 32 by a hydrogen transfer reduction with Pd(OH)₂/C and 1,4-cyclohexadiene in methanol.

20



- 5 The amide hydrogen of a compound of formula VI can be deprotonated using standard bases, for example LDA, NaH, or NaHMDS, followed by addition of an alkylating agent $\text{R}^6\text{-LG}$ wherein LG is an appropriate leaving group, for example halide, mesylate, or triflate, in an appropriate solvent to
- 10 provide a compound of formula VII, see Scheme 11. A similar synthetic sequence may be applied to protected intermediate XV, which may then be converted to VII using the methods described above.

Scheme 11



5 All references cited herein, are hereby incorporated by reference in their entirety unless otherwise stated.

Examples

Chemical abbreviations used in the Examples are defined as follows: "DMPU" for 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidone, "TBTU" for O-(1H-benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate, "BOP" for benzotriazol-1-yloxytris-(dimethylamino)-phosphonium hexafluorophosphate, "EDC" for 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride, "HOBT" for 1-hydroxybenzotriazole hydrate, "TEA" for triethyl amine, "LiHMDS" for lithium bis(trimethylsilyl)amide, "HMPA" for hexamethylphosphoramide, "LDA" for lithium diisopropylamide, "DCC" for 1,3-dicyclohexylcarbodiimide, "PyBoP" for benzotriazole-1-yl-oxy-tris-pyrrolidino-phosphonium hexafluorophosphate, and "HATU" for O-(7-azabenzotriazol-1-yl)-N, N, N', N'-tetramethyluronium hexafluorophosphate. "HPLC" is an abbreviation used herein for high pressure liquid chromatography. Reverse-phase HPLC can be carried out using a Vydac C-18 column with gradient elution from 10% to 100 % buffer B in buffer A (buffer A: water containing 0.1% trifluoroacetic acid, buffer B: 10% water, 90% acetonitrile containing 0.1% trifluoroacetic acid). If necessary, organic layers can be dried over sodium sulfate unless otherwise indicated. However, unless otherwise indicated, the following conditions are generally applicable.

Melting points were determined on a Mel-Temp II apparatus and are uncorrected. IR spectra were obtained on a single-beam Perkin-Elmer Spectrum 1000 FT-IR spectrometer using 4 accumulations at a resolution of 4.00 cm⁻¹ on samples prepared in a pressed disc of KBr or as a film on NaCl plates. Proton NMR spectra (500 MHz, referenced to tetramethylsilane) were obtained on a Bruker AMX 500 spectrometer or on a Bruker AC 300 spectrometer (300MHz, referenced to tetramethylsilane). Mass spectra were obtained on a Shimadzu QP-5000 mass spectrometer (CI or EI), a Perkin Elmer Sciex 100 atmospheric pressure ionization (API) mass spectrometer or a Finnigan LCQ Duo LCMS ion trap electrospray ionization (ESI) mass spectrometer. HPLC analyses were obtained using a Rainin Dynamax C₁₈ column with UV detection at 223 nm using a standard solvent gradient program as follows:


HPLC solvent conditions: Samples were dissolved in methanol (1 mg/mL) and run using the following gradient program with a solvent flow rate of 1.0 mL/min.

<u>Time (min)</u>	<u>Acetonitrile</u>	<u>H₂O</u>
	<u>(0.05% TFA)</u>	<u>(0.05% TFA)</u>
Initial	10	90
20.0	90	10
20-30	90	10

Elemental analyses were performed by Quantitative Technologies, Inc. (Whitehouse, NJ).

Example 1

1-[(1R)-3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclopent-3-enecarboxylic amide



Following the disclosure of Scheme 10:

- 5 **Step 1:** Preparation of 2-Allyl-2-[3-methyl-1-(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-ylcarbamoyl)-butyl]-pent-4-enoic acid tert-butyl ester 29. Compound 28 (1.4g, 4.5mmol) in 50ml DMF was added HATU (O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium
10 hexafluorophosphate) (2.2g, 5.8mmol) followed by DIEA (N,N-diisopropylethylamine) (800ul, 4.6mmol). The solution was stirred at RT for one hour. A mixture of compound 11 (2.3g, 4.6mmol) and DIEA (920ul, 5.2mmol) in 50ml DMF was added to the above solution over several minutes. The
15 resulting solution was stirred at ambient temperature overnight, then quenched with 20ml water. Removal of volatiles gave a yellow oil which was taken up in ethyl acetate/water (1:1). The organic layer was washed with water twice, then brine, dried over Na₂SO₄, filtered, and
20 concentrated. The residue was purified by silica gel chromatography with hexane/ethyl acetate (8:2) to give 300mg of product 29 (15%). ¹H NMR (CDCl₃) 0.96-0.89 (dd, 6H), 1.28-1.22 (m, 1H), 1.50 (s, 9H), 1.65-1.50 (m, 1H), 2.00-1.85 (m, 1H), 3.80-2.40 (m, 5H), 3.45 (s, 3H), 5.31-
25 5.00 (m, 5H), 5.59-5.57 (d, 1H), 6.00-5.85 (m, 2H), 7.40-7.20 (m, 9H). ESI ES+ = 558.3 (M+1).

Preparation of 2-Diallyl-3-isobutyl-succinic acid 1-tert-butyl ester 28. To a solution of 2-allyl-3-isobutyl-succinic acid 1-tert-butyl ester (24, 3.1g, 11.5mmol) in
30 50ml THF at -78C was added 0.2M LDA in THF (29mmol, 2.5 eq). After 1hr, a solution of allyl bromide (2.1g, 17.4mmol, 1.5 eq) in 30ml THF was added slowly. The mixture

was allowed to warm to ambient temperature overnight. The reaction was quenched with 30ml of methanol, and the solvent was removed under reduced pressure to provide a yellow oil, which was taken up in ethyl acetate. The organic layer was washed twice with 1.0N citric acid and then with brine, then dried over anhydrous Na₂SO₄.

Filtration and removal of solvent provided the title compound. NMR analysis indicated that the crude product contained a mixture of product 28 (>80%) and starting material 24 (<20%). The mixture was utilized in Step 1 without further purification. The ¹H NMR signals of 28 are overlapped with those of 24. ¹³C NMR for 2: (CDCl₃) 21.2, 23.9, 26.6, 27.9, 36.4, 37.8, 37.9, 48.9, 50.6, 81.5, 118.5, 118.6, 133.3, 134.0, 173.1, 180.0. ESI ES- = 619.5 (2M-1).

Step 2: Preparation of 2-Allyl-2-[3-methyl-1-(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-ylcarbamoyl)-butyl]-pent-4-enoic acid 30. A solution of compound 29 (300mg, 0.54mmol) in 50ml TFA/CH₂Cl₂ (1:1) was stirred for 2hr at ambient temperature, then concentrated under vacuum. Two cycles of toluene addition and removal of volatiles gave 250mg of 30 as a white solid (92%) which was used in the next step without purification. ESI ES+ = 502.3 (M +1)

Step 3: Preparation of 2,2-Diallyl-3-isobutyl-N4-(1-methyl-2-oxo-5-phenyl-2,3-dihydro-1H-benzo[e][1,4]diazepin-3-yl)-succinamide 31. To a solution of 30 (250mg, 0.5mmol) in 30ml DMF was added HATU (250mg, 0.66mmol) and DIEA (400 uL, 2.2 mmol). After stirring for 2hrs, ammonia gas was bubbled into the solution for 5 minutes using a glass inlet tube. The mixture was then stirred at ambient temperature overnight. Water was added (10 mL) and the solvent was removed under vacuum to give a yellow oil which was taken up in ethyl acetate and water (1:1). The organic layer was washed twice with water and then brine, and dried over Na₂SO₄. After filtration and concentration, the crude


product was purified using silica gel chromatography (CH₂Cl₂/methanol 10:0.5) to give compound **31** (120 mg) as a white solid (48%). ¹H NMR (CDCl₃) 0.95-0.88 (dd, 6H), 1.32-1.28 (m, 1H), 1.64-1.50 (m, 1H), 1.90-1.80 (m, 1H), 2.35-2.00 (m, 1H), 2.80-2.54 (m, 4H), 3.48 (s, 3H), 5.38-5.14 (m, 5H), 5.54-5.51 (d, 1H), 5.90-5.78 (m, 2H), 7.65-7.25 (m, 9H), 8.52 (s, 1H). ESI ES+ = 523.2 (M + Na).

Step 4: Preparation of 1-[(1R)-3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclopent-3-enecarboxylic amide **32**. To a solution of **31** (110mg, 0.22mmol) in 20ml of toluene and CH₂Cl₂ (1:1) at 100 degree was added (bis(tricyclohexylphosphine)-benzylidene ruthenium (IV) dichloride) (24mg, 0.029 mmol). The mixture was stirred at 100 degree for 4hrs and another 24mg Grubbs' catalyst was added. The mixture was then stirred at 100 degree for another 4 hrs. Then the solution was cooled to RT, after which 100 mg of charcoal was added. The mixture was filtered through celite to provide a slightly yellow solution, then concentrated. The residue was purified in silica gel chromatography with CH₂Cl₂/methanol (10:0.5) to give 48mg of **32** (46%). ¹H NMR (CDCl₃) 0.97-0.90 (dd, 6H), 1.30-1.25 (m, 1H), 1.65-1.57 (m, 1H), 2.00-1.85 (m, 1H), 2.40-2.30 (d, 1H), 2.80-2.65 (m, 2H), 3.00-2.80 (d, 1H), 3.40-3.20 (m, 1H), 3.47 (s, 3H), 5.40-5.20 (s, 1H), 5.52-5.49 (d, 1H), 5.80-5.71 (m, 2H), 7.63-7.20 (m, 9H), 7.90 (s, 1H). API AP+ 473.1 (M+1).

Example 1a

Synthesis of cyclic succinate intermediate 6 (Scheme 12).

Scheme 12



Step 1: Preparation of diallylsuccinate mono-acid 2.

A solution 300 mmole of LDA was prepared by adding 120 mL of 2.5M n-BuLi in hexanes to 45ml (320 mmol) DIPA in 200ml THF at $-78^\circ C$ followed by stirring in an ice bath for 30 minutes. This was added to a solution of syn-succinate 1 (34.0g, 126mmol) in 100ml THF at $-78^\circ C$ to give a clear yellow solution, which was stirred at that temperature for one hour. A solution of allyl bromide (21.0g, 170 mmol) in 100ml THF was added to the above solution over 20 minutes, and the resulting yellow solution was stirred and allowed warm to room temperature overnight. The reaction mixture was quenched with 50ml methanol and 50ml water. The solvents were evaporated to give a yellow viscous oil which was taken up in EtOAc (400ml) and 200ml 1.0N HCl. The organic layer was washed with 100ml 1.0N HCl, and brine and dried over sodium sulfate. The solution was concentrated to give 2 as a slightly yellow oil (34.2g, 110.3mmol, 88%). ^{13}C NMR (300MHz, $CDCl_3$) 179.8, 173.0, 134.0, 133.3, 118.7,

118.4, 81.4, 50.6, 48.8, 37.8, 36.3, 27.8, 26.6, 23.8, 21.1.

Step 2: Preparation of diallylsuccinate diester 3

5 To a solution of 2 (34.2g, 110 mmol) in 400ml acetone was added potassium carbonate (28.0g, 200mmol) and benzyl bromide (28.0g, 163mmol). The solution was heated to reflux for 2 hours. Evaporation of the solvent gave a yellow oil which was taken up in EtOAc and water. The organic layer
10 was washed with brine, dried over sodium sulfate and concentrated to give diester 3 (44.5g, 100%). ¹³C NMR (300MHz, CDCl₃) 173.7, 172.8, 134.5, 133.6, 129.0, 128.8, 128.5, 128.4, 128.3, 128.2, 118.5, 118.2, 81.0, 66.2, 50.9, 49.0, 38.5, 37.5, 37.9, 28.0, 26.7, 23.9, 21.3.

15

Step 3: Preparation of cyclic succinate diester 4


To a solution of 3 (3.0g, 7.5mmol) in 700ml solution of methylene chloride-toluene (1:1) was added 200 mg tricyclohexylphosphine[1,3-bis(2,4,6-trimethylphenyl)-4,5-
20 dihydroimidazol-2-ylidene]benzylidene]ruthenium (IV) dichloride 200mg (0.24mmol). The reaction mixture was refluxed for 4hrs at 100°C, then evaporated to give a dark oil which was purified by chromatography on silica gel with EtOAc:hexane (5:95) to give 4 as a yellow oil (1.7g,
25 4.6mmol, 62%). ¹H NMR (300 MHz, CDCl₃) 0.8-0.9 (m, 6H), 0.9-1.1 (m, 1H), 1.4 (s, 9H), 1.6-1.8 (m, 2H), 2.6-3.0 (m, 4H), 5.0-5.2 (m, 2H), 5.4-5.6 (d, 2H), 7.2-7.4 (m, 5H).

Step 4: Preparation of cyclic succinate HOBT ester 6

30 Triethylsilane was added to a solution of palladium acetate (170mg, 0.75mmol) in 10ml of methylene chloride and the resulting mixture was stirred at RT for 30 minutes. Triethylamine (0.2ml, 1.4mmol) was added, followed by 4 (2.8g, 7.5mmol) in 10 ml of methylene chloride. The mixture
35 was stirred at RT overnight, filtered through a short silica gel column, then concentrated to give 5 as a colorless oil which was used without further purification. To a solution of 5 in 100ml methylene chloride was added

triethylamine (2.0ml, 14.3 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide HCl (EDC, 2.2g, 11.5 mmol) and 1-hydroxybenzotriazole hydrate (HOBT, 2.2g, 16.3mmol). The mixture was stirred for 4 hours. The solvents were removed under reduced pressure and the resulting oil was taken up in EtOAc and water. The organic layer was washed with water and brine, then dried over sodium sulfate and concentrated. Purification by flash chromatography on silica gel with EtOAc:hexane (10:90) gave 6 (1.95g, 4.9mmol, 65%). ¹HNMR (300 MHz, CDCl₃) 0.9-1.1 (m, 6H), 1.2-1.4 (m, 1H), 1.8-2.0 (m, 2H), 2.5-3.2 (m, 5H), 5.6-5.8 (m, 2H), 7.3-7.6 (m, 3H), 8.0 (d, 1H). MS: 422.1 (M+Na), 463.2 (M+Na+CH₃CN), 821.4 (2M+Na).

15 Synthesis of Formula I compounds 11 and diastereomer 12
(Scheme 13)



Step 1: Preparation of benzodiazepinesuccinamide 8

A solution of racemic aminobenzodiazepine HBr salt 7 (2.6g, 7.8mmol), 6 (6.0g, 15mmol) and diisopropylethyleamine (DIPEA, 1.4ml, 8.0mmol) in 200ml DMF was heated at 60°C for 3 hours. The solvents were removed under reduced pressure to give a viscous oil which was taken up in EtOAc and water. The organic layer was washed with water and brine, then dried over sodium sulfate. The solvents were evaporated under reduced pressure to give an oil which was purified by flash chromatography (30:70 EtOAc:hexane), giving 8 as a white solid (2.1g, 4.0mmol, 51%). ¹HNMR (300 MHz, CDCl₃) 0.8-1.2 (m, 6H), 1.2-1.4 (m, 1H), 1.4-1.6 (m, 9H), 1.6-2.0 (m, 2H), 2.6-3.0 (m, 5H), 5.4-5.8 (m, 3H), 7.0-7.6 (m, 9H). MS: 516.2 (M+H), 538.2 (M+Na).

Step 2: Preparation of N-methylactam 9

To a solution of 8 (2.1g, 4.0mmol) in 100ml DMF was added iodomethane (2.0g, 14.0mmol) and potassium carbonate (1.0g, 7.2 mmol). The reaction mixture was stirred at RT overnight. The solvents were evaporated under reduced pressure and the residue was taken up in EtOAc and water. The organic layer was washed with water and brine, then dried over sodium sulfate. Evaporation of solvents gave 9 as a white solid which was used without purification. ¹HNMR (300 MHz, CDCl₃) 0.8-1.2 (m, 6H), 1.2-1.4 (m, 1H), 1.4-1.6 (m, 9H), 1.6-2.0 (m, 2H), 2.6-3.0 (m, 5H), 3.4 (s, 3H), 5.4-5.7 (m, 3H), 7.2-7.6 (m, 9H). MS: 530.2 (M+H), 552.2 (M+Na).

Step 3: Preparation of succinamide free acid 10


A solution of 9 in 100ml TFA-methylene chloride (1:1) was stirred for 3 hours. The solvents were removed under reduced pressure, and the resulting oil was dissolved in 50 mL of toluene and concentrated provide 10 as a yellow solid (2.0g, 4.2mmol). MS: 474.2 (M+H), 586.2 (M+CF₃COO-).

Step 4: Synthesis of diastereomeric succinamides 11 and 12 (Scheme 2)

A solution of 10 (2.0g, 4.2mmol), O-(7-azabenzotriazol-1-yl)-N,N,N,N-tetramethyluronium hexafluorophosphate (HATU, 2.0g, 5.3mmol) and DIEPA (1ml, 5.7mmol) in 100ml DMF was treated with ammonium gas for 5 minutes. The reaction mixture was stirred overnight, then quenched with water. The solvents were removed under reduced pressure and the resulting oil was taken up in EtOAc and water. The organic layer was washed with brine and dried over sodium sulfate. Evaporation of solvents provided the crude mixture of diastereomers which were separated by flash chromatography using EtOAc: hexane (1:1). Diastereomer 11 (900mg) eluted first: ¹HNMR (300 MHz, CDCl₃) 0.8-1.0 (dd, 6H), 1.2-1.4 (m, 1H), 1.5-1.7 (m, 1H), 1.8-2.0 (m, 1H), 2.3-2.4 (m, 1H), 2.6-3.0 (m, 3H), 3.2-3.4 (m, 1H), 3.47 (s, 3H), 5.25 (s, 1H), 5.4-5.5 (d, 1H), 5.6-5.8 (m, 2H), 7.2-7.8 (m, 9H), 7.9 (s, 1H). MS: 473.2 (M+H). Diastereomer 12 (680mg) eluted second: ¹HNMR (300 MHz, CDCl₃) 0.9-1.2 (m, 6H), 1.2-1.4 (m, 1H), 1.6-1.8 (m, 1H), 1.9-2.1 (m, 1H), 2.35-2.45 (m, 1H), 2.6-2.9 (m, 3H), 3.2 (m, 1H), 3.5 (s, 3H), 5.4 (s, 1H), 5.5 (d, 1H), 5.6-5.8 (m, 2H), 7.2-7.8 (m, 10H). MS: 473.2 (M+H), 495.1 (M+Na).

Example 2


1-[(1R)-3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclopentanecarboxylic amide



Step 5 : Preparation of 1-[(1R)-3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclopentanecarboxylic amide 33 (Scheme 10). To a

solution of compound **32** (Scheme 10) (45mg, 0.1mmol) in 20ml of methanol was added 400 mg of Pd(OH)₂/C (20% weight, water < 50%), followed by 2ml (21mmol) of 1,4-cyclohexadiene. The solution was heated to gentle reflux at (65°C) for 1hr and then cooled to ambient temperature. The solution was filtered and concentrated to provide a white solid, which was purified using silica gel chromatography with CH₂Cl₂/methanol (10:0.5) to give 30mg of **33** (Scheme 10) as a white solid, 8 (63%). ¹H NMR (CDCl₃) 0.96-0.89 (dd, 6H), 1.50-1.20 (m, 2H), 2.10-1.60 (m, 8H), 2.65-2.42 (m, 2H), 3.48 (s, 3H), 5.25 (s, 1H), 5.52-5.50 (d, 1H), 7.65-7.20 (m, 9H), 8.20 (s, 1H). ESI ES+ = 475.2 (M+1).

15

Example 2aSynthesis of Formula I compound 13 (Scheme 14)


20

To a solution of 11 (200mg, 0.42mmol) in 20 mL of methanol was added 200mg palladium hydroxide (20% on carbon, wet) and 1,4-cyclohexadiene (2ml, 21mmol). The reaction mixture was slowly heated to reflux (65°C), stirred 2 hours, then filtered through celite and evaporated. Flash chromatography using methanol:methylene chloride (5:95) provided 13 as a white solid (128mg, 0.27mmol). ¹H NMR (300 MHz, CDCl₃) 0.8-1.0 (dd, 6H), 1.2-2.2 (m, 10H), 2.4-2.7 (m, 2H), 3.47(s, 3H), 5.3 (s, 1H), 5.5 (d, 1H), 7.2-7.6 (m, 9H), 8.2 (s, 1H). MS: 475.2 (M+H), 497.2 (M+Na). The corresponding diastereomer epimeric at the benzodiazepine


carbon was prepared using the same procedure beginning with
12. ¹HNMR (300 MHz, CDCl₃) 0.9-1.1 (m, 6H), 1.2-2.0 (m,
10H), 2.4-2.7 (m, 2H), 3.5 (s, 3H), 5.3 (s, 1H), 5.5 (d,
1H), 7.2-7.7 (m, 9H), 8.2 (s, 1H). MS: 475.2 (M+H), 497.2
5 (M+Na).

Example 3

1-[(1R)-3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-(4-
chlorophenyl)-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-
10 cyclohexanecarboxylic amide



Scheme 17



Step A: Synthesis of intermediate 2 (Scheme 17).

The starting material **1** (prepared as the procedure described in *J. Chem. Res.* 1981, 1772-1783) (10 mmol, 2.00 g) was taken up in 50 mL of anhydrous acetonitrile and cooled to 0° C in an ice bath. To this was added DBU (10 mmol, 1.49 mL) and the reaction was stirred for 10 minutes before adding benzyl bromide (10 mmol, 1.19 mL). The reaction was allowed to slowly warm to room temperature and stir 16 hrs. The reaction was then evaporated to an oil, taken up in methylene chloride and washed with 5% NaHSO₄, saturated NaHCO₃ and brine. The organic layer was dried over Na₂SO₄, filtered and evaporated to give the desired product (93%, 2.70 g). ¹H-NMR (300 MHz, CDCl₃) δ: 1.40 - 1.60 (m, 10H), 2.00 - 2.10 (m, 2H), 2.67 (s, 2H), 3.60 (s, 3H), 5.10 (s, 2H), 7.40 (s, 5H); (M + H⁺) 351.3.

Step B: Synthesis of intermediate 4 (Scheme 17).

The starting material 2 (11 mmol, 3.10 g) was taken up in 50 mL anhydrous THF, cooled to -78°C and treated with lithium bis(trimethylsilyl) amide (22 mmol, 21.2 mL of 1.0 M solution in THF) under an inert atmosphere for 1.0 hr.

5 Bromomethylpropene (33 mmol) was added slowly at -78°C and the reaction was allowed to gradually warm to room temperature and stir overnight. The reaction was then diluted with ethyl acetate and washed with water followed by saturated NaHCO_3 . The organic layer was dried over
10 Na_2SO_4 , filtered and evaporated to give the desired product (96%, 3.51 g). ^1H -NMR (300 MHz, CDCl_3) δ : 1.00 - 2.00 (m, 10H), 3.40 - 3.70 (m, 6H), 4.50 - 4.70 (m, 2H), 4.80 - 5.20 (m, 2H), 7.19 - 7.35 (m, 5H).

15 Step C: Synthesis of intermediate 5 (Scheme 17).

The product obtained from the previous reaction 4 (10 mmol, 3.51 g) was taken up in 50 mL ethyl acetate and treated with 30% by weight (.525 g) of 10% palladium on activated carbon and in a Parr shaker overnight under 50
20 psi H_2 . The reaction was then purged with nitrogen, filtered through celite and evaporated in vacuo. The resulting residue was taken up in ethyl acetate and washed with saturated NaHCO_3 . The aqueous layer was separated, acidified by the dropwise addition of concentrated HCl and
25 extracted with ethyl acetate. The ethyl acetate extract was evaporated in vacuo to give the desired product 5 (76%, 2.01 g). ^1H -NMR (300 MHz, CDCl_3) δ : 0.80 - 0.99 (m, 6H), 1.10 - 1.80 (m, 10H), 1.80 - 2.20 (m, 2H), 3.77 (s, 3H); (M + H^+) 257.3.

30

Step D: Synthesis of intermediate 7 (Scheme 17).

Carboxylic acid 5 (0.27 mmol, 70 mg) was taken up in 5.0 mL of anhydrous DMF and treated with HATU (0.27 mmol, 104 mg). After 30 minutes, Hunig's base (0.54 mmol, 0.1
35 mL) and the benzodiazepine amine 6 (0.27 mmol, 61 mg) were added. The reaction was stirred at room temperature under a nitrogen atmosphere overnight, then diluted with ethyl acetate and washed with 5% NaHSO_4 followed by saturated

NaHCO₃ and brine. The organic layer was evaporated in vacuo to give a crude yellow oil which was purified by flash chromatography using 75% hexane and 25% ethyl acetate to give the desired product (100%, 160 mg). H¹-NMR (300 MHz, CDCl₃) δ: 0.8 - 0.9 (m, 6H), 1.10 - 1.95 (m, 10H), 2.50 - 2.60 (m, 1H), 3.49 (s, 3H), 3.76 (d, 3H), 5.52 - 5.55 (m, 1H), 7.10 - 7.61 (m, 8H); (M + H⁺) 538.3.

Step E: Synthesis of intermediate 8 (Scheme 17).

The ester 7 (0.28 mmol, 150 mg) was taken up in 7.0 mL of THF and was treated with LiOH-H₂O (1.4 mmol, 65 mg) in 1.0 ml H₂O. Methanol was added to ensure homogeneity and the reaction was stirred overnight at room temperature. The reaction was then diluted with ethyl acetate and extracted with H₂O. The aqueous phase was separated and acidified by the dropwise addition of concentrated HCl. This was then extracted with ethyl acetate to give the desired product (46%, 67 mg). H¹-NMR (300 MHz, CDCl₃) δ: 0.8 - 1.00 (m, 6H), 1.10 - 2.00 (m, 10H), 3.50 (s, 1H), 5.50 - 5.60 (m, 1H), 7.30 - 7.65 (m, 8H); (M + H⁺) 524.2.

Step F: Synthesis of intermediate 9 (Scheme 17).

The acid 8 obtained from the above reaction (0.13 mmol, 67 mg) was taken up in 7.0 mL of anhydrous DMF, was added HATU (0.13 mmol, 49 mg) and stirred for 30 min. before bubbling in NH₃ (g) for 30 min. The reaction was then capped and allowed to stand at room temperature overnight. Afterwards, the reaction was diluted with ethyl acetate and washed with 5% NaHSO₄, saturated NaHCO₃ and brine. The organic layer was dried over Na₂SO₄, filtered and evaporated to give crude yellow oil which was purified by rotary TLC using 75% ethyl acetate and 25% hexane to give the desired product, 9 as a mixture of diastereomers (14%, 10 mg). H¹-NMR (300 MHz, CDCl₃) δ: 0.8 - 1.00 (m, 6H), 1.20 - 1.85 (m, 10H), 2.30 - 2.35 (d, 1H), 2.40 - 2.55 (m, 1H), 3.50 (s, 3H), 5.52 (d, 2H), 7.10 - 7.90 (m, 8H); (M + H⁺) 523.3.


Additional examples of the present invention are illustrated and prepared according to the procedures described herein using starting materials appropriate for the desired products.

5

Example 4

1-[(1R)-3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclohexanecarboxylic amide, 12

10




The title compound 12 was prepared using methods similar to those employed in Example 3. The product was obtained as an oil. ¹H-NMR (300 MHz, CDCl₃) δ: 0.85 - 1.05 (m, 6H), 1.15 - 1.90 (m, 10H), 3.41 - 3.55 (m, 3H), 5.30 - 5.60 (m, 3H), 7.38 - 7.90 (m, 9H); (M - H⁺) 487.2.

15

Example 5

1-[(1R)-3-Methyl-1-(5-methyl-6-oxo-6,7-dihydro-5H-dibenzo[b,d]-azepin-7-ylcarbamoyl)-butyl]-cyclohexanecarboxylic acid amide, 15


20



The title compound **15** was prepared using methods similar to those employed in Example 3. The product was obtained as an oil. ¹H-NMR (300 MHz, CDCl₃) δ: 0.88 (d, 3H), 0.97 (d, 3H), 1.20 - 1.90 (m, 10H), 2.00 (d, 1H), 2.35 (d, 1H), 2.55 (d, 1H), 3.39 (s, 3H), 5.25 - 5.40 (m, 2H), 7.30 - 7.70 (m, 8H); (M + H⁺) 462.3.

Examples 6 and 6a

- 10 1-[(1R)-3-Methyl-1-(2-oxo-2,3,4,5-tetrahydro-1H-benzo[b]azepin-3-ylcarbamoyl)-butyl]-cyclopent-3-enecarboxylic acid amide, **20**




The reaction procedures used to make compound 19 are shown in Scheme 4a and in the Equation above. Compound 16 is obtained from literature procedures (J. Med. Chem. 1999, **42**, p 2621.) Compound 19, a mixture of diastereomers, was separated by flash chromatography using 100% EtOAc to give the final products Example 6 (20) and diastereomer Example 6a, respectively. Compound 20 was the first eluting isomer: ¹H NMR (300MHz, CDCl₃) 7.27-7.02 (m, 4H), 5.53 (m, 2H), 4.41-4.34 (q, 1H), 2.94-2.34 (m, 8H), 2.18-2.02 (m, 1H), 1.80-1.66 (m, 1H), 1.60-1.42 (m, 1H), 1.00 (m, 1H), 0.91-0.81 (q, 6H); MS [M + H]⁺ 384.

Example 6a was the second eluting isomer: ^1H NMR (300MHz, CDCl_3) 7.26-7.02 (m, 4H), 5.54 (m, 2H), 4.43-4.36 (q, 1H), 2.96-2.36 (m, 8H), 2.20-2.04 (m, 1H), 1.80-1.68 (m, 1H), 1.44-1.30 (m, 1H), 1.06 (m, 1H), 0.90-0.78 (q, 6H); MS $[\text{M} + \text{H}]^+$ 384.

Example 8

1-(1-{1-[3-(2-Fluoro-phenoxy)-benzyl]-2-oxo-2,3,4,5-tetrahydro-1H-benzo[b]azepin-3-ylcarbamoyl}-3-methyl-butyl)-cyclopent-3-enecarboxylic acid amide, 23




5

To a solution of 20 (11 mg) in CH_3CN (2 mL) was added K_2CO_3 (8 mg) and 3-(2-fluorophenoxy)benzyl bromide 22 (12 mg). The reaction mixture was stirred at rt for 20 hrs before the solvent was removed under pressure. The crude product was purified by flash chromatography to give compound 23 (77%, 13 mg). ^1H NMR (300MHz, CDCl_3) 7.32-6.80 (m, 12H), 5.72 (m, 1H), 5.58 (m, 1H), 5.34 (d, 1H), 4.72 (d, 1H), 4.48 (m, 1H), 3.18 (m, 1H), 2.70-2.20 (m, 8H), 1.95-1.78 (m, 2H), 1.42 (m, 1H), 1.12 (m, 1H), 0.92-0.80 (q, 6H); MS $[\text{M} + \text{H}]^+$ 584.

15

Example 9




20

The compound of 24 was made in a similar manner as described above for Example 8. ^1H NMR (300MHz, CDCl_3) 7.30-6.78 (m, 12H), 5.68 (m, 1H), 5.60 (m, 1H), 5.3 (d, 1H), 4.76 (d, 1H), 4.50 (m, 1H), 3.15 (m, 1H), 2.66-2.20

(m, 8H), 1.96-1.80 (m, 2H), 1.44 (m, 1H), 1.15 (m, 1H), 0.94-0.84 (q, 6H); MS $[M + H]^+$ 584.

Example 10

- 5 1-(3-Methyl-1-[2-oxo-1-(3-phenylamino-benzyl)-2,3,4,5-tetrahydro-1H-benzo[b]azepin-3-ylcarbamoyl]-butyl)-cyclopent-3-enecarboxylic acid amide, 27




27

- 10 (3-Phenylamino-phenyl)-methanol (Eur. Pat. Appl. 1994, 55) was converted to the (3-Phenylamino)benzyl-bromide intermediate by mixing with PPh_3 and CBr_4 in CH_2Cl_2 at rt for 3 hours. The crude product was used for the synthesis of 27. The title compound 27 was prepared using methods similar to those described in Example 8. The product was obtained as an oil. 1H NMR (300MHz, $CDCl_3$) 7.38-6.80 (m, 13H), 5.68 (m, 1H), 5.58 (m, 1H), 5.24 (d, 1H), 4.72 (d, 1H), 4.48 (m, 1H), 3.15 (m, 1H), 2.70-2.20 (m, 8H), 1.98-1.86 (m, 2H), 1.42 (m, 1H), 1.10 (m, 1H), 0.90-0.80 (q, 6H);
- 15 MS $[M + H]^+$ 565.
- 20

Example 11

- 1-(3-Methyl-1-[2-oxo-1-(3-phenylamino-benzyl)-2,3,4,5-tetrahydro-1H-benzo[b]azepin-3-ylcarbamoyl]-butyl)-cyclopentanecarboxylic acid amide, 28
- 25




28

The title compound **28** was prepared by reduction of the Example 10 using 10% palladium on activated carbon in a Parr shaker overnight under 50 psi H₂. The product was obtained as an oil. ¹H NMR (300MHz, CDCl₃) 7.30-6.80 (m, 13H), 5.20 (d, 1H), 4.64 (d, 1H), 4.42 (m, 1H), 2.65-2.22 (m, 6H), 1.92-1.05 (m, 10H), 1.42 (m, 1H), 1.10 (m, 1H), 0.86-0.72 (q, 6H); MS [M + H]⁺ 567.

Using the methods described herein, the following examples of a Compound of Formula (I) were prepared.

Example 12

1-[2-Cyclopropyl-1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-ylcarbamoyl]-ethyl]-cyclopent-3-enecarboxylic amide



(Diastereomer A)


Optical Rotation $[\alpha]_D^{25} +45.3^\circ$ (c 0.10, Methanol)
Melting Point 238-240°C
Mass Ion 471 (M+H)

(Diastereomer B)

Optical Rotation $[\alpha]_D^{25} -18.2^\circ$ (c 0.10, Methanol)
Melting Point 382-383 °C
Mass Ion 471 M+H)

Example 14

1-[2-Cyclopropyl-1-[1,3-dihydro-1-methyl-2-oxo-5-(2-trifluoromethylphenyl)-2H-1,4-benzodiazepin-3-ylcarbamoyl]-ethyl]-cyclopent-3-enecarboxylic amide



(Diastereomer B)

Optical Rotation $[\alpha]_D^{25}$ -47.3° (c 0.20, Methanol)

Melting Point $235-236^\circ\text{C}$

5 Mass Ion 539 (M+H)

(Diastereomer A)

Optical Rotation $[\alpha]_D^{25}$ $+93.7^\circ$ (c 0.05, Methanol)


Melting Point $344-345^\circ\text{C}$

10 Mass Ion 539 (M+H)

Following the procedures outlined in Scheme 19 and Scheme 19a, the title compound Example 14 was prepared.

15

Scheme 19



Data for compound 3; Scheme 19.

- 5 Collected 3 as a white solid: ¹H NMR (CDCl₃) δ 7.42–7.13 (m, 5 H), 4.68 (m, 1 H), 4.18 (m, 2 H), 3.31 (dd, *J* = 13.3, 3.3 Hz, 1 H), 4.59 (m, 2 H), 2.79 (dd, *J* = 15.0, 3.7 Hz, 1 H), 1.60 (m, 2 H), 0.79 (m, 1 H), 0.47 (m, 2 H), 0.18–0.00 (m, 2 H).

10

Data for compound 5; Scheme 19.

- 15 Collected 5 (45 g, 55%) as a white solid: ¹H NMR (CDCl₃) δ 7.40–7.10 (m, 5 H), 4.65 (m, 1 H), 4.32 (m, 1 H), 4.16 (m, 2 H), 3.34 (dd, *J* = 13.3, 3.2 Hz, 1 H), 2.96–2.53 (m, 3 H), 1.65–1.30 (m, 11 H), 0.75 (m, 1 H), 0.43 (m, 2 H), 0.02–0.00 (m, 2 H).

Data for compound 6; Scheme 19.

Collected **6** (22.5 g, 77%) as a clear viscous oil: ^1H NMR (CDCl_3) δ 2.92 (m, 1 H), 2.68 (dd, $J = 16.6, 9.3$ Hz, 1 H), 2.50 (dd, $J = 16.5, 5.2$ Hz, 1 H), 1.67-1.38 (m, 2 H), 1.44 (s, 9 H), 0.72 (m, 1 H), 0.48 (m, 2 H), 0.08 (m, 2 H).

5

Data for compound 7; Scheme 19.

Collected **7** (5.1 g, 87%) as a clear, viscous oil: ^1H NMR (CDCl_3) δ 5.74 (m, 1 H), 5.07 (m, 2 H), 2.76 (m, 2 H), 2.39 (m, 2 H), 1.64 (m, 1 H), 1.47 (m, 1 H), 1.43 (s, 9 H), 0.74 (m, 1 H), 0.48 (m, 2 H), 0.08 (m, 2 H).

10

Data for compound 8; Scheme 19.

Collected **8** (9.2 g, 93%) as a clear, viscous oil: ^1H NMR (CDCl_3) δ 5.77 (m, 2 H), 5.10 (m, 4 H), 2.79 (dd, $J = 11.8, 2.9$ Hz, 1 H), 2.55 (dd, $J = 14.5, 6.4$ Hz, 1 H), 2.44 (m, 2 H), 2.28 (dd, $J = 14.4, 8.2$ Hz, 1 H), 1.76 (ddd, $J = 11.9, 11.8, 6.3$ Hz, 1 H), 1.40 (s, 9 H), 1.29 (m, 1 H), 0.69 (m, 1 H), 0.45 (m, 2 H), 0.02-0.00 (m, 2 H).

15

20 *Data for compound 9; Scheme 19.*

Collected **9** (4.8 g, 86%) as a viscous oil: ^1H NMR (CDCl_3) δ 7.43-7.29 (m, 5 H), 5.77 (m, 2 H), 5.20-4.95 (m, 6 H), 2.83 (dd, $J = 11.8, 2.6$ Hz, 1 H), 2.49 (ddd, $J = 14.2, 6.3$ Hz, 1 H), 2.41 (m, 2 H), 2.25 (dd, $J = 14.3, 7.9$ Hz, 1 H), 1.81 (m, 1 H), 1.43 (s, 9 H), 1.35 (m, 1 H), 0.55 (m, 1 H), 0.35 (m, 2 H), 0.01 (m, 2 H).

25

Data for compound 11; Scheme 19.

Collected **11** (4.6 g, >95%) as a viscous oil: ^1H NMR (CDCl_3) δ 7.36 (m, 5 H), 5.51 (m, 2 H), 5.10 (m, 2 H), 3.01 (m, 1 H), 2.91-2.52 (m, 4 H), 1.78 (m, 1 H), 1.41 (s, 9 H), 1.07 (m, 1 H), 0.64 (m, 1 H), 0.38 (m, 2 H), 0.01 (m, 2 H).


30

Data for compound 12; Scheme 19.

Collected **12** (1.9 g, >98%) as a clear viscous oil: ^1H NMR (CDCl_3) δ 5.60 (s, 2 H), 2.89 (m, 3 H), 2.58 (m, 2 H), 1.78 (m, 1 H), 1.44 (s, 9 H), 1.12 (m, 1 H), 0.78 (m, 1 H), 0.48 (m, 2 H), 0.20-0.00 (m, 2 H).

35

Scheme 19a



5

Preparation of compound 14; Scheme 19a.

To a solution of **12** (70 mg, 0.25 mmol) and ClCH₂CH₂Cl (1.25 mL) at 0 °C was added HATU (109 mg, 0.29 mmol), DIPEA (174 μL, 1.0 mmol) and **13** (84 mg, 0.25 mmol). The reaction was warmed to rt and stirred for 18 h. The reaction was diluted with sat. NH₄Cl (70 mL). The aqueous layer was extracted with EtOAc (2 x 70 mL). The organic layers were combined and washed with NaHCO₃ (70 mL) and brine (70 mL). The organic layer was dried over MgSO₄ and filtered. The crude material was purified by SiO₂ chromatography (5% ethyl acetate in hexane) to afford **14** (110 mg, 74%) as a white solid: ¹H NMR (CDCl₃) δ 7.70-7.46 (m, 5 H), 7.36 (dd, *J* = 8.3, 4.0 Hz, 1 H), 7.12 (t, *J* = 7.7 Hz, 1 H), 7.00 (m, 1 H), 5.66-5.49 (m, 3 H), 3.49 (m, 3 H), 2.97-2.62 (m, 6 H), 1.95 (m, 2 H), 1.45 (s, 9 H), 1.02-0.70 (m, 1 H), 0.65-0.39 (m, 2 H), 0.20-0.00 (m, 2 H).

20

Preparation of compounds 15a and 15b; Scheme 19a.

To a round bottom flask containing 14 (110 mg, 0.19 mmol) was added CH_2Cl_2 (1 mL) and TFA (1 mL). The solution was stirred at rt for 2 h. The solvent was removed under reduced pressure and the intermediate was dissolved in CH_2Cl_2 (5 mL). The solution was treated with HATU (76 mg, 0.2 mmol) and DIPEA (131 μM , 0.8 mmol). Ammonia gas was bubbled through the solution for 5 min. and the reaction was stirred for 18 h.

The reaction was diluted with CH_2Cl_2 (50 mL) and washed with 10% citric acid (30 mL), NaHCO_3 (30 mL) and brine (30 mL). The organic layer was dried over MgSO_4 and filtered. The solvent was removed under reduced pressure. The crude product was purified by SiO_2 chromatography (5% methanol in CH_2Cl_2) to give 15a,b (66 mg, 68%). The preparatory HPLC separation was conducted with a Waters Symmetry C18 column (5 cm x 25 cm, 118 mL/min, 240 nm, 50:50 $\text{H}_2\text{O}/\text{CH}_3\text{CN}$; 2% TFA).

Data for compound 15a; Scheme 19a.

Collected **15a** (22 mg, 22%) as a white powder: mp 344-345 $^\circ\text{C}$; ^1H NMR (CDCl_3) δ 7.98 (m, 1 H), 7.74-7.51 (m, 6 H), 7.39 (d, J = 7.6 Hz, 1 H), 7.16 (t, J = 7.2 Hz, 1 H), 7.05 (dd, J = 7.9, 1.5 Hz, 1 H), 5.71 (m, 1 H), 5.63 (m, 1 H), 5.58 (d, J = 7.8 Hz, 1 H), 5.35 (m, 1 H), 3.50 (s, 3 H), 3.24 (dt, J = 16.4, 1.4 Hz, 1 H), 2.74 (m, 2 H), 2.28 (m, 1 H), 1.98 (ddd, J = 13.9, 11.3, 5.9 Hz, 1 H), 1.67 (m, 1 H), 1.23 (m, 1 H), 0.85 (m, 1 H), 0.48 (m, 2 H), 0.15-0.00 (m, 2 H); IR (ATR) 3339, 2923, 1657, 1600, 1490, 1377, 1311, 1110, 1035, 912, 764, 731, 685 cm^{-1} ; ESI MS m/z = 539 [$\text{C}_{29}\text{H}_{29}\text{F}_3\text{N}_4\text{O}_3 + \text{H}$] $^+$; $[\alpha]^{25}_{\text{D}}$ +93.7 (c 0.05, Methanol); HPLC >95% t_r = 16.20 min.


Data for compound 15b; Scheme 19a.

Collected **15b** (24 mg, 25%) as a white powder: mp 235-236 $^\circ\text{C}$; ^1H NMR (CDCl_3) δ 8.16 (m, 1 H), 7.75-7.53 (m, 7 H), 7.40 (d, J = 8.2 Hz, 1 H), 7.16 (t, J = 7.5 Hz, 1 H), 7.04 (d, J = 6.7 Hz, 1 H), 5.70 (m, 1 H), 5.64 (m, 1 H), 5.57 (d, J = 8.5 Hz, 1 H), 3.51 (s, 3 H), 3.18 (d, J = 16.7 Hz, 1 H),


2.78 (d, $J = 9.6$ Hz, 1 H), 2.70 (m, 2 H), 2.35 (d, $J = 16.0$ Hz, 1 H), 1.86 (m, 1 H), 1.34 (m, 1 H), 0.80 (m, 1 H), 0.48 (m, 2 H), 0.16 (m, 1 H), 0.03 (m, 1 H); IR (ATR) 3331, 3068, 2924, 1652, 1491, 1448, 1313, 1159, 1111, 763 cm^{-1} ;
 5 ESI MS $m/z = 539$ [$\text{C}_{29}\text{H}_{29}\text{F}_3\text{N}_4\text{O}_3 + \text{H}$] $^+$; $[\alpha]^{25}_{\text{D}} -47.3$ (c 0.2, Methanol); HPLC >95% $t_r = 17.04$ min.

Example 16

10 1-[2-Cyclopropyl-1-(5-methyl-6-oxo-6,7-dihydro-5H-dibenzo[b,d]azepin-7-ylcarbamoyl)-ethyl]-cyclopent-3-enecarboxylic acid amide



Scheme 20




20 Compound 17 was prepared by the same method as compound 14 in Example 14. Collected 17 (87 mg, 73%) as a white solid: ^1H NMR (CDCl_3) δ 7.63-7.30 (m, 8 H), 5.72-5.46 (m, 3 H), 5.36 (d, $J = 6.6$ Hz, 1 H), 3.37 (s, 3 H), 3.05-2.57 (m, 5 H), 2.10-1.90 (m, 1 H), 1.46 (s, 9 H), 0.97 (m, 1 H), 0.78 (m, 1 H), 0.54 (m, 2 H), 0.12 (m, 2 H).

Compound 18 was prepared by the same method as compound 15a and 15b in Example 14. Collected 18 (32 mg, 33%) as a white powder: mp 217-222 °C; ^1H NMR (CDCl_3) δ 7.12-7.34 (m, 8 H), 5.72 (m, 1 H), 5.65 (m, 1 H), 5.38 (d, $J = 6.6$ Hz, 1 H), 5.27 (m, 1 H), 3.37 (s, 3 H), 3.18 (m, 1 H), 2.93 (dd, $J = 11.0, 2.9$ Hz, 1 H), 2.60 (m, 2 H), 2.35 (m, 1 H), 2.00 (ddd, $J = 14.0, 11.1, 6.3$ Hz, 1 H), 1.27 (ddd, $J = 14.1, 8.0, 3.1$ Hz, 1 H), 0.85 (m, 1 H), 0.57 (m, 2 H), 0.15 (m, 2 H); IR 3346, 3229, 1652, 1600, 1498, 1382, 1301, 917 cm^{-1} ; ESI MS $m/z = 444$ [$\text{C}_{27}\text{H}_{29}\text{N}_3\text{O}_3 + \text{H}$] $^+$; $[\alpha]^{25}_{\text{D}} +95.2$ (c 0.07, Methanol); HPLC >95% tr = 16.19 min.
(Diastereomer A)

15


Example 17

1-{3-Methyl-1-[2-oxo-1-(3-o-tolylamino-benzyl)-azepan-3-ylcarbamoyl]-butyl}-cyclopent-3-enecarboxylic acid amide



20

Scheme 21



Preparation of Compound 4, Scheme 21. A solution of BINAP (0.22 g, 0.35 mmol) in toluene (12.5 mL) was degassed with Ar and refluxed at 80 °C for 1 min. After cooling the solution to room temperature, Pd(OAc)₂ (0.05 g, 0.23 mmol) was added and the mixture was stirred for 10 min. Next, 2 (1.03 g, 2.32 mmol) in toluene (3 mL) was added followed by *p*-toluidine (1.24 mL, 11.6 mmol) and NaOt-Bu (0.27 g, 2.8 mmol). The reaction was refluxed under Ar at 100 °C overnight. The reaction was diluted with H₂O (20 mL) and extracted with EtOAc (2 x 40 mL). The organic extracts were dried over MgSO₄, filtered through Celite, and concentrated under reduced pressure. The crude material was subjected to flash chromatography on silica (hexanes/EtOAc) to provide 4 (0.45 g, 46%) as a pink solid: ¹H NMR (500 MHz, CDCl₃) δ 7.25-6.71 (m, 8 H), 6.03 (m, 1 H), 5.38 (s, 1 H), 4.77 (d, *J* = 14.68 Hz, 1 H), 4.41 (m, 1 H), 4.33 (d, *J* = 14.67 Hz, 1 H), 3.42 (m, 1 H), 3.23 (dd, *J* = 14.97, 4.49 Hz, 1 H), 2.24 (s, 3 H), 2.08-1.21 (m, 6 H), 1.45 (m, 9 H); APCI MS *m/z* = 424 [C₂₅H₃₃N₃O₃]⁺.

Preparation of Compound 5, Scheme 21. To saturated HCl/EtOH (2.6 mL) at 0 °C was added 4 (0.44 g, 1.0 mmol). The mixture was stirred overnight at room temperature and the solvent was removed under reduced pressure giving 5 (1.7 g, 94%) as an off-white solid: ¹H NMR (500 MHz, CD₃OD) δ 7.32-6.83 (m, 8 H), 4.67 (q, *J* = 28.2, 14.61 Hz, 2 H), 4.40 (d, *J* = 10.98 Hz, 1 H), 3.72-3.47 (m, 2 H), 2.33 (s, 3 H), 2.13-1.31 (m, 6 H); APCI MS *m/z* = 324 [C₂₀H₂₆N₃O₃Cl+H]⁺.

Preparation of Compound 7, Scheme 21. To a solution of 5 (0.2, 0.56 mmol) in CH₂Cl₂ (2.0 mL) cooled to 0 °C was added HATU (0.25 g, 0.65 mmol) and a solution of succinate 6 (0.16 g, 0.57 mmol) in CH₂Cl₂ (0.7 mL). DIPEA (0.39 mL, 2.3 mmol) was added, and the reaction was warmed to rt and stirred overnight. The reaction was concentrated, added to H₂O (10 mL), and extracted with EtOAc (2 x 20 mL). The

organic extract was washed successively with H₂O (3 x 6 mL), brine (8 mL), dried over MgSO₄, and filtered. Solvent removal under reduced pressure afforded the crude product which was subjected to flash chromatography on silica

5 (Hexanes/EtOAc) to provide 7 (0.21 g, 64%) as an off-white solid: ¹H NMR (500 MHz, CDCl₃) δ 7.27-6.74 (m, 9 H), 5.59-5.42 (m, 3 H), 4.68-4.49 (m, 3 H), 3.48-3.22 (m, 2 H), 2.86-1.00 (m, 14 H), 2.24 (s, 3 H), 1.47 (s, 9 H), 0.88 (m, 6 H); APCI MS m/z = 588 [C₃₆H₄₉N₃O₄+H]⁺.

10

Preparation of Compound 8, Scheme 21. A solution of 7 (0.20 g, 0.34 mmol) in 1:1 TFA/CH₂Cl₂ (12 mL) was stirred at rt overnight. The solvent was removed under reduced pressure at 60 °C to afford 8 (0.19 g, >95%) as a brown semisolid: ¹H NMR (500 MHz, CD₃OD) δ 7.22-6.61 (m, 8 H), 5.67-5.49 (m, 2 H), 4.62 (d, J = 11.3 Hz, 1 H), 4.56-4.39 (m, 2 H), 3.47-1.02 (m, 16 H), 2.15 (s, 3 H), 0.90-0.77 (m, 6 H). APCI MS m/z = 532 [C₃₂H₄₁N₃O₄+H]⁺.

20

Preparation of Compound 1, Scheme 21. To a solution of DIPEA (0.75 mL, 4.3 mmol) in DMF (6.9 mL) was added 8 (0.19 g, 0.34 mmol) and HATU (0.22 g, 0.58 mmol). Ammonia gas was bubbled in the solution for 10 min, and the reaction was stirred overnight. The reaction was concentrated under reduced pressure and partitioned between EtOAc (10 mL) and H₂O (10 mL). The aqueous portion was extracted with EtOAc (2 x 10 mL). The combined organic extracts were washed successively with H₂O (3 x 4 mL) and brine (4 mL), dried over MgSO₄, and filtered. Solvent removal under reduced pressure afforded the crude product which was subjected to flash chromatography on silica (Hexanes/EtOAc) then (EtOAc/MeOH) to provide 1 as an off-white solid (0.086 g, 48%): mp 90-91 °C; ¹H NMR (500 MHz, CD₃OD) δ 7.20-6.68 (m, 8 H), 5.59 (s, 2 H), 4.65 (d, J = 10.45 Hz, 1 H), 4.59 (d, J = 14.68 Hz, 1 H), 4.46 (d, J = 14.65 Hz, 1 H), 3.54 (m, 1 H), 3.30 (m, 1 H), 2.84 (d, J = 11.8 Hz, 1 H), 2.74 (s, 2 H), 2.61 (d, J = 17.2 Hz, 1 H), 2.21 (s, 3 H), 1.91-1.03 (m, 10 H), 0.90 (dd, J = 13.65,


35

6.51 Hz, 6 H); ESI MS m/z = 531 $[C_{32}H_{42}N_4O_3+H]^+$; IR (KBr) 3320 (br.), 1630 cm^{-1} ; HPLC >95%, t_r = 21.31 min. Optical Rotation $[\alpha]_D^{25}$ -3.3 (c 0.10, Methanol)

- 5 Using the methods described herein, the following examples of a Compound of Formula (I) were prepared.

Example 18


- 10 1-[3-Methyl-1-(5-methyl-6-oxo-6,7-dihydro-5H-dibenzo[b,d]azepin-7-ylcarbamoyl)-butyl]-cyclopent-3-enecarboxylic acid amide.



- 15 1H NMR (300 MHz, $CDCl_3$) 0.8-1.0 (dd, 6H), 1.0-1.2 (m, 1H), 1.4-1.6 (m, 1H), 1.7-1.9 (m, 1H), 2.3-2.8 (m, 4H), 3.0-3.2 (m, 1H), 5.2 (s, 1H), 5.3 (s, 1H), 5.6-5.8 (m, 2H), 7.2-7.6 (m, 8H). MS: 446.4 (M+H).


20 Example 19

- 1-[3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-(4-chlorophenyl)-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclopent-3-enecarboxylic acid amide.



25 Synthesis of Formula I compound 17 (Scheme 15)

Scheme 15



- 5 Step 1: Preparation of diallylsuccinate HOBT ester 14.
 To a solution of 2 (30.5g, 98.4mmol) in 500ml methylene
 chloride was added HOBT (27g, 200mmol), EDC (25g, 130mmol)
 and triethylamine (21ml, 150mmol). The solution was stirred
 at RT for three hours. The solvents were removed under
 reduced pressure, providing an oil which was taken up in
 EtOAc and water. The organic layer was washed with water
 and brine, dried over sodium sulfate, then concentrated to
 give the crude product as an oil. Purification by column
 chromatography on silica gel with EtOAc:hexane (5:95)
 provided 14 as a colorless oil (32.2g, 75.4mmol, 77%). ¹HNMR
 (300 MHz, CDCl₃) 0.9-1.1 (m, 6H), 1.2-1.4 (m, 1H), 1.4 (s,
 9H), 1.8-2.0 (m, 2H), 2.3-2.8 (m, 4H), 3.2 (m, 1H), 5.0-5.2
 (m, 4H), 5.6-5.9 (m, 2H), 7.3-8.1 (m, 4H).

Step 2: Preparation of benzodiazepinesuccinamide 16


Following the procedures described for the synthesis of 11, diallylsuccinamide intermediate 16 was prepared from 14 in reasonable yield. 14 (2.5g, 5.9mmol), p-chloro-BZD (2.34g, 6.2mmol) and DIPEA (1ml, 5.8mmol) in 30ml DMF was heated to 40-50 degree overnight.

Step 3: Preparation of Formula I compound 17

A solution of 16 (700mg, 1.3mmol) in 100ml toluene and methylene chloride (1:1) and tricyclohexylphosphine[1,3-bis(2,4,6-trimethylphenyl)-4,5-dihydroimidazol-2-ylidene]benzylidene]ruthenium (IV) dichloride (100mg, 0.1mmol) was heated at 60°C for two hours. The solvents were evaporated to give a dark oil which was purified by flash chromatography using EtOAc:hexane (1:1) to give 17 as a solid (390mg, 1077mmol, 59%). ¹HNMR (300 MHz, CDCl₃) 0.8-1.0 (dd, 6H), 1.2-1.3 (m, 1H), 1.5-1.7 (m, 1H), 1.8-2.0 (m, 1H), 2.3-3.0 (m, 4H), 3.2 (m, 1H), 3.5 (s, 3H), 5.3 (s, 1H), 5.5 (d, 1H), 5.6-5.8 (m, 2H), 7.2-7.7 (m, 8H), 7.8 (s, 1H). MS: 507.4 (M+H), 529.3 (M+Na).

Example 20

1-[3-Methyl-1-(5-methyl-6-oxo-6,7-dihydro-5H-dibenzo[b,d]azepin-7-ylcarbonyl)-butyl]-cyclopentanecarboxylic acid amide.




¹HNMR (300 MHz, CDCl₃) 0.8-1.0 (dd, 6H), 1.0-2.0 (m, 10H), 2.4-2.7 (m, 2H), 3.3 (s, 3H), 5.3 (d, 1H), 5.6 (s, 1H), 7.2-7.8 (m, 8H), 8.1 (s, 1H). MS: 448.4 (M+H), 470.4 (M+Na).

Example 21

1-[3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-(4-chlorophenyl)-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclopentanecarboxylic amide.


5



A solution of Example 19 (300mg, 0.59mmol) in 50ml ethanol with 300mg of chlorotris-(triphenylphosphine) rhodium(I), was shaken under H₂ (~50psi) overnight. The solvents were removed under reduced pressure, and the resulting residue was purified by chromatography on silica gel in 5% methanol/CH₂Cl₂. The desired product (18) was isolated as a solid (247mg, 0.49mmol). ¹HNMR (300 MHz, CDCl₃) 0.8-1.0 (dd, 6H), 1.2-2.0 (m, 10H), 2.4-2.6 (m, 2H), 3.5 (s, 3H), 5.3 (s, 1H), 5.5 (d, 1H), 7.2-7.7 (m, 8H), 8.1 (s, 1H). MS: 509.4 (M+H).

Example 22

1-[3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-(4-trifluoromethylphenyl)-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclopent-3-enecarboxylic amide.




¹H NMR (300 MHz, CDCl₃) 0.8-1.0 (dd, 6H), 1.1-1.3 (m, 1H), 1.5-1.7 (m, 1H), 1.8-2.0 (m, 1H), 2.2-3.0 (m, 4H), 3.0-3.3 (m, 1H), 3.5 (s, 3H), 5.3 (s, 1H), 5.5 (d, 1H), 5.6-5.8 (m, 2H), 7.2-7.8 (m, 8H). MS: 541.5 (M+H), 563.5 (M+Na).

5


Example 23**Example 23a**

1-[3-methyl-1-[1,3-dihydro-1-(i-propyl)-2-oxo-5-(2-fluorophenyl)-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclopent-3-enecarboxylic amide.

10



Example 23




Example 23a

15 Synthesis of Formula I compounds 21 and 22.

Diastereomers 21 and 22 were prepared from diallylsuccinamide intermediate 19 according to the methods outlined in Scheme 16.

20

Scheme 16



Synthesis of diallylsuccinamide intermediate 19

Intermediate 19 was made according to the procedures

- 5 outlined in Scheme 4, using the appropriate
aminobenzodiazepine. 1H NMR (300 MHz, $CDCl_3$) 0.8-1.0 (m, 6H), 1.0-2.0 (m, 3H), 2.2-2.8 (m, 5H), 5.0-5.3 (m, 4H), 5.4-5.6 (d, 1H), 5.6-6.0 (m, 2H), 7.0-7.6 (m, 8H). MS: 505.3 (M+H), 527.3 (M+Na).

10

Synthesis of intermediate 20 by alkylation of 19.

To a solution of 19 (150mg, 0.3mmol) in 20ml DMF was added potassium carbonate (90mg, 0.65mmol) and 2-iodopropane (120mg, 0.69mmol). The reaction mixture was stirred at RT
15 overnight. The solvents were removed under reduced pressure and the residue was taken up in EtOAc and water. The organic layer was washed with brine, dried over sodium sulfate and concentrated, and the resulting solid was purified by column chromatography on silica gel in EtOAc:hexane (70:30)
20 to provide 20 as a white solid (105mg, 0.19mmol). 1H NMR (300 MHz, $CDCl_3$) 0.8-1.0 (m, 6H), 1.0-1.3 (m, 3H), 1.4-1.5 (d, 3H), 1.6-2.0 (m, 2H), 2.0-2.4 (m, 3H), 2.4-2.8 (m, 2H), 4.4-4.6 (m, 1H), 5.0-5.5.2 (m, 4H), 5.3 (s, 1H), 5.4 (d, 1H), 5.6-6.0 (m, 2H), 6.8-7.6 (m, 8H), 8.4 (s, 1H). MS: 547.2 (M+H).


25

Synthesis of Formula I compounds 21 and 22 by RCM cyclization of intermediate 20.

To a solution of 20 (70mg, 0.13mmol) in 100ml toluene-methylene chloride (1:1) was added
 5 tricyclohexylphosphine[1,3-bis(2,4,6-trimethylphenyl)-4,5-dihydroimidazol-2-ylidene]benzylidene]ruthenium (IV) dichloride (70mg). The reaction mixture was refluxed for 2 hrs at 65°C, then evaporated to an oil which was purified
 10 by chromatography on silica gel with EtOAc:hexane (1:1) to collect 21 as the first eluting spot (26mg, 0.05mmol) ¹HNMR (300MHz, CDCl₃) 0.7-0.9 (dd, 6H), 1.1-1.3 (m, 4H), 1.4-1.6 (m, 4H), 1.8-2.0 (m, 1H), 2.2-3.0 (m, 4H), 3.2 (m, 1H), 4.4-4.6 (m, 1H), 5.2 (s, 1H), 5.4 (d, 1H), 5.6-5.8 (m, 2H),
 15 6.9-7.6 (m, 8H), 7.8 (s, 1H). MS: 519.5 (M+H), 541.5 (M+Na). Succinamide 22 was collected as the second eluting spot (10mg, 0.02mmol) ¹HNMR (300MHz, CDCl₃) 0.8-1.0 (m, 6H), 1.2-1.4 (m, 4H), 1.5 (d, 3H), 1.7 (m, 1H), 1.8-2.0 (m, 1H), 2.2-2.4 (m, 1H), 2.6-2.8 (m, 3H), 3.1-3.3 (m, 1H),
 20 4.4-4.6 (m, 1H), 5.3 (s, 1H), 5.4 (d, 1H), 5.6-5.8 (m, 2H), 6.9-7.8 (m, 9H). MS: 519.5 (M+H), 541.5 (M+Na).

Example 24


1-[3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-(4-trifluoromethylphenyl)-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclopentanecarboxylic amide.



¹HNMR (300 MHz, CDCl₃) 0.8-1.0 (dd, 6H), 1.2-2.0 (m, 10H),
 30 2.4-2.6 (m, 2H), 3.4 (s, 3H), 5.3 (s, 1H), 5.5 (d, 1H),
 7.2-7.8 (m, 8H), 8.1 (s, 1H). MS: 521.5 (M+H).

Example 25

1-[3-methyl-1-[1,3-dihydro-1-(2-cyclopropylethyl)-2-oxo-5-(2-fluorophenyl)-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclopent-3-enecarboxylic amide.




Following the procedures in Scheme 16, the following Formula I compounds were prepared. Both examples eluted as the first isomer upon chromatography on silica gel using 1:1 EtOAc:hexane.

¹HNMR (300 MHz, CDCl₃) 0.0 (m, 2H), 0.2-0.5 (m, 2H), 0.5-0.6 (m, 1H), 0.8-1.0 (dd, 6H), 1.2-1.8 (m, 4H), 1.8-2.0 (m, 1H), 2.2-3.0 (m, 4H), 3.2-3.3 (m, 1H), 3.6-3.8 (m, 1H), 4.4-4.6 (m, 1H), 5.3 (s, 1H), 5.5 (s, 1H), 5.6-5.8 (m, 2H), 7.0-7.8 (m, 8H), 7.9 (s, 1H). MS: 545.5 (M+H), 567.4 (M+Na).

Example 26

1-[3-methyl-1-[1,3-dihydro-1-(2-methylpropyl)-2-oxo-5-(2-fluorophenyl)-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclopent-3-enecarboxylic amide.




Following the procedures in Scheme 16, the following
Formula I compounds were prepared. Both examples eluted as
5 the first isomer upon chromatography on silica gel using
1:1 EtOAc:hexane.

¹HNMR (300 MHz, CDCl₃) 0.67 (d, 3H), 0.80 (d, 3H), 0.87 (d,
3H), 0.93 (d, 3H), 1.1-1.3 (m, 1H), 1.6-2.0 (m, 2H), 2.2-
10 3.0 (m, 4H), 3.2 (d, 1H), 3.4-3.6 (m, 1H), 4.2-4.4 (m, 1H),
5.3 (s, 1H), 5.5 (d, 1H), 5.6-5.8 (m, 2H), 6.9-7.8 (m, 8H),
7.9 (s, 1H). MS: 533.3 (M+H), 555.4 (M+Na).


Example 27

15 1-[3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-(4-
chlorophenyl)-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-
cyclobutanecarboxylic amide.



20

Synthesis of Formula I compounds XVIa and XVIb (Scheme 18)
Scheme 18



Step 1. Preparation of diester X (Scheme 18)

A 2.5M solution of *n*-butyllithium in hexanes (4mL) was added dropwise to a -78°C solution of DIEA (10mmol) in 40mL dry THF under N₂. The mixture was stirred at 0°C for 30min, then cooled to -78°C. Ethyl cyclobutanecarboxylate (1.26mL, 9.1mmol) was added dropwise and the mixture was stirred at -78°C for 45min. *tert*-Butyl bromoacetate (1.4mL, 9.5mmol) was added and the mixture was stirred at RT for 4hrs. Ethanol (0.5mL) was added and the reaction mixture was concentrated *in vacuo*, then partitioned between CH₂Cl₂ and cold 0.5N HCl. The organic layer was dried over MgSO₄,

filtered and evaporated in vacuo, resulting in 1.8g of a dark amber oil (intermediate X) which was used without further purification. $^1\text{H-NMR}$ (300MHz, CDCl_3): δ 4.15 (quart, 2H), 2.77 (s, 2H), 2.48 (m, 2H), 1.94 (m, 4H),
5 1.4 (s, 9H), 1.26 (t, 3H).

Step 2. Deprotection of diester X to give free acid XI.
(Scheme 18)

TFA (15mL) was added to a solution of 1.2g (4.95mmol)
10 of X in 15mL CH_2Cl_2 and the mixture stirred for 3hrs. The mixture was concentrated in vacuo. Toluene was added and the mixture was again concentrated in vacuo. The residue was partitioned between Et_2O and 5% aq. NaHCO_3 . The Et_2O layer was washed with 5% aq. NaHCO_3 and the aqueous layers
15 were combined and carefully acidified to a pH between 3-4 using conc. HCl. This mixture was extracted twice with EtOAc . The EtOAc layers were combined, dried (MgSO_4) and evaporated in vacuo to give 875mg (95% yield) of acid XI. $^1\text{H-NMR}$ (300MHz, CDCl_3): δ 4.18 (quart, 2H), 2.91 (s, 2H),
20 2.55 (m, 2H), 2.0 (m, 4H), 1.26 (t, 3H).

Step 3. Synthesis of benzyl ester XII. (Scheme 18)

K_2CO_3 (4.48g, 32.4mmol) was added to a solution of 2.74g (14.71mmol) of XI in 50mL DMF. Benzyl bromide
25 (1.92mL, 16.18mmol) was added and the reaction mixture was stirred at room temperature for 16hrs. The mixture was partitioned between Et_2O and water. The water layer was washed with Et_2O and the organic layers were combined, dried (MgSO_4) and evaporated in vacuo. The residue was
30 purified by flash chromatography on silica gel (100% Hexane to 5% EtOAc /Hex gradient) to give intermediate XII. $^1\text{H-NMR}$ (300MHz, CDCl_3): δ 7.33 (m, 5H), 5.09 (s, 2H), 4.1 (quart, 2H), 2.9 (s, 2H), 2.54 (m, 2H), 1.97 (m, 4H), 1.19 (t, 3H).

35 Step 4. Alkylation of diester XII. (Scheme 18)

3.86mL of a 1M LiHMDS in THF solution was added to a cooled (-78°C) solution of intermediate XII (970mg, 3.51mmol) in 10mL dry THF. After 15min., 0.39mL (3.86mmol)

of 3-bromo-2-methylpropene was added and the reaction mixture was stirred for 16hrs at RT. The mixture was partitioned between Et₂O and water. The water layer was washed with Et₂O and the organic layers were combined, dried (MgSO₄) and evaporated in vacuo. 1.11g of product XIII of suitable purity was isolated. ¹H-NMR(300MHz, CDCl₃): δ 7.28(m,5H), 5.04(ABquart,2H), 4.62(m,2H), 4.05(m,2H), 3.0(dd,1H), 2.44-2.2(m,4H), 2.1-1.6(m,4H), 1.65(s,3H), 1.16(t,3H).

Step 5. Preparation of free acid XIV. (Scheme 18)

To a degassed solution of intermediate XIII in 20mL of ethanol in a Parr bomb was added 100mg of 5% Pd on carbon, and the mixture was agitated under 50psi of H₂ for 2hrs.

The reaction mixture was filtered through celite and washed with ethanol. The alcoholic filtrate was concentrated in vacuo and the residue partitioned between CH₂Cl₂ and 1N NaOH. The layers were separated and the basic layer was acidified using conc. HCl. The acidic mixture was extracted twice with CH₂Cl₂ and these organic layers were combined, dried (MgSO₄) and evaporated in vacuo to give intermediate XIV. ¹H-NMR(300MHz, CDCl₃): δ 4.2(m,2H), 2.85(dd,1H), 2.5-2.25(m,3H), 2.15-1.5(m,5H), 1.27(t,3H), 1.05(m,1H), 0.9(m,6H).

Step 6. Coupling of acid XIV with aminobenzodiazepine A to give succinamide Xva. (Scheme 18)

To a solution of XIV (218mg, 0.9mmol) in 5mL of DMF were added HATU (380mg, 1mmol), aminobenzodiazepine hydrobromide A (362mg, 0.95mmol), and DIEA (0.35mL, 2mmol). The reaction mixture was stirred at room temperature for 4hrs, then partitioned between EtOAc and water. The organic layer was dried (MgSO₄) and evaporated in vacuo. The residue was purified by flash chromatography on silica gel (EtOAc/Hexane gradient) to give 183mg of product Xva. ¹H-NMR(300MHz, CDCl₃): δ 7.6-7.5(m,4H), 7.4-7.3(m,4H), 7.25-7.2(m,1H), 5.53(d,1H), 4.23(m,2H), 3.45(s,3H), 2.85(dd,1H), 2.5-2.4(m,3H), 2.38-2.27(m,1H), 2.0-1.77(m,3H), 1.55(m,1H),

1.3 (t, 3H), 0.99 (m, 1H), 0.9 (dd, 6H). MS: APcI (M+H)⁺ = 524.1.


Step 7. Synthesis of Formula I compound XVIa. (Scheme 18)

5 To a solution of intermediate XVa (183mg, 0.35mmol) in 5mL of THF was added a solution of LiOH.H₂O (30mg, 0.7mmol) in 2mL of water. 1mL of MeOH was added and the mixture was stirred at room temperature for 16hrs. The mixture was partitioned between EtOAc and 1N HCl. The aqueous layer
10 was washed with EtOAc and the organic layers were combined, dried (MgSO₄) and evaporated in vacuo. The residue was dissolved in 9mL of DMF, and HATU (138mg, 0.36mmol) and DIEA (0.063mL, 0.36mmol) were added. NH₃ gas was bubbled into the mixture for 30min., then the flask was stoppered
15 and stirred at room temperature for 72hrs. The mixture was partitioned between EtOAc and water. The organic layer was dried (MgSO₄) and evaporated in vacuo. The residue was purified by flash chromatography on silica gel (10% to 40% EtOAc/Hexane gradient) to give 28mg of product XVIa. ¹H-
20 NMR: • 7.83 (br s, 1H), 7.65-7.51 (m, 4H), 7.42-7.34 (m, 4H), 7.26 (m, 1H), 5.48 (d, 1H), 5.33 (br s, 1H), 3.48 (s, 3H), 2.89-2.80 (m, 2H), 2.4-2.35 (m, 2H), 2.04-1.6 (m, 5H), 1.26 (m, 1H), 0.93 (dd, 6H). MS: APcI (M+H)⁺ = 495.1.

25

Example 28

1-[3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-yl]carbamoyl]-butyl]-cyclobutanecarboxylic amide.



Following the procedures of Example 27 and disclosed in Scheme 18, the title compound was prepared.

Step 8. Synthesis of succinamide ester XVb

5 To a solution of intermediate XIV (250mg, 1.03mmol) in 5mL of DMF was added HATU (418mg, 1.1mmol), aminobenzodiazepine hydrobromide B (380mg, 1.1mmol), and DIEA (0.38mL, 2.2mmol). The reaction mixture was stirred at room temperature for 16hrs, then partitioned between
10 EtOAc and water. The organic layer was dried (MgSO₄) and evaporated in vacuo. The residue was purified by flash chromatography on silica gel (EtOAc/Hexane gradient) to give 122mg of product XVb. MS: APcI (M+H)⁺ = 490.1.


15 Step 9. Synthesis of Formula I compound XVIb

To a solution of intermediate XVb (122mg, 0.25mmol) in 3mL of THF was added a solution of LiOH.H₂O (42mg, 1mmol) in 2mL of water. 1mL of MeOH was added and the mixture was stirred at room temperature for 4hrs. The mixture was
20 partitioned between EtOAc and 1N HCl. The aqueous layer was washed with EtOAc and the organic layers were combined, dried (MgSO₄) and evaporated in vacuo. The residue was dissolved in 5mL of DMF, and HATU (103mg, 0.27mmol) and DIEA (0.052mL, 0.3mmol) were added. NH₃ gas was bubbled
25 into the mixture for 30min., then the flask was stoppered and stirred at room temperature for 72hrs. The mixture was partitioned between EtOAc and water. The organic layer was dried (MgSO₄) and evaporated in vacuo. The residue was purified by flash chromatography on silica gel
30 (EtOAc/Hexane gradient) to give 8mg of product XVIb. ¹H-NMR: δ 7.88(br s, 1H), 7.61-7.34(m, 9H), 7.24(m, 1H), 5.48(d, 1H), 5.33(br s, 1H), 3.46(s, 3H), 2.88-2.8(m, 2H), 2.43-2.28(m, 2H), 2.02-1.6(m, 5H), 1.25(m, 1H), 0.92(dd, 6H). MS: APcI (M+H)⁺ = 461.1

35


Example 29

1-[3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-cycloheptyl-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclopent-3-enecarboxylic amide.



5


Step 1. Preparation of 2-aminophenyl cycloheptyl ketone 3



To magnesium (8.26 g, 0.34 mol) was added ether (30 mL), followed by addition of 2 (50 g, 0.28 mol) in ether (200 mL) at a rate that maintained the reflux over 2 h. The reaction mixture was then heated at reflux for 20 h. A solution of 1 (11.8 g, 0.10 mol) in ether (100 mL) was added to the reaction mixture slowly with vigorous stirring. After refluxing for 3 h, the reaction mixture was cooled to 0°C. 10% aq. HCl (300 mL) was added slowly and stirring was continued overnight. To the reaction mixture was added 30% aq. NaOH at 0 °C to pH-11. The resulting mixture was filtered through a pad of celite. The aqueous layer was extracted with ethyl acetate (3x200 mL). The extracts were combined and washed with brine, then dried over Na₂SO₄. After evaporation of the solvent, the residue was purified on silica gel, using 10% ethyl acetate-hexane, to afford 3 (16.43 g, 76%) as an orange oil. MS m/z 218.4 (MH⁺).

25


Step 2. Preparation of Cbz-protected aminobenzodiazepine 6



To a solution of 4 (27.09 g, 83 mmol) in THF (anhydrous, 200 mL) at 0 °C was added oxalyl chloride (7.24 mL, 83 mmol) via syringe over 5 min., followed by DMF (anhydrous, 0.24 mL). Stirring was continued for 2 h at 0 °C. To the above yellow reaction mixture was added a solution of 3 (16.40 g, 75.5 mmol) and *N*-methylmorpholine (18.26 mL, 166 mmol) in dry THF (100 mL) over ca. 20 min. Precipitates were formed and the reaction mixture was slowly warmed to room temperature and stirred for 2 h. The precipitates were filtered and washed with THF (50 mL). Ammonia was passed through the filtrate for 30 min. with stirring at room temperature. The reaction mixture was diluted with methanol (150 mL), saturated with ammonia and stirred for 1 h. After evaporation of solvents, the residue was dissolved in ethyl acetate (1200 mL) and washed with NaOH (1N, 2 x 200 mL). The aqueous layer was extracted with ethyl acetate (2 x 200 mL). The extracts were combined and washed with water (4 x 100 mL), brine (150 mL), and dried (K_2CO_3). Evaporation of the solvent gave a yellow solid 5. The crude solid 5 was dissolved in acetic acid (400 mL), followed by addition of ammonium acetate (20 g). The reaction mixture was stirred at ambient temperature for 24 h, and then concentrated. To the residue was added ethyl acetate (300 mL) and aqueous NaOH (1N) until pH>8, cooled in an ice bath. The solid was filtered, washed with water and cold ether sequentially,


and dried in air overnight to give a white solid **6** (12 g, 40% over three steps). MS m/z 406.5 (MH^+).

Step 3. N-Methylation of benzodiazepine **6**




To a mixture of **6** (1.0 g, 2.47 mmol) and potassium carbonate (0.68 g, 4.93 mmol) in dry DMF (10 mL) was added iodomethane (0.46 mL, 7.41 mmol). The reaction mixture was stirred at ambient temperature for 20 h, then poured into water (200 mL). The precipitates were filtered, washed with water and dried in air to afford **7** (1.01 g, 97%) as a white solid. MS m/z 420.4 (MH^+).

Step 4. Synthesis of aminobenzodiazepine **8**



The crude product **7** (1.0 g, 2.4 mmol) from the above reaction was dissolved in methylene chloride (10 mL), treated with anhydrous HBr gas for 1 h until disappearance of **7** on TLC. The reaction mixture was diluted with ether (100 mL) and decanted. The precipitates were washed with ether and decanted (3 x 100 mL). The remaining ether was removed in vacuo and the residue was dried under vacuum to afford **8** (0.88 g, 100%) as a white solid. MS m/z 366.0 (MH^+), 286.1 ($MH^+ - HBr$).

Step 5. Coupling of 8 with diallylsuccinate 9




5 8 (230 mg, 0.63 mmol), 9 (195 mg, 0.63 mmol) and 1-hydroxybenzotriazole hydrate (HOBT, 102 mg, 0.76 mmol) were dissolved in CH_2Cl_2 and cooled to 0 °C, then 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC, 242 mg, 1.26 mmol) and triethylamine (0.22 mL, 1.58

10 mmol) were added. After stirring for 2 days at ambient temperature, the reaction mixture was diluted with ethyl acetate, washed with water and brine then dried (Na_2SO_4). After evaporation of the solvent, the residue was purified on silica gel (increasing gradient of 20-50% ethyl


15 acetate/hexane) to afford product 10 (268 mg, 29%). MS m/z 578.6 (MH^+).

Step 6. Preparation of succinate free acid 11




10 (268 mg, 0.46 mmol) was dissolved in $\text{CH}_2\text{Cl}_2/\text{TFA}$ (2 mL, 1:1) and stirred for 5 h at ambient temperature. Evaporation of the solvent gave product **11** (258 mg) as a sticky oil which was used for next step without
5 purification.

Step 7. Synthesis of diallylsuccinate **12**



11 (240 mg, 0.46 mmol) and HATU (350 mg, 0.92 mmol) were dissolved in DMF (2 mL) and diisopropylethylamine (0.16 mL, 0.92 mmol) was added. The mixture was stirred
15 for 15 min at ambient temperature, and then treated with anhydrous ammonia for 20 min. Stirring was continued overnight. DMF was removed *in vacuo*, and the residue was diluted with ethyl acetate, washed with water and brine, then dried over MgSO_4 . After evaporation of the solvent,
20 the residue was purified on silica gel (5% methanol/methylene chloride) to afford product **12** as a white solid. **12**, a mixture of two diastereomers, was then separated on silica gel, using increasing gradient of 20-50% ethyl acetate-methylene chloride, to give first eluting
25 isomer **12a** (43 mg, 18%) and the second eluting isomer **12b** (61 mg, 25%) both as a white solids. **12a**: MS m/z 521.2 (MH^+). **12b**: MS m/z 521.2 (MH^+).


Step 8. Synthesis of Formula I compound **13**.



To a solution of **12a** (30 mg, 0.06 mmol) in CH₂Cl₂-toluene (4 mL, 1:1) was added tricyclohexylphosphine[1,3-bis(2,4,6-trimethylphenyl)-4,5-dihydroimidazol-2-ylidene][benzylidene]ruthenium (IV) dichloride (5 mg, 0.006 mmol), and the reaction mixture was heated at 60 °C for 4 h. The reaction mixture was concentrated and the residue was purified on silica gel, using 40% ethyl acetate-methylene chloride, to afford **13** (19 mg, 63%) as a white solid. ¹H NMR (300 MHz, CDCl₃) δ : 0.91 (dd, J = 16.8, 6.2 Hz, 6H), 1.18-2.04 (m, br, 15H), 2.25-2.38 (m, 1H), 2.60-2.71 (m, 2H), 2.72-2.83 (m, 1H), 2.95-3.05 (m, 1H), 3.18-3.28 (m, 1H), 3.4 (s, 3H), 5.23-5.35 (m, 2H), 5.62-5.76 (m, 2H), 7.20-7.42 (m, 3H), 7.45-7.60 (m, 2H), 7.86 (s, br, 1H); MS m/z 493.5 (MH⁺).


Example 30

1-[3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-(2-trifluoromethylphenyl)-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclopent-3-enecarboxylic amide



Example 30 can be prepared by the methods disclosed herein using the intermediates disclosed in Scheme 22

Scheme 22



5 Preparation of Compound 3, Scheme 22. To a solution of compound 1 (87.7 g, 453 mmol) in Et_2O (910 mL) at -78°C

under a nitrogen atmosphere was added *t*-BuLi (1.7 M solution in pentane, 605 mL) while maintaining a solution temperature of -78 °C. The solution was allowed to warm to -10 °C and maintained at that temperature for 3 h. A solution of 2 (79 g, 453 mmol) in Et₂O (100 mL) was added and the solution was stirred at -10 °C for an additional 2 h. The resulting solution was allowed to warm to room temperature while stirring for 17 h. An aqueous solution of saturated NH₄Cl (400 mL) was added and the layers were separated. Acid (1 N HCl) was added until a pH = 1 was obtained, the aqueous layer was extracted with ethyl acetate (3 x 200 mL). The organic extracts were combined, washed with brine, filtered, dried over anhydrous Na₂SO₄, and concentrated to yield a orange oil. Addition of a 1/1 (v/v) solution of Et₂O/Hexanes resulted in precipitation of 3 (86 g 65%) as a tan solid that was used without further purification: ¹H NMR (300 MHz, CDCl₃) 7.89-6.52 (m, 10 H); ESI MS *m/z* = 294 [C₁₅H₁₀F₃NO₂+H]⁺.

Preparation of Compound 4, Scheme 22. A solution of compound 3 (23.2 g, 79.2 mmol) and LiOH (16.5 g, 0.4 mole) in a 1/1 (V/V) of dioxane/H₂O (250 mL each) was refluxed for 18 h. Methylene chloride was added to the solution and the layers were separated. The aqueous layer was extracted with additional CH₂Cl₂ (2 x 100 mL). The organic layers were combined, dried over anhydrous Na₂SO₄, filtered, and concentrated to yield 4 (12.5 g, 59%) as a yellow oil that was used without further purification: ¹H NMR (300 MHz, CDCl₃) 9.42 (s, 2 H), 7.89-6.52 (m, 9 H); ESI MS *m/z* = 268 [C₁₄H₁₂F₃NO+H]⁺.

Preparation of Compound 5, Scheme 22. To a solution of compound 4 (12.4 g, 46.4 mmol), in CH₂Cl₂ (300 mL) was added MnO₂ (21.1 g, 0.232 mole) and the resulting heterogenous solution was stirred for 14 h at room temperature, filtered through celite, and concentrated to yield an dark orange oil. This residue was further purified

by column chromatography (silica gel, 80:20 Hexanes/EtOAc) to yield **5** (6.68 g, 54%) as a yellow solid: ^1H NMR (300 MHz, CDCl_3) 7.82-6.35 (m, 10 H); ESI MS m/z = 266 $[\text{C}_{14}\text{H}_{10}\text{F}_3\text{NO}+\text{H}]^+$.

5

Preparation of compound **6**, Scheme 22. To a solution of **5** (32.8 g, 0.12 mole) at 0 °C in CH_2Cl_2 was added bromoacetyl bromide (27.5 g, 0.14 mole). A 10% aqueous solution of Na_2CO_3 (150 mL) was added slowly with stirred for 30 min. The organic layer was separated, washed with water, dried over anhydrous Na_2SO_4 , filtered, and concentrated to yield **6** (35.1 g, 73%) as a cream colored solid: ^1H NMR (300 MHz, CDCl_3) 8.88 (d, J = 7.1 Hz, 1 H), 7.83-7.04 (m, 8 H), 4.04 (s, 2 H); ESI MS m/z = 386 $[\text{C}_{16}\text{H}_{11}\text{BrF}_3\text{NO}+\text{H}]^+$.

15

Preparation of Compound **8**, Scheme 22. Ammonia (500 mL) was condensed with a dry ice condensor and maintained at -78 °C. To this was added a solution of **6** (35 g, 91 mmol) in CH_2Cl_2 (600 mL). After 8 h, the dry ice condensor was removed and the reaction was warmed to room temperature allowing the ammonia to evaporate. The remaining solution was washed with brine, dried over anhydrous Na_2SO_4 , filtered and concentrated to yield a white solid. This solid was dissolved in EtOH (500 mL) and AcOH (15 mL), and the resulting solution was refluxed for 24 h, and concentrated to yield a yellow-red oil. Addition of Et_2O (300 mL) to this oil resulted in the precipitation of **8** (13 g, 47%) as a white powder: ^1H NMR (300 MHz, CDCl_3) δ 9.32 (s, 1 H), 7.83-7.05 (m, 8 H), 4.39 (s, 2 H); ESI MS m/z = 305 $[\text{C}_{15}\text{H}_{11}\text{F}_3\text{N}_2\text{O}+\text{H}]^+$.

25

30

Preparation of Compound **9**, Scheme 22. To a solution of **8** (3.4 g, 11.2 mmol) and Cs_2CO_3 (5.46 g, 16.6 mmol) in DMF (30 mL) was added MeI (1.0 mL, 16.6 mmol) and the solution was stirred at room temperature for 4 h. At this

35

time, the solution was diluted with ethyl acetate, (200 mL) washed with aqueous 5% LiCl (3 x 100 mL), brine, dried over Na₂SO₄, filtered, and concentrated to yield **9** (3.1 g, 87%) as a pale blue solid: ¹H NMR (300 MHz, CDCl₃) δ 7.82-6.97 (m, 8 H), 4.42 (q_{ab}, J = 6.5 Hz, 2 H), 3.42 (s, 3 H); ESI MS m/z = 319 [C₁₇H₁₃F₃N₂O+H]⁺.

Preparation of Compound **10**, Scheme 22. To a solution of **9** (2.4 g, 7.8 mmol) in THF (56 mL) at -78 °C was added KO-*t*-Bu (15.8 mL, 1 M solution in THF) and the solution was stirred for 5 min. To this solution was added a solution of trisyl azide (3.1 g, 8.7 mmol) in THF (27 mL) at -78 °C. After stirring for 5 min at -78 °C, AcOH (2.1 g, 34.7 mmol) was added and the solution was allowed to warm to room temperature while stirring for 18 h. This solution was diluted with CH₂Cl₂ (200 mL), washed with sat. NaHCO₃ (2 x 100 mL), brine, dried over anhydrous Na₂SO₄, filtered, and concentrated to yield a yellow powder. This powder was triturated with hexanes/EtOAc 5/1 (V/V) to yield 2.1 g of a white powder. This solid was dissolved in THF (50 mL) and H₂O (5 mL). Triphenylphosphine 4.6 g, 17.5 mmol) was added and the solution was stirred at room temperature for 24 h. The solution was concentrated under reduced pressure. The residue was dissolved in Et₂O (100 mL) and extracted with 1 N HCl (2 x 100 mL). The acidic extracts were adjusted to pH 14 with 1 N NaOH and extracted with CH₂Cl₂ (3 x 75 mL). The CH₂Cl₂ extracts were washed with brine (1 x 100 mL), dried over anhydrous Na₂SO₄, filtered, and concentrated to yield **10** (1.6 g, 62%) as pale yellow solid: ¹H NMR (300 MHz, CDCl₃) δ 7.79-6.93 (m, 8 H), 4.52 (s, 1 H), 3.47 (s, 3 H), 2.46 (br. s, 2 H); ESI MS m/z = 334 [C₁₇H₁₄F₃N₃O+H]⁺.

Preparation of **12**, Scheme 22. To a solution of **10** (380 mg, 1.1 mmol), DIPEA (1 mL, 5.7 mmol), and **11** (0.37 g, 1.2 mmol) in DMF (30 mL) was added HATU (0.43 g, 1.1 mmol) and the solution was allowed to stir for 24 h at room

temperature. The solution was diluted with EtOAc (150 mL) and washed with 5% aqueous solution of LiCl (3 x 100 mL), 1 N HCl (1 x 100 mL), saturated aqueous NaHCO₃ (2 x 100 mL), brine, dried over anhydrous Na₂SO₄, filtered, and

5 concentrated to yield a pale yellow oil. This residue was further purified by column chromatography (silica gel, 3:1 Hexanes/EtOAc) to yield 12 (0.31 g, 44%) as a white foam: ¹H NMR (300 MHz, CDCl₃) δ 7.86-6.91 (m, 9 H), 5.82 (m, 2 H), 5.67 (m, 1 H), 5.09 (m, 4 H), 3.48 (s, 3 H), 2.73-2.28 (m, 10 5 H), 1.92 (m, 1 H), 1.61 (m, 1 H), 1.42 (s, 9 H), 1.27 (m, 1 H), 0.93 (m, 6 H); ESI MS m/z = 626 [C₃₅H₄₂F₃N₃O₄+H]⁺.

Preparation of 13, Scheme 22. To a solution of 12 (300 mg, 0.5 mmol) in CH₂Cl₂ (4 mL) was added TFA (10 mL) 15 and the solution was allowed to stir for 24 h at room temperature. The solution was concentrated under reduced pressure, the residue was redissolved in toluene and concentrated (3 x 10 mL). Ammonia gas was bubbled through a solution of the residue, DIPEA (0.4 mL, 2.5 mmol), HATU 20 (180 mg, 0.5 mmol) in DMF (5 mL) for 30 min and the solution was allowed to stir for 24 h at room temperature. The contents of the flask were partitioned between EtOAc and a 5% LiCl solution (50 mL each), the organic phase washed with 5% LiCl (3 x 50 mL), and dried over anhydrous 25 Na₂SO₄, filtered, and concentrated to yield a white solid. The diastereomers were separated by column chromatography (silica gel, 2:1 EtOAc/Hexanes) to yield Diastereomer A and Diastereomer B (99 mg, 35%) as white powders:

Diastereomer A: mp 218-219 °C; ¹H NMR (300 MHz, 30 CDCl₃) δ 8.48 (s, 1 H), 7.67-7.01 (m, 8 H), 5.79 (m, 2 H), 5.53 (d, J = 7.9 Hz, 2 H), 5.33 (s, 1 H), 5.18-5.04 (m, 4 H), 3.47 (s, 3 H), 2.57-2.38 (m, 5 H), 1.79 (m, 1 H), 1.28 (m, 2 H), 0.89 (d, J = 6.5 Hz, 3 H), 0.85 (d, J = 6.5 Hz, 3 H); IR (ATR) 3142, 2956, 1706, 1647, 1599, 1538, 1312, 1142 35 cm⁻¹; ESI MS m/z = 569 [C₃₁H₃₅F₃N₄O₃ + H]⁺; HPLC 96.9 %, t_r = 21.6 min; Optical Rotation [α]_D²⁵ +2.5° (c 0.01, Methanol).


Diastereomer B: mp 200-202 °C; ¹H NMR (300 MHz, CDCl₃) δ 8.42 (s, 1 H), 7.67-7.05 (m, 8 H), 5.73 (m, 2 H),

5.59 (d, $J = 8.1$ Hz, 2 H), 5.33 (s, 1 H), 5.16-5.02 (m, 4 H), 3.49 (s, 3 H), 2.56-2.41 (m, 5 H), 1.73 (m, 1 H), 1.21 (m, 2 H), 0.91 (d, $J = 6.8$ Hz, 3 H), 0.87 (d, $J = 6.8$ Hz, 3 H); IR (ATR) 3308, 3150, 2928, 1712, 1650, 1603, 1310 cm^{-1} ;
5 ESI MS $m/z = 569$ [$\text{C}_{31}\text{H}_{35}\text{F}_3\text{N}_4\text{O}_3 + \text{H}$] $^+$; HPLC 98.8%, $t_r = 21.2$ min; $[\alpha]^{25}_D -7.4^\circ$ (c 0.01, Methanol).

Compound 13 may be converted to the title compound, Example 30, by using the RCM method according to the step 4 of
10 Example 1.

EXAMPLE 500

Preparation of 3-amino-8-bromo-1,5-dimethyl-1,3-dihydro-2H-1,4-benzodiazepin-2-one




15

Step 1: Preparation of 1-(2-amino-4-bromophenyl)ethanone
Under an argon atmosphere a solution of 3-bromoaniline (31.3 g, 181.8 mmol) and acetonitrile (75g, 1.818 mol) in
20 anhydrous toluene (120 ml) was added dropwise over 2.5 hours to a stirred solution of boron trichloride (23.4 g, 200 mmol) in (200 ml) hexanes cooled in an ice bath. After the addition was completed, aluminum chloride (26.6g, 200 mmol) was added portion wise over 30 minutes. The mixture
25 was allowed to warm to ambient temperature and then heated at reflux for 16 hours with stirring. The reaction mixture was then cooled to 10 °C and 100 mL of a 3N HCl solution was added dropwise with continued stirring. After the addition was complete, the mixture was heated at reflux for 3.5
30 hours, then cooled to room temperature, and the layers separated. The aqueous layer was extracted with chloroform (3 x 250 ml). Organic layers were combined, dried over magnesium sulfate, filtered, and concentrated to give the title compound (9.58g, 25 %). ^1H NMR (CDCl_3 , 300 MHz): δ

7.54 (d, 1H, J=8.8 Hz), 6.83 (d, 1H, J = 1.9 Hz), 6.75 (dd, 1H, J = 8.4, 1.8 Hz), 2.54 (s, 3H) ppm.

Step 2: Preparation of benzyl 8-bromo-5-methyl-2-oxo-2,3-dihydro-1H-1,4-benzodiazepin-3-ylcarbamate




1b

Under an argon atmosphere 1H-1,2,3-benzotriazol-1-yl{[(benzyloxy)carbonyl]amino}acetic acid (6.71 g, 20.6 mmol) was suspended in anhydrous methylene chloride (92 ml) and cooled to 0°C in an ice bath. Oxalyl chloride (2.61g, 20.6 mmol), then N,N-dimethylformamide (38 ml) were added dropwise to the suspension. The reaction mixture was stirred at 0°C in an ice bath for 30 minutes, at which point no further gas evolution was noted. Then a solution of 1-(2-amino-4-bromophenyl)ethanone (4.0g, 18.7 mmol) and 4-methylmorpholine (2.84g, 28.0 mmol) in anhydrous methylene chloride (60 ml) was added dropwise. The reaction mixture was allowed to warm to room temperature over 12 hours, then quenched with water (200 ml), and extracted with ethyl acetate (3x250 ml). The organic layers were combined, dried over magnesium sulfate, filtered and concentrated to give a semi-solid which was dissolved in tetrahydrofuran (120 ml) and methyl alcohol (35 ml). Ammonia gas was bubbled through this solution for 2.5 hours. The reaction was then concentrated to a viscous light brown oil. The oil was dissolved in acetic acid (120 ml) and ammonium acetate (4.3g, 56.1 mmol) was added in one portion and stirred for 12 hours. The reaction mixture was diluted with water (100 ml) and then made basic (pH = 10) with 25% sodium hydroxide, while stirring in an ice bath. The aqueous solution was then extracted with ethyl acetate (3 x 500 ml) and the organic layers combined, dried over magnesium sulfate, filtered and concentrated. The

resulting residue was purified on silica gel, eluting with 40% ethyl acetate in hexanes to give the title compound (4 g, 53%). ¹H NMR (CDCl₃, 300 MHz): δ 9.90 (s-br, 1H), 7.42-7.31 (m, 6H), 7.12 (d, 1H, 1.5 Hz), 7.06-7.03 (m, 1H), 5.18-5.08 (m, 3H), 2.50 (s, 3H) ppm.


Step 3: Preparation of benzyl 8-bromo-1,5-dimethyl-2-oxo-2,3-dihydro-1H-1,4-benzodiazepin-3-ylcarbamate



1c

The product of Step 2 (2.0 g, 4.98 mmol) was dissolved in anhydrous N,N-dimethylformamide (10 ml). To this solution was added potassium carbonate (1.72g, 12.44 mmol) and iodomethane (0.847g, 5.97 mmol), and the reaction mixture was sealed in a pressure flask and stirred for 12 hours at room temperature, then diluted with water and ethyl acetate (20/70 ml). The aqueous solution was then extracted with ethyl acetate (3 x 20 ml). The organic layers were combined, washed with water (1 x 100 ml), dried over magnesium sulfate, filtered and concentrated to give the title compound (1.58 g, 77.5%). ¹H NMR (CDCl₃, 300 MHz): δ 7.43-7.28 (m, 7H), 6.68 (d, 1H, J = 8.1 Hz), 5.15-5.05 (m, 3H), 3.38 (s, 3H), 2.45 (d, 3H, 1.5 Hz) ppm.

Step 4: Preparation of 3-amino-8-bromo-1,5-dimethyl-1,3-dihydro-2H-1,4-benzodiazepin-2-one




1d

The product of Step 3 (0.831 g, 2.00 mmol) was dissolved in anhydrous anisole (16 ml) and then methanesulfonic acid (3.84g, 40 mmol) was added in one portion. The reaction mixture was heated to 40°C for 30 minutes with stirring, then cooled to 0°C in an ice bath and made basic (pH = 10) with concentrated ammonium hydroxide. The aqueous solution was then extracted with chloroform (3 x 50 ml) and the organic layers combined, dried over magnesium sulfate, filtered and concentrated to give the crude product. Purification on silica gel, eluting with 10% methyl alcohol in chloroform, providing the title compound as an XXXX (0.463 g, 82%). ¹H NMR (CDCl₃, 300 MHz): δ 7.32-7.23 (m, 3H), 4.12 (d, 1H, J = 1.1 Hz), 3.27 (s, 3H), 2.30 (d, 3H, J = 1.5 Hz) ppm.

Example 501

Scheme 23



Preparation of compound 20, Scheme 23.

A three neck 3L round bottom flask was charged with **19** (100 g, 876 mmol), benzylbromide (104 mL, 876 mmol) and DMF (876

mL). The reaction was cooled to 0 °C and to the reaction was added solid K₂CO₃ (170 g, 1.2 mol) in portions. The ice bath was removed and the reaction was stirred vigorously for 2 h. To the reaction was added H₂O (1800
5 mL). The aqueous layer was extracted with Hexanes (3 x 1000 mL). The organic layers were combined and dried over MgSO₄. The solution was filtered and the solvent was removed under reduced pressure to afford **20** (174 g, 97%) as a clear oil: ¹H NMR (CDCl₃) δ 7.34 (m, 5 H), 5.12 (s, 2 H),
10 2.79 (m, 1 H), 1.92-1.57 (m, 8 H).

Preparation of compound 21, Scheme 23.

A 2L round bottom flask equipped with a mechanical stirring apparatus was charged with 1 M LiHMDS in hexanes (860 mL,
15 860 mmol) and anhydrous THF (302 mL). The solution was cooled to -78 °C and treated with **20** in THF (100 mL) added slowly through an addition funnel. After 1h, allylbromide (90 mL, 1.0 mol) was added slowly to the reaction. The reaction was warmed slowly to rt and then stirred for 12 h.
20 The reaction was poured into 1 N HCl (1000 mL) and stirred for 10 min. The layers were separated. The organic layer was washed with 1 N HCl (100 mL), aq. NaHCO₃ (500 mL) and brine (500 mL). The solution was dried over MgSO₄ and filtered. The solvent was removed under reduced pressure
25 to afford **21** (189 g, 95%) as a pale yellow oil: ¹H NMR (CDCl₃) δ 7.35 (m, 5 H), 5.77-5.63 (m, 1 H), 5.11 (s, 2 H), 5.02 (m, 1 H), 4.97 (m, 1 H), 2.38 (m, 2 H), 2.11 (m, 2 H), 1.68-1.51 (m, 6 H).

30 Preparation of compound 22, Scheme 23.

A solution of RuO₂·H₂O (2.9 g, 22 mmol) and NaIO₄ (44 g, 205 mmol) in 1:1 CHCl₃-H₂O (820 mL) was shaken in a separatory funnel. The layers were separated. The organic layer was poured into a 5L 3N round bottom flask equipped with a
35 mechanical stirring apparatus. To the reaction was added a solution of **21** (100 g, 410 mmol) in acetone (410 mL). A separate solution of NaIO₄ (569 g, 2.67 mol) was added in

portions to a stirred solution of the reactant components. The resulting slurry was stirred vigorously for 24 h.

5 The reaction was filtered through a 1L Buchner funnel to remove the excess NaIO_4 remaining in the reaction. The filtrate was diluted with Et_2O (400 mL) and 1 N NaHSO_4 (400 mL). The organic layer was separated and washed with 1 N NaHSO_4 (1 x 400 mL). The organic layer was diluted with Et_2O (1100 mL) and filtered through a Buchner funnel to
10 afford **22** (66 g, 62%) as a light purple oil: ^1H NMR (CDCl_3) δ 7.34 (m, 5 H), 5.14 (s, 2 H), 3.30 (s, 2 H), 2.31-2.16 (m, 2 H), 1.81-1.50 (m, 6 H).

Preparation of compound **24**, Scheme 23.

15 To **22** (62 g, 238 mmol) in CH_2Cl_2 (238 mL) at rt was added oxalyl chloride (28 mL, 310 mmol). The reaction was stirred for 2 h until gas evolution subsided. The solvent was removed under reduced pressure to provide the intermediate acid chloride.

20 A 3L 3N round bottom flask equipped with a mechanical stirring apparatus was charged with **23** (41 g, 230 mmol) and anhydrous THF (700 mL). The solution was cooled to -78°C and treated with 2.5 M $n\text{-BuLi}$ in hexanes (92 mL, 230 mmol)
25 added through an addition funnel. After 20 min, the acid chloride intermediate in THF (67 mL) was added slowly to the solution. After addition was complete, the reaction was warmed to rt and stirred for 2 h.

30 The reaction was poured into sat. NH_4Cl (500 mL). The layers were separated. The aqueous layer was extracted with Et_2O (1000 mL). The organic layers were combined and washed with NaHCO_3 (500 mL) and brine (500 mL). The solution was dried over MgSO_4 and filtered. The solvents
35 were removed under reduced pressure. The crude material was purified by SiO_2 chromatography (CH_2Cl_2) to afford **24** (68 g, 68%) as an off-white solid: ^1H NMR (CDCl_3) δ 7.40-7.12 (m, 10 H), 5.15 (s, 2 H), 4.49 (m, 1 H), 4.13 (m, 2

H), 3.34 (m, 2 H), 3.15 (dd, $J = 13.4, 3.2$ Hz, 1 H), 2.65 (dd, $J = 13.4, 9.6$ Hz, 1 H), 2.29 (m, 2 H), 1.85-1.50 (m, 6 H).

5 Preparation of compound 25, Scheme 23.

A 3L 3N round bottom flask equipped with a mechanical stirring apparatus was charged with 24 (32 g, 78 mmol) and anhydrous THF (155 mL). The reaction was cooled to -60°C and treated with 1 M LiHMDS in hexanes (78 mL, 78 mmol).

10 The reaction was warmed to -20°C and stirred for an 4 h.

To a 1L 3N round bottom flask was added hexane (516 mL), 2,6-lutidine (38 mL, 326 mmol) and methallyl alcohol (26 mL, 311 mmol). The solution was stirred vigorously using a
15 mechanical stirring apparatus. The solution was cooled to -20°C and treated with TiF_2O (52 mL, 311 mmol). During the addition of TiF_2O , a white solid precipitated from the solution. After 1 h, the contents of the flask were poured into a Buchner funnel and transferred to a pre-cooled (-40°C)
20 1000 mL round bottom flask. The solution was carefully transferred to the main reaction via cannula. Stirring was continued for 18 h at -20°C .

The reaction was poured into a stirred solution of NH_4Cl
25 (1000 mL). The layers were separated. The aqueous layer was extracted with EtOAc. The organic layers were combined, washed with 10% aq. citric acid (400 mL) and brine (400 mL). The organic layer was dried over MgSO_4 and filtered. The solvents were removed under reduced pressure
30 to afford 25 (28 g, 87%) as viscous oil: ^1H NMR (CDCl_3) δ 7.41-7.15 (m, 10 H), 5.15 (s, 2 H), 4.72 (m, 2 H), 4.58 (m, 1 H), 4.10-3.95 (m, 2 H), 3.27 (dd, $J = 13.4, 3.1$ Hz, 1 H), 2.67 (dd, $J = 13.5, 11.9$ Hz, 1 H), 2.54 (dd, $J = 13.3, 10.2$ Hz, 1 H), 2.25 (m, 2 H), 1.80-1.49 (m, 11 H).

35

Preparation of compound 26, Scheme 23.

To 24 (80 mg, 0.17 mmol) in 2:1 DME- H_2O (3 mL) was added 30% wt H_2O_2 (104 μL) and then $\text{LiOH}\cdot\text{H}_2\text{O}$ (14 mg, 0.34 mmol) in

H₂O (0.3 mL). The reaction was stirred for 72 h at rt. To the solution was added 30% NaHSO₃ (0.5 mL). The reaction was diluted with NH₄Cl (20 mL) and extracted with EtOAc (50 mL). The organic layer was dried over MgSO₄ and filtered.


- 5 The solvent was removed under reduced pressure. The crude material was purified by column chromatography (50% ethyl acetate in hexanes) to provide 26 (22 mg, 41%) as a clear, viscous oil: ¹H NMR (CDCl₃) δ 7.35 (m, 5 H), 5.14 (m, 2 H), 4.72 (d, J = 12.0 Hz, 1 H), 4.71 (s, 1 H), 3.07 (dd, J =
- 10 11.6, 2.9 Hz, 1 H), 2.43 (dd, J = 14.5, 11.9 Hz, 1 H), 2.19 (m, 2 H), 2.05 (m, 1 H), 1.90-1.50 (m, 9 H).

Table 2 demonstrates representative compounds envisaged within the scope of the present invention. Each formulae at the start of Table 2 are intended to be paired with each entry in the table which follows.


For example the compound 1-[(1R)-3,3-dimethyl-1-[6,7-dihydro-5-methyl-6-oxo-5H-dibenz[b,d]azepin-7-ylcarbamoyl]-butyl]-cyclopentanecarboxylic amide is represented by Example #500-B-y, which comprises the core B, succinate y, and entry #500.

For example the compound 1-[(1R)-3-methyl-1-[(3S)-1,3-dihydro-1-ethyl-2-oxo-5-phenyl-7-chloro-2H-1,4-benzodiazepin-3-yl]-butyl]-cyclohexanecarboxylic amide is represented by Example #510-D-ab, which comprises the core D, succinate ab, and entry #510.


Table 2




A




B




C











D







E












F



**P****Q****R****S**



















5










wherein R^3 , R^{3a} and R^5 are described, respectively, in the following moieties:

**a****b****c****d****e****f****g****h****i**

10

15

**j****k****l****m****n****o****p****q****r****s****t****u****v****w****x****y****z****aa**

abacadaeafagahaiaj

5

10

Table 2(cont.)

Ex#	core	R3/R5	R13	Z
500	A - S	a - aj	H	methyl
501	C - S	a - aj	F	methyl
502	C - S	a - aj	Cl	methyl
503	C - S	a - aj	OH	methyl
504	C - S	a - aj	-CH ₃	methyl
505	C - S	a - aj	-CH ₂ CH ₃	methyl
506	C - S	a - aj	-OCH ₃	methyl
507	C - S	a - aj	-CF ₃	methyl
508	A - S	a - aj	H	ethyl
509	C - S	a - aj	F	ethyl
510	C - S	a - aj	Cl	ethyl
511	C - S	a - aj	OH	ethyl
512	C - S	a - aj	-CH ₃	ethyl
513	C - S	a - aj	-CH ₂ CH ₃	ethyl
514	C - S	a - aj	-OCH ₃	ethyl

515	C - S	a - aj	-CF ₃	ethyl
516	A - S	a - aj	H	i-propyl
517	C - S	a - aj	F	i-propyl
518	C - S	a - aj	Cl	i-propyl
519	C - S	a - aj	OH	i-propyl
520	C - S	a - aj	-CH ₃	i-propyl
521	C - S	a - aj	-CH ₂ CH ₃	i-propyl
522	C - S	a - aj	-OCH ₃	i-propyl
523	C - S	a - aj	-CF ₃	i-propyl
524	A - S	a - aj	H	n-propyl
525	C - S	a - aj	F	n-propyl
526	C - S	a - aj	Cl	n-propyl
527	C - S	a - aj	OH	n-propyl
528	C - S	a - aj	-CH ₃	n-propyl
529	C - S	a - aj	-CH ₂ CH ₃	n-propyl
530	C - S	a - aj	-OCH ₃	n-propyl
531	C - S	a - aj	-CF ₃	n-propyl
532	A - S	a - aj	H	n-butyl
533	C - S	a - aj	F	n-butyl
534	C - S	a - aj	Cl	n-butyl
535	C - S	a - aj	OH	n-butyl
536	C - S	a - aj	-CH ₃	n-butyl
537	C - S	a - aj	-CH ₂ CH ₃	n-butyl
538	C - S	a - aj	-OCH ₃	n-butyl
539	C - S	a - aj	-CF ₃	n-butyl
540	A - S	a - aj	H	i-butyl
541	C - S	a - aj	F	i-butyl
542	C - S	a - aj	Cl	i-butyl
543	C - S	a - aj	OH	i-butyl
544	C - S	a - aj	-CH ₃	i-butyl
545	C - S	a - aj	-CH ₂ CH ₃	i-butyl
546	C - S	a - aj	-OCH ₃	i-butyl
547	C - S	a - aj	-CF ₃	i-butyl
548	A - S	a - aj	H	s-butyl
549	C - S	a - aj	F	s-butyl
550	C - S	a - aj	Cl	s-butyl

551	C - S	a - aj	OH	s-butyl
552	C - S	a - aj	-CH ₃	s-butyl
553	C - S	a - aj	-CH ₂ CH ₃	s-butyl
554	C - S	a - aj	-OCH ₃	s-butyl
555	C - S	a - aj	-CF ₃	s-butyl
556	A - S	a - aj	H	t-butyl
557	C - S	a - aj	F	t-butyl
558	C - S	a - aj	Cl	t-butyl
559	C - S	a - aj	OH	t-butyl
560	C - S	a - aj	-CH ₃	t-butyl
561	C - S	a - aj	-CH ₂ CH ₃	t-butyl
562	C - S	a - aj	-OCH ₃	t-butyl
563	C - S	a - aj	-CF ₃	t-butyl
564	A - S	a - aj	H	allyl
565	C - S	a - aj	F	allyl
566	C - S	a - aj	Cl	allyl
567	C - S	a - aj	OH	allyl
568	C - S	a - aj	-CH ₃	allyl
569	C - S	a - aj	-CH ₂ CH ₃	allyl
570	C - S	a - aj	-OCH ₃	allyl
571	C - S	a - aj	-CF ₃	allyl
572	A - S	a - aj	H	cyclopropyl
573	C - S	a - aj	F	cyclopropyl
574	C - S	a - aj	Cl	cyclopropyl
575	C - S	a - aj	OH	cyclopropyl
576	C - S	a - aj	-CH ₃	cyclopropyl
577	C - S	a - aj	-CH ₂ CH ₃	cyclopropyl
578	C - S	a - aj	-OCH ₃	cyclopropyl
579	C - S	a - aj	-CF ₃	cyclopropyl
580	A - S	a - aj	H	cyclopropyl-CH ₂ -
581	C - S	a - aj	F	cyclopropyl-CH ₂ -
582	C - S	a - aj	Cl	cyclopropyl-CH ₂ -
583	C - S	a - aj	OH	cyclopropyl-CH ₂ -
584	C - S	a - aj	-CH ₃	cyclopropyl-CH ₂ -
585	C - S	a - aj	-CH ₂ CH ₃	cyclopropyl-CH ₂ -
586	C - S	a - aj	-OCH ₃	cyclopropyl-CH ₂ -

587	C - S	a - aj	-CF ₃	cyclopropyl-CH ₂ -
588	A - S	a - aj	H	cyclobutyl
589	C - S	a - aj	F	cyclobutyl
590	C - S	a - aj	Cl	cyclobutyl
591	C - S	a - aj	OH	cyclobutyl
592	C - S	a - aj	-CH ₃	cyclobutyl
593	C - S	a - aj	-CH ₂ CH ₃	cyclobutyl
594	C - S	a - aj	-OCH ₃	cyclobutyl
595	C - S	a - aj	-CF ₃	cyclobutyl
596	A - S	a - aj	H	cyclobutyl-CH ₂ -
597	C - S	a - aj	F	cyclobutyl-CH ₂ -
598	C - S	a - aj	Cl	cyclobutyl-CH ₂ -
599	C - S	a - aj	OH	cyclobutyl-CH ₂ -
600	C - S	a - aj	-CH ₃	cyclobutyl-CH ₂ -
601	C - S	a - aj	-CH ₂ CH ₃	cyclobutyl-CH ₂ -
602	C - S	a - aj	-OCH ₃	cyclobutyl-CH ₂ -
603	C - S	a - aj	-CF ₃	cyclobutyl-CH ₂ -
604	A - S	a - aj	H	cyclopentyl
605	C - S	a - aj	F	cyclopentyl
606	C - S	a - aj	Cl	cyclopentyl
607	C - S	a - aj	OH	cyclopentyl
608	C - S	a - aj	-CH ₃	cyclopentyl
609	C - S	a - aj	-CH ₂ CH ₃	cyclopentyl
610	C - S	a - aj	-OCH ₃	cyclopentyl
611	C - S	a - aj	-CF ₃	cyclopentyl
612	A - S	a - aj	H	cyclopentyl-CH ₂ -
613	C - S	a - aj	F	cyclopentyl-CH ₂ -
614	C - S	a - aj	Cl	cyclopentyl-CH ₂ -
615	C - S	a - aj	OH	cyclopentyl-CH ₂ -
616	C - S	a - aj	-CH ₃	cyclopentyl-CH ₂ -
617	C - S	a - aj	-CH ₂ CH ₃	cyclopentyl-CH ₂ -
618	C - S	a - aj	-OCH ₃	cyclopentyl-CH ₂ -
619	C - S	a - aj	-CF ₃	cyclopentyl-CH ₂ -
620	A - S	a - aj	H	cyclohexyl
621	C - S	a - aj	F	cyclohexyl
622	C - S	a - aj	Cl	cyclohexyl

623	C - S	a - aj	OH	cyclohexyl
624	C - S	a - aj	-CH ₃	cyclohexyl
625	C - S	a - aj	-CH ₂ CH ₃	cyclohexyl
626	C - S	a - aj	-OCH ₃	cyclohexyl
627	C - S	a - aj	-CF ₃	cyclohexyl
628	A - S	a - aj	H	cyclohexyl-CH ₂ -
629	C - S	a - aj	F	cyclohexyl-CH ₂ -
630	C - S	a - aj	Cl	cyclohexyl-CH ₂ -
631	C - S	a - aj	OH	cyclohexyl-CH ₂ -
632	C - S	a - aj	-CH ₃	cyclohexyl-CH ₂ -
633	C - S	a - aj	-CH ₂ CH ₃	cyclohexyl-CH ₂ -
634	C - S	a - aj	-OCH ₃	cyclohexyl-CH ₂ -
635	C - S	a - aj	-CF ₃	cyclohexyl-CH ₂ -
636	A - S	a - aj	H	phenyl
637	C - S	a - aj	F	phenyl
638	C - S	a - aj	Cl	phenyl
639	C - S	a - aj	OH	phenyl
640	C - S	a - aj	-CH ₃	phenyl
641	C - S	a - aj	-CH ₂ CH ₃	phenyl
642	C - S	a - aj	-OCH ₃	phenyl
643	C - S	a - aj	-CF ₃	phenyl
644	A - S	a - aj	H	2-F-phenyl
645	C - S	a - aj	F	2-F-phenyl
646	C - S	a - aj	Cl	2-F-phenyl
647	C - S	a - aj	OH	2-F-phenyl
648	C - S	a - aj	-CH ₃	2-F-phenyl
649	C - S	a - aj	-CH ₂ CH ₃	2-F-phenyl
650	C - S	a - aj	-OCH ₃	2-F-phenyl
651	C - S	a - aj	-CF ₃	2-F-phenyl
652	A - S	a - aj	H	3-F-phenyl
653	C - S	a - aj	F	3-F-phenyl
654	C - S	a - aj	Cl	3-F-phenyl
655	C - S	a - aj	OH	3-F-phenyl
656	C - S	a - aj	-CH ₃	3-F-phenyl
657	C - S	a - aj	-CH ₂ CH ₃	3-F-phenyl
658	C - S	a - aj	-OCH ₃	3-F-phenyl

659	C - S	a - aj	-CF ₃	3-F-phenyl
660	A - S	a - aj	H	4-F-phenyl
661	C - S	a - aj	F	4-F-phenyl
662	C - S	a - aj	Cl	4-F-phenyl
663	C - S	a - aj	OH	4-F-phenyl
664	C - S	a - aj	-CH ₃	4-F-phenyl
665	C - S	a - aj	-CH ₂ CH ₃	4-F-phenyl
666	C - S	a - aj	-OCH ₃	4-F-phenyl
667	C - S	a - aj	-CF ₃	4-F-phenyl
668	A - S	a - aj	H	3-Cl-phenyl
669	C - S	a - aj	F	3-Cl-phenyl
670	C - S	a - aj	Cl	3-Cl-phenyl
671	C - S	a - aj	OH	3-Cl-phenyl
672	C - S	a - aj	-CH ₃	3-Cl-phenyl
673	C - S	a - aj	-CH ₂ CH ₃	3-Cl-phenyl
674	C - S	a - aj	-OCH ₃	3-Cl-phenyl
675	C - S	a - aj	-CF ₃	3-Cl-phenyl
676	A - S	a - aj	H	4-Cl-phenyl
677	C - S	a - aj	F	4-Cl-phenyl
678	C - S	a - aj	Cl	4-Cl-phenyl
679	C - S	a - aj	OH	4-Cl-phenyl
680	C - S	a - aj	-CH ₃	4-Cl-phenyl
681	C - S	a - aj	-CH ₂ CH ₃	4-Cl-phenyl
682	C - S	a - aj	-OCH ₃	4-Cl-phenyl
683	C - S	a - aj	-CF ₃	4-Cl-phenyl
684	A - S	a - aj	H	3-Me-phenyl
685	C - S	a - aj	F	3-Me-phenyl
686	C - S	a - aj	Cl	3-Me-phenyl
687	C - S	a - aj	OH	3-Me-phenyl
688	C - S	a - aj	-CH ₃	3-Me-phenyl
689	C - S	a - aj	-CH ₂ CH ₃	3-Me-phenyl
690	C - S	a - aj	-OCH ₃	3-Me-phenyl
691	C - S	a - aj	-CF ₃	3-Me-phenyl
692	A - S	a - aj	H	4-Me-phenyl
693	C - S	a - aj	F	4-Me-phenyl
694	C - S	a - aj	Cl	4-Me-phenyl

695	C - S	a - aj	OH	4-Me-phenyl
696	C - S	a - aj	-CH ₃	4-Me-phenyl
697	C - S	a - aj	-CH ₂ CH ₃	4-Me-phenyl
698	C - S	a - aj	-OCH ₃	4-Me-phenyl
699	C - S	a - aj	-CF ₃	4-Me-phenyl
700	A - S	a - aj	H	3-MeO-phenyl
701	C - S	a - aj	F	3-MeO-phenyl
702	C - S	a - aj	Cl	3-MeO-phenyl
703	C - S	a - aj	OH	3-MeO-phenyl
704	C - S	a - aj	-CH ₃	3-MeO-phenyl
705	C - S	a - aj	-CH ₂ CH ₃	3-MeO-phenyl
706	C - S	a - aj	-OCH ₃	3-MeO-phenyl
707	C - S	a - aj	-CF ₃	3-MeO-phenyl
708	A - S	a - aj	H	4-MeO-phenyl
709	C - S	a - aj	F	4-MeO-phenyl
710	C - S	a - aj	Cl	4-MeO-phenyl
711	C - S	a - aj	OH	4-MeO-phenyl
712	C - S	a - aj	-CH ₃	4-MeO-phenyl
713	C - S	a - aj	-CH ₂ CH ₃	4-MeO-phenyl
714	C - S	a - aj	-OCH ₃	4-MeO-phenyl
715	C - S	a - aj	-CF ₃	4-MeO-phenyl
716	A - S	a - aj	H	3-F ₃ C-phenyl
717	C - S	a - aj	F	3-F ₃ C-phenyl
718	C - S	a - aj	Cl	3-F ₃ C-phenyl
719	C - S	a - aj	OH	3-F ₃ C-phenyl
720	C - S	a - aj	-CH ₃	3-F ₃ C-phenyl
721	C - S	a - aj	-CH ₂ CH ₃	3-F ₃ C-phenyl
722	C - S	a - aj	-OCH ₃	3-F ₃ C-phenyl
723	C - S	a - aj	-CF ₃	3-F ₃ C-phenyl
724	A - S	a - aj	H	4-F ₃ C-phenyl
725	C - S	a - aj	F	4-F ₃ C-phenyl
726	C - S	a - aj	Cl	4-F ₃ C-phenyl
727	C - S	a - aj	OH	4-F ₃ C-phenyl
728	C - S	a - aj	-CH ₃	4-F ₃ C-phenyl
729	C - S	a - aj	-CH ₂ CH ₃	4-F ₃ C-phenyl
730	C - S	a - aj	-OCH ₃	4-F ₃ C-phenyl

731	C - S	a - aj	-CF ₃	4-F ₃ C-phenyl
-----	-------	--------	------------------	---------------------------

UTILITY

A β production has been implicated in the pathology of Alzheimer's Disease (AD). The compounds of the present invention have utility for the prevention and treatment of AD by inhibiting A β production. Methods of treatment target formation of A β production through the enzymes involved in the proteolytic processing of β amyloid precursor protein. Compounds that inhibit β or γ secretase activity, either directly or indirectly, control the production of A β . Such inhibition of β or γ secretases reduces production of A β , and is expected to reduce or prevent the neurological disorders associated with A β protein, such as Alzheimer's Disease.

Cellular screening methods for inhibitors of A β production, testing methods for the *in vivo* suppression of A β production, and assays for the detection of secretase activity are known in the art and have been disclosed in numerous publications, including *J. Med. Chem.* 1999, 42, 3889-3898, PCT publication number WO 98/22493, EPO publication number 0652009, US patent 5703129 and US patent 5593846; all hereby incorporated by reference.

The compounds of the present invention have utility for the prevention and treatment of disorders involving A β production, such as cerebrovascular disorders.

Compounds of Formula (I) are expected to possess γ -secretase inhibitory activity. The γ -secretase inhibitory activity of the compounds of the present invention is demonstrated using assays for such activity, for example, using the assay described below. Compounds of the present invention have been shown to inhibit the activity of γ -secretase, as determined by the A β immunoprecipitation assay.

Compounds provided by this invention should also be useful as standards and reagents in determining the ability of a potential pharmaceutical to inhibit A β production.

These would be provided in commercial kits comprising a compound of this invention.

As used herein "µg" denotes microgram, "mg" denotes milligram, "g" denotes gram, "µL" denotes microliter, "mL" denotes milliliter, "L" denotes liter, "nM" denotes nanomolar, "µM" denotes micromolar, "mM" denotes millimolar, "M" denotes molar, "nm" denotes nanometer, "SDS" denotes sodium dodecyl sulfate, and "DMSO" denotes dimethyl sulfoxide, and "EDTA" denotes ethylenediaminetetraacetate.

A compound is considered to be active if it has an IC₅₀ or K_i value of less than about 100µM for the inhibition of Aβ production. Preferably the IC₅₀ or K_i value is less than about 10µM; more preferably the IC₅₀ or K_i value is less than about 0.1µM. Compounds of the present invention have been shown to inhibit Aβ protein production with an IC₅₀ or K_i value of less than 100µM.

β amyloid precursor protein accumulation assay

A novel assay to evaluate the accumulation of Aβ protein was developed to detect potential inhibitors of secretase. The assay uses the N 9 cell line, characterized for expression of exogenous APP by immunoblotting and immunoprecipitation.

The effect of test compounds on the accumulation of Aβ in the conditioned medium is tested by immunoprecipitation. Briefly, N 9 cells are grown to confluency in 6-well plates and washed twice with 1 x Hank's buffered salt solution. The cells are starved in methionine/cysteine deficient media for 30 min, followed by replacement with fresh deficient media containing 150uCi S35 Translabel (Amersham). Test compounds dissolved in DMSO (final concentration 1%) are added together with the addition of radiolabel. The cells are incubated for 4 h at 37°C in a tissue culture incubator.

At the end of the incubation period, the conditioned medium is harvested and pre-cleared by the addition of 5 µl normal mouse serum and 50ul of protein A Sepharose

(Pharmacia), mixed by end-over-end rotation for 30 minutes at 4°C, followed by a brief centrifugation in a microfuge. The supernatant is then harvested and transferred to fresh tubes containing 5µg of a monoclonal antibody (clone 1101.1; directed against an internal peptide sequence in Aβ) and 50 µl protein A Sepharose. After incubation overnight at 4°C, the samples are washed three times with high salt washing buffer (50mM Tris, pH 7.5, 500mM NaCl, 5mM EDTA, 0.5% Nonidet P-40), three times with low salt wash buffer (50mM Tris, pH 7.5, 150mM NaCl, 5mM EDTA, 0.5% Nonidet P-40), and three times with 10mM Tris, pH 7.5. The pellet after the last wash is resuspended in SDS sample buffer (Laemmli, 1970) and boiled for 3 minutes. The supernatant is then fractionated on either 10-20% Tris/Tricine SDS gels or on 16.5% Tris/Tricine SDS gels. The gels are dried and exposed to X-ray film or analyzed by phosphorimaging. The resulting image is analyzed for the presence of Aβ polypeptides. The steady-state level of Aβ in the presence of a test compound is compared to wells treated with DMSO (1%) alone. A typical test compound blocks Aβ accumulation in the conditioned medium, and is therefore considered active, with an IC₅₀ less than 100 µM.

C-Terminus β Amyloid Precursor Protein Accumulation Assay

The effect of test compounds on the accumulation of C-terminal fragments is determined by immunoprecipitation of APP and fragments thereof from cell lysates. N 9 cells are metabolically labeled as above in the presence or absence of test compounds. At the end of the incubation period, the conditioned medium are harvested and cells lysed in RIPA buffer (10 mM Tris, pH 8.0 containing 1% Triton X-100, 1% deoxycholate, 0.1% SDS, 150mM NaCl, 0.125% NaN₃). Again, lysates are precleared with 5µl normal rabbit serum / 50µl protein A Sepharose, followed by the addition of BC-1 antiserum (15µl;) and 50µl protein A Sepharose for 16 hours at 4°C. The immunoprecipitates are washed as above, bound proteins eluted by boiling in SDS sample buffer and fractionated by Tris/Tricine SDS-PAGE. After exposure to

X-ray film or phosphorimager, the resulting images are analyzed for the presence of C-terminal APP fragments. The steady-state level of C-terminal APP fragments is compared to wells treated with DMSO (1%) alone. A typical test compound stimulates C-terminal fragment accumulation in the cell lysates, and is therefore considered active, with an IC_{50} less than 100 μM .

A β Immunoprecipitation Assay

This immunoprecipitation assay is specific for γ secretase (i.e., proteolytic activity required to generate the C-terminal end of A β either by direct cleavage or generating a C-terminal extended species which is subsequently further proteolyzed). N 9 cells are pulse labeled in the presence of a reported γ secretase inhibitor (MDL 28170) for 1 h, followed by washing to remove radiolabel and MDL 28170. The media is replaced and test compounds are added. The cells are chased for increasing periods of times and A β is isolated from the conditioned medium and C-terminal fragments from cell lysates (see above). The test compounds are characterized whether a stabilization of C-terminal fragments is observed and whether A β is generated from these accumulated precursor. A typical test compound prevents the generation of A β out of accumulated C-terminal fragments and is considered active with an IC_{50} less than 100 μM .

Dosage and Formulation

The compounds of the present invention can be administered orally using any pharmaceutically acceptable dosage form known in the art for such administration. The active ingredient can be supplied in solid dosage forms such as dry powders, granules, tablets or capsules, or in liquid dosage forms, such as syrups or aqueous suspensions. The active ingredient can be administered alone, but is generally administered with a pharmaceutical carrier. A valuable treatise with respect to pharmaceutical dosage

forms is Remington's Pharmaceutical Sciences, Mack Publishing.

The compounds of the present invention can be administered in such oral dosage forms as tablets, capsules (each of which includes sustained release or timed release formulations), pills, powders, granules, elixirs, tinctures, suspensions, syrups, and emulsions. Likewise, they may also be administered in intravenous (bolus or infusion), intraperitoneal, subcutaneous, or intramuscular form, all using dosage forms well known to those of ordinary skill in the pharmaceutical arts. An effective but non-toxic amount of the compound desired can be employed to prevent or treat neurological disorders related to β -amyloid production or accumulation, such as Alzheimer's disease and Down's Syndrome.

The compounds of this invention can be administered by any means that produces contact of the active agent with the agent's site of action in the body of a host, such as a human or a mammal. They can be administered by any conventional means available for use in conjunction with pharmaceuticals, either as individual therapeutic agents or in a combination of therapeutic agents. They can be administered alone, but generally administered with a pharmaceutical carrier selected on the basis of the chosen route of administration and standard pharmaceutical practice.

The dosage regimen for the compounds of the present invention will, of course, vary depending upon known factors, such as the pharmacodynamic characteristics of the particular agent and its mode and route of administration; the species, age, sex, health, medical condition, and weight of the recipient; the nature and extent of the symptoms; the kind of concurrent treatment; the frequency of treatment; the route of administration, the renal and hepatic function of the patient, and the effect desired. An ordinarily skilled physician or veterinarian can readily determine and prescribe the effective amount of the drug

required to prevent, counter, or arrest the progress of the condition.

Advantageously, compounds of the present invention may be administered in a single daily dose, or the total daily
5 dosage may be administered in divided doses of two, three, or four times daily.

The compounds for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal routes, using those
10 forms of transdermal skin patches well known to those of ordinary skill in that art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

15 In the methods of the present invention, the compounds herein described in detail can form the active ingredient, and are typically administered in admixture with suitable pharmaceutical diluents, excipients, or carriers (collectively referred to herein as carrier materials)
20 suitably selected with respect to the intended form of administration, that is, oral tablets, capsules, elixirs, syrups and the like, and consistent with conventional pharmaceutical practices.

For instance, for oral administration in the form of a
25 tablet or capsule, the active drug component can be combined with an oral, non-toxic, pharmaceutically acceptable, inert carrier such as lactose, starch, sucrose, glucose, methyl cellulose, magnesium stearate, dicalcium phosphate, calcium sulfate, mannitol, sorbitol and the
30 like; for oral administration in liquid form, the oral drug components can be combined with any oral, non-toxic, pharmaceutically acceptable inert carrier such as ethanol, glycerol, water, and the like. Moreover, when desired or necessary, suitable binders, lubricants, disintegrating
35 agents, and coloring agents can also be incorporated into the mixture. Suitable binders include starch, gelatin, natural sugars such as glucose or β -lactose, corn sweeteners, natural and synthetic gums such as acacia,

tragacanth, or sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes, and the like. Lubricants used in these dosage forms include sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride, and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum, and the like.

The compounds of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles, and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine, or phosphatidylcholines.

Compounds of the present invention may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamide-phenol, polyhydroxyethylaspartamidephenol, or polyethyleneoxide-polylysine substituted with palmitoyl residues. Furthermore, the compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polyglycolic acid, copolymers of polylactic and polyglycolic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacylates, and crosslinked or amphipathic block copolymers of hydrogels.

Gelatin capsules may contain the active ingredient and powdered carriers, such as lactose, starch, cellulose derivatives, magnesium stearate, stearic acid, and the like. Similar diluents can be used to make compressed tablets. Both tablets and capsules can be manufactured as sustained release products to provide for continuous release of medication over a period of hours. Compressed tablets can be sugar coated or film coated to mask any unpleasant taste and protect the tablet from the atmosphere, or enteric coated for selective disintegration in the gastrointestinal tract.


Liquid dosage forms for oral administration can contain coloring and flavoring to increase patient acceptance. In general, water, a suitable oil, saline, aqueous dextrose (glucose), and related sugar solutions and glycols such as propylene glycol or polyethylene glycols are suitable carriers for parenteral solutions. Solutions for parenteral administration preferably contain a water soluble salt of the active ingredient, suitable stabilizing agents, and if necessary, buffer substances. Antioxidizing agents such as sodium bisulfite, sodium sulfite, or ascorbic acid, either alone or combined, are suitable stabilizing agents. Also used are citric acid and its salts and sodium EDTA. In addition, parenteral solutions can contain preservatives, such as benzalkonium chloride, methyl- or propyl-paraben, and chlorobutanol.

Suitable pharmaceutical carriers are described in Remington's Pharmaceutical Sciences, Mack Publishing Company, a standard reference text in this field.

CLAIMS

What is claimed is:

- 5 1. A compound of Formula (I):



- 10 or a pharmaceutically acceptable salt or prodrug thereof,

wherein:

Q is $-NR^1R^2$;

15

R^1 is H, C_1 - C_4 alkyl, phenyl, benzyl, C_3 - C_6 cycloalkyl, or
(C_3 - C_6 cycloalkyl)methyl-;

20

R^2 is H, C_1 - C_4 alkyl, OH, C_1 - C_4 alkoxy, phenyl, benzyl, C_3 -
 C_6 cycloalkyl, or (C_3 - C_6 cycloalkyl)methyl-;

R^3 and R^{3a} may be combined to form a 3-8 membered
carbocyclic moiety;

25

wherein said 3-8 membered carbocyclic moiety is
saturated or partially unsaturated;

wherein said 3-8 membered carbocyclic moiety may
optionally contain a heteroatom selected from -O-,
-S-, -S(=O)-, -S(=O)₂-, -N=, -NH-, and -N(R^{20})-, and

30

wherein said 3-8 membered carbocyclic moiety is
substituted with 0-4 R^4 ;

additionally, two R^4 substituents on adjacent atoms may be
combined to form a benzo fused radical; wherein said
benzo fused radical is substituted with 0-4 R^{23} ;

35

5 additionally, two R^4 substituents on adjacent atoms may be combined to form a 5 to 6 membered heteroaryl fused radical, wherein said 5 to 6 membered heteroaryl fused radical comprises 1 or 2 heteroatoms selected from N, O, and S; wherein said 5 to 6 membered heteroaryl fused radical is substituted with 0-3 R^{23} ;

10 additionally, two R^4 substituents on the same or adjacent carbon atoms may be combined to form a C_3 - C_6 carbocycle substituted with 0-3 R^{23} ;

15 alternatively, R^3 is H; C_1 - C_6 alkyl substituted with 0-3 R^4 ; C_2 - C_6 alkenyl substituted with 0-3 R^4 ; or C_2 - C_6 alkynyl substituted with 0-3 R^4 ; and R^{3a} is H, C_1 - C_6 alkyl, or C_2 - C_6 alkenyl;

20 R^4 , at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO_2 , CF_3 , acetyl, SCH_3 , $S(=O)CH_3$, $S(=O)_2CH_3$, $NR^{15}R^{16}$, OR^{14a} , C_1 - C_4 alkyl, C_2 - C_6 alkenyl, alkynyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl, C_1 - C_4 haloalkoxy, and C_1 - C_4 haloalkyl-S-, C_3 - C_6 carbocycle, aryl, and a 5 to 6 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur;

25 R^5 is H, C_1 - C_6 alkoxy; C_1 - C_6 alkyl substituted with 0-3 R^{5b} ; C_2 - C_6 alkenyl substituted with 0-3 R^{5b} ; 30 C_2 - C_6 alkynyl substituted with 0-3 R^{5b} ; C_3 - C_{10} carbocycle substituted with 0-3 R^{5c} ; C_6 - C_{10} aryl substituted with 0-3 R^{5c} ; or 5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle 35 is substituted with 0-3 R^{5c} ;

R^{5a} is H, C_1 - C_4 alkyl, or C_2 - C_4 alkenyl;

R^{5b}, at each occurrence, is independently selected from:
H, C₁-C₆ alkyl, CF₃, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂,
NR¹⁵R¹⁶;
5 C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};
C₆-C₁₀ aryl substituted with 0-3 R^{5c}; or
5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered heterocycle
10 is substituted with 0-3 R^{5c};

R^{5c}, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
S(=O)CH₃, S(=O)₂CH₃,
15 C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,
C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

alternatively, R⁵ and R^{5a} may be combined to form a 3-8
membered carbocyclic moiety;
20 wherein said 3-8 membered carbocyclic moiety is
saturated or partially unsaturated;
wherein said 3-8 membered carbocyclic moiety may
optionally contain a heteroatom selected from -O-,
-S-, -S(=O)-, -S(=O)₂-, -N=, -NH-, and -N(R²⁰)-, and
25 wherein said 3-8 membered carbocyclic moiety is
substituted with 0-4 R^{5c};

provided at least:

- 1) R³ and R^{3a} are combined to form a 3-8 membered
30 carbocyclic moiety; or
- 2) R⁵ and R^{5a} are combined to form a 3-8 membered
carbocyclic moiety; or
- 3) R³ and R^{3a} are combined to form a 3-8 membered
carbocyclic moiety and R⁵ and R^{5a} are combined to form
35 a 3-8 membered carbocyclic moiety;

R⁶ is H;

C₁-C₆ alkyl substituted with 0-3 R^{6a};

C₃-C₁₀ carbocycle substituted with 0-3 R^{6b}; or
C₆-C₁₀ aryl substituted with 0-3 R^{6b};

5 R^{6a}, at each occurrence, is independently selected from H,
C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶,
aryl or CF₃;

10 R^{6b}, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, C₁-C₆ alkyl,
C₁-C₄ alkoxy, C₁-C₄ haloalkyl, and C₁-C₄ haloalkoxy;

Ring B is a 6, 7, or 8 membered lactam,
wherein the lactam is saturated, partially saturated
or unsaturated;
15 wherein each additional lactam carbon is substituted
with 0-2 R¹¹; and,
optionally, the lactam contains an additional
heteroatom selected from -O-, -S-, -S(=O)-, -S(=O)₂-
, -N=, -NH-, and -N(R¹⁰)-;

20 additionally, two R¹¹ substituents on adjacent atoms may be
combined to form a benzo fused radical; wherein said
benzo fused radical is substituted with 0-4 R¹³;

25 additionally, two R¹¹ substituents on adjacent atoms may be
combined to form a 5 to 6 membered heteroaryl fused
radical, wherein said 5 to 6 membered heteroaryl fused
radical comprises 1 or 2 heteroatoms selected from N,
O, and S; wherein said 5 to 6 membered heteroaryl
30 fused radical is substituted with 0-3 R¹³;

additionally, two R¹¹ substituents on the same or adjacent
carbon atoms may be combined to form a C₃-C₆
carbocycle substituted with 0-3 R¹³;

35 R¹⁰ is H, C(=O)R¹⁷, C(=O)OR¹⁷, C(=O)NR¹⁸R¹⁹,
S(=O)₂NR¹⁸R¹⁹, S(=O)₂R¹⁷;
C₁-C₆ alkyl optionally substituted with 0-3 R^{10a};

- C₆-C₁₀ aryl substituted with 0-4 R^{10b};
C₃-C₁₀ carbocycle substituted with 0-3 R^{10b}; or
5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered heterocycle
is substituted with 0-3 R^{10b};
- R^{10a}, at each occurrence, is independently selected from H,
C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶,
CF₃, aryl substituted with 0-4 R^{10b}; or a 5-6 membered
heterocycle substituted with 0-4 R^{10b};
- R^{10b}, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄
haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;
- R¹¹, at each occurrence, is independently selected from
H, C₁-C₄ alkoxy, Cl, F, Br, I, =O, CN, NO₂, NR¹⁸R¹⁹,
C(=O)R¹⁷, C(=O)OR¹⁷, C(=O)NR¹⁸R¹⁹, S(=O)₂NR¹⁸R¹⁹, CF₃;
C₁-C₆ alkyl optionally substituted with 0-3 R^{11a};
C₆-C₁₀ aryl substituted with 0-3 R^{11b};
C₃-C₁₀ carbocycle substituted with 0-3 R^{11b}; or
5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered heterocycle
is substituted with 0-3 R^{11b};
- R^{11a}, at each occurrence, is independently selected from
H, C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂,
NR¹⁵R¹⁶, CF₃;
phenyl substituted with 0-3 R^{11b};
C₃-C₆ cycloalkyl substituted with 0-3 R^{11b}; and
5 to 6 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 6 membered heterocycle
is substituted with 0-3 R^{11b};

R^{11b} , at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

W is a bond or $-(CR^8R^{8a})_p-$;

p is 0, 1, 2, 3, or 4;

R^8 and R^{8a} , at each occurrence, are independently selected from H, F, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl and C₃-C₈ cycloalkyl;

X is a bond;

C₆-C₁₀ aryl substituted with 0-3 R^{Xb} ;

C₃-C₁₀ carbocycle substituted with 0-3 R^{Xb} ; or

5 to 10 membered heterocycle substituted with 0-2 R^{Xb} ;

R^{Xb} , at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ halothioalkoxy;

Y is a bond or $-(CR^9R^{9a})_t-V-(CR^9R^{9a})_u-$;

t is 0, 1, or 2;

u is 0, 1, or 2;

R^9 and R^{9a} , at each occurrence, are independently selected from H, F, C₁-C₆ alkyl or C₃-C₈ cycloalkyl;

V is a bond, -C(=O)-, -O-, -S-, -S(=O)-, -S(=O)₂-, -N(R¹⁹)-, -C(=O)NR^{19b}-, -NR^{19b}C(=O)-, -NR^{19b}S(=O)₂-, -S(=O)₂NR^{19b}-, -NR^{19b}S(=O)-, -S(=O)NR^{19b}-, -C(=O)O-, or -OC(=O)-;

Z is H;

C₁-C₈ alkyl substituted with 0-3 R^{12a};
C₂-C₆ alkenyl substituted with 0-3 R^{12a};
C₂-C₆ alkynyl substituted with 0-3 R^{12a};
5 C₆-C₁₀ aryl substituted with 0-4 R^{12b};
C₃-C₁₀ carbocycle substituted with 0-4 R^{12b}; or
5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered heterocycle
10 is substituted with 0-3 R^{12b};

R^{12a}, at each occurrence, is independently selected from
H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, -C(=O)NR¹⁵R¹⁶,
CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃,
15 C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,
C₁-C₄ haloalkoxy, C₁-C₄ haloalkyl-S-,
C₆-C₁₀ aryl substituted with 0-4 R^{12b};
C₃-C₁₀ carbocycle substituted with 0-4 R^{12b}; or
5 to 10 membered heterocycle containing 1 to 4
20 heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered heterocycle
is substituted with 0-3 R^{12b};

R^{12b}, at each occurrence, is independently selected from
25 H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl,
SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₃-C₆ cycloalkyl,
C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,
C₁-C₄ haloalkoxy, C₁-C₄ haloalkyl-S, and
aryl substituted with 0-3 R^{12c};

30 R^{12c}, at each occurrence, is independently selected from
H, methyl, ethyl, propyl, methoxy, ethoxy, amino,
hydroxy, Cl, F, Br, I, CF₃, SCH₃, S(O)CH₃, SO₂CH₃,
-N(CH₃)₂, N(CH₃)H, CN, NO₂, OCF₃, C(=O)CH₃, CO₂H,
35 CO₂CH₃, and C₁-C₃ haloalkyl;

R¹³, at each occurrence, is independently selected from

H, OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, and CF₃;

5 R¹⁴ is H, phenyl, benzyl, C₁-C₆ alkyl, C₂-C₆ alkoxyalkyl, or C₃-C₆ cycloalkyl;

R^{14a} is H, phenyl, benzyl, or C₁-C₄ alkyl;

10 R¹⁵, at each occurrence, is independently selected from H, C₁-C₆ alkyl, benzyl, phenethyl, (C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;

15 R¹⁶, at each occurrence, is independently selected from H, C₁-C₆ alkyl, benzyl, phenethyl, (C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;

20 R¹⁷ is H, C₁-C₆ alkyl, C₂-C₆ alkoxyalkyl, aryl substituted by 0-4 R^{17a}, or -CH₂-aryl substituted by 0-4 R^{17a};

R^{17a} is H, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, butoxy, -OH, F, Cl, Br, I, CF₃, OCF₃, SCH₃, S(O)CH₃, SO₂CH₃, -NH₂, -N(CH₃)₂, or C₁-C₄ haloalkyl;

25 R¹⁸, at each occurrence, is independently selected from H, C₁-C₆ alkyl, phenyl, benzyl, phenethyl, (C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;

30 R¹⁹, at each occurrence, is independently selected from H, OH, C₁-C₆ alkyl, phenyl, benzyl, phenethyl, (C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;

35 additionally, R¹⁸ and R¹⁹, when substituents on the same atom, may be combined to form a 3 to 6 membered heterocyclic ring;

R^{19b}, at each occurrence, is independently is H or C₁-C₄ alkyl;

R²⁰ is H, C(=O)R¹⁷, C(=O)OR¹⁷, C(=O)NR¹⁸R¹⁹,
S(=O)₂NR¹⁸R¹⁹, S(=O)₂R¹⁷;

C₁-C₆ alkyl optionally substituted with 0-3 R^{20a}; or
5 C₆-C₁₀ aryl substituted with 0-4 R^{20b};

R^{20a}, at each occurrence, is independently selected from H,
C₁-C₄ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶,
CF₃, or aryl substituted with 0-4 R^{20b};


10

R^{20b}, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
S(=O)CH₃, S(=O)₂CH₃, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄
haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

15

R²³, at each occurrence, is independently selected from
H, OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN,
NO₂, NR¹⁵R¹⁶, and CF₃.

20 2. A compound of claim 1 of Formula (II):



25 or a pharmaceutically acceptable salt or prodrug thereof,
wherein:

R³ and R^{3a} may be combined to form a 3-8 membered
carbocyclic moiety;

30 wherein said 3-8 membered carbocyclic moiety is
saturated or partially unsaturated; and

R⁵ and R^{5a} may be combined to form a 3-8 membered
carbocyclic moiety;

wherein said 3-8 membered carbocyclic moiety is saturated or partially unsaturated;

provided at least:

- 5 1) R^3 and R^{3a} are combined to form a 3-8 membered carbocyclic moiety; or
2) R^5 and R^{5a} are combined to form a 3-8 membered carbocyclic moiety; or
3) R^3 and R^{3a} are combined to form a 3-8 membered carbocyclic moiety and R^5 and R^{5a} are combined to form
10 a 3-8 membered carbocyclic moiety.

3. A compound of claim 2 of Formula (II), wherein:

- 15 R^3 and R^{3a} may be combined to form a 3-8 membered carbocyclic moiety selected from cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, and cyclohexenyl;

- 20 alternatively, R^3 and R^{3a} are independently selected from the group H, C_1-C_6 alkyl, C_2-C_6 alkenyl, and C_2-C_6 alkynyl;

R^5 is H;

- 25 C_1-C_4 alkyl substituted with 0-1 R^{5b} ;
 C_2-C_4 alkenyl substituted with 0-1 R^{5b} ;
 C_2-C_4 alkynyl substituted with 0-1 R^{5b} ;

R^{5a} is H, C_1-C_4 alkyl, or C_2-C_4 alkenyl;

30

R^{5b} is selected from:

H, methyl, ethyl, propyl, butyl, CF_3 , Cl, F, $NR^{15}R^{16}$, cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl;

- 35 alternatively, R^5 and R^{5a} may be combined to form a 3-8 membered carbocyclic moiety selected from cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, and cyclohexenyl;

provided at least:

- 1) R^3 and R^{3a} are combined to form a 3-8 membered carbocyclic moiety; or
- 5 2) R^5 and R^{5a} are combined to form a 3-8 membered carbocyclic moiety; or
- 3) R^3 and R^{3a} are combined to form a 3-8 membered carbocyclic moiety and R^5 and R^{5a} are combined to form a 3-8 membered carbocyclic moiety;

10

Ring B is a 7 membered lactam,
wherein the lactam is saturated, partially saturated
or unsaturated;
wherein each additional lactam carbon is substituted
15 with 0-2 R^{11} ; and,
optionally, the lactam contains an additional
heteroatom selected from -O-, -S-, -S(=O)-, -S(=O)₂-
, -N=, -NH-, and -N(R^{10})-;


- 20 additionally, two R^{11} substituents on adjacent atoms may be
combined to form a benzo fused radical; wherein said
benzo fused radical is substituted with 0-4 R^{13} ;

25 additionally, two R^{11} substituents on adjacent atoms may be
combined to form a 5 to 6 membered heteroaryl fused
radical, wherein said 5 to 6 membered heteroaryl fused
radical comprises 1 or 2 heteroatoms selected from N,
O, and S; wherein said 5 to 6 membered heteroaryl
fused radical is substituted with 0-3 R^{13} ;

30

additionally, two R^{11} substituents on the same or adjacent
carbon atoms may be combined to form a C₃-C₆
carbocycle substituted with 0-3 R^{13} .

- 35 4. A compound of Claim 1 of Formula (I):



(I)

or a pharmaceutically acceptable salt or prodrug thereof,
 5 wherein:

Q is $-NR^1R^2$;

10 R^1 is H, methyl, ethyl, propyl, butyl, phenyl, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl, or cyclohexylmethyl;

15 R^2 is H, methyl, ethyl, propyl, butyl, OH, methoxy, ethoxy, propoxy, phenyl, benzyl, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclopropylmethyl, cyclobutylmethyl, cyclopentylmethyl, or cyclohexylmethyl;

20 R^3 and R^{3a} are combined to form a 3-8 membered carbocyclic moiety;
 wherein said 3-8 membered carbocyclic moiety is saturated or partially unsaturated;
 wherein said 3-8 membered carbocyclic moiety may
 25 optionally contain a heteroatom selected from -O-, -S-, -S(=O)-, -S(=O)₂-, -N=, -NH-, and -N(R^{20})-, and
 wherein said 3-8 membered carbocyclic moiety is substituted with 0-4 R^4 ;

30 additionally, two R^4 substituents on adjacent atoms may be combined to form a benzo fused radical; wherein said benzo fused radical is substituted with 0-4 R^{23} ;

35 additionally, two R^4 substituents on adjacent atoms may be combined to form a 5 to 6 membered heteroaryl fused

radical, wherein said 5 to 6 membered heteroaryl fused radical comprises 1 or 2 heteroatoms selected from N, O, and S; wherein said 5 to 6 membered heteroaryl fused radical is substituted with 0-3 R²³;

5

additionally, two R⁴ substituents on the same or adjacent carbon atoms may be combined to form a C₃-C₆ carbocycle substituted with 0-3 R²³;

10 R⁴, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, NR¹⁵R¹⁶, OR^{14a}, C₁-C₄ alkyl, C₂-C₆ alkenyl, alkynyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-,
15 C₃-C₆ carbocycle, aryl, and a 5 to 6 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur;

20 R⁵ is H, C₁-C₆ alkoxy;
C₁-C₆ alkyl substituted with 0-3 R^{5b};
C₂-C₆ alkenyl substituted with 0-3 R^{5b};
C₂-C₆ alkynyl substituted with 0-3 R^{5b};
C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};
25 C₆-C₁₀ aryl substituted with 0-3 R^{5c}; or
5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{5c};

30

R^{5a} is H, C₁-C₄ alkyl, or C₂-C₄ alkenyl;

R^{5b}, at each occurrence, is independently selected from:
H, C₁-C₆ alkyl, CF₃, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂,

35

NR¹⁵R¹⁶;
C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};
C₆-C₁₀ aryl substituted with 0-3 R^{5c}; or

5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered heterocycle
is substituted with 0-3 R^{5c};

5

R^{5c}, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
S(=O)CH₃, S(=O)₂CH₃,

10

C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,
C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

R⁶ is H;

C₁-C₆ alkyl substituted with 0-3 R^{6a};

C₃-C₁₀ carbocycle substituted with 0-3 R^{6b}; or

15

C₆-C₁₀ aryl substituted with 0-3 R^{6b};

R^{6a}, at each occurrence, is independently selected from H,
C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶,
aryl or CF₃;

20

R^{6b}, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, C₁-C₆ alkyl,
C₁-C₄ alkoxy, C₁-C₄ haloalkyl, and C₁-C₄ haloalkoxy;

25

Ring B is a 7 membered lactam,

wherein the lactam is saturated, partially saturated
or unsaturated;

wherein each additional lactam carbon is substituted
with 0-2 R¹¹; and,

30

optionally, the lactam contains an additional

heteroatom selected from -O-, -S-, -S(=O)-, -S(=O)₂-
, -N=, -NH-, and -N(R¹⁰)-;

additionally, two R¹¹ substituents on adjacent atoms may be
combined to form a benzo fused radical; wherein said
benzo fused radical is substituted with 0-4 R¹³;

35

5 additionally, two R¹¹ substituents on adjacent atoms may be combined to form a 5 to 6 membered heteroaryl fused radical, wherein said 5 to 6 membered heteroaryl fused radical comprises 1 or 2 heteroatoms selected from N, O, and S; wherein said 5 to 6 membered heteroaryl fused radical is substituted with 0-3 R¹³;

10 additionally, two R¹¹ substituents on the same or adjacent carbon atoms may be combined to form a C₃-C₆ carbocycle substituted with 0-3 R¹³;

R¹⁰ is H, C(=O)R¹⁷, C(=O)OR¹⁷, C(=O)NR¹⁸R¹⁹, S(=O)₂NR¹⁸R¹⁹, S(=O)₂R¹⁷;
C₁-C₆ alkyl optionally substituted with 0-3 R^{10a};
15 C₆-C₁₀ aryl substituted with 0-4 R^{10b};
C₃-C₁₀ carbocycle substituted with 0-3 R^{10b}; or
5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle
20 is substituted with 0-3 R^{10b};

R^{10a}, at each occurrence, is independently selected from H, C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶, CF₃, aryl substituted with 0-4 R^{10b}; or a 5-6 membered
25 heterocycle substituted with 0-4 R^{10b};

R^{10b}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄
30 haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

R¹¹, at each occurrence, is independently selected from H, C₁-C₄ alkoxy, Cl, F, Br, I, =O, CN, NO₂, NR¹⁸R¹⁹, C(=O)R¹⁷, C(=O)OR¹⁷, C(=O)NR¹⁸R¹⁹, S(=O)₂NR¹⁸R¹⁹, CF₃;
35 C₁-C₆ alkyl optionally substituted with 0-3 R^{11a};
C₆-C₁₀ aryl substituted with 0-3 R^{11b};
C₃-C₁₀ carbocycle substituted with 0-3 R^{11b}; or

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{11b};

5

R^{11a}, at each occurrence, is independently selected from H, C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶, CF₃;

phenyl substituted with 0-3 R^{11b};

10

C₃-C₆ cycloalkyl substituted with 0-3 R^{11b}; and

5 to 6 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R^{11b};

15

R^{11b}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃,

C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,

20

C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

W is a bond or -(CR⁸R^{8a})_p-;

p is 0, 1, 2, 3, or 4;

25

R⁸ and R^{8a}, at each occurrence, are independently selected from H, F, C₁-C₄ alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl and C₃-C₈ cycloalkyl;

30 X is a bond;

C₆-C₁₀ aryl substituted with 0-3 R^{Xb};

C₃-C₁₀ carbocycle substituted with 0-3 R^{Xb}; or

5 to 10 membered heterocycle substituted with 0-2 R^{Xb};

35

R^{Xb}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ halothioalkoxy;

Y is a bond or $-(CR^9R^{9a})_t-V-(CR^9R^{9a})_u-$;

t is 0, 1, or 2;

5

u is 0, 1, or 2;

R^9 and R^{9a} , at each occurrence, are independently selected from H, F, C_1-C_6 alkyl or C_3-C_8 cycloalkyl;

10

V is a bond, $-C(=O)-$, $-O-$, $-S-$, $-S(=O)-$, $-S(=O)_2-$, $-N(R^{19})-$, $-C(=O)NR^{19b}-$, $-NR^{19b}C(=O)-$, $-NR^{19b}S(=O)_2-$, $-S(=O)_2NR^{19b}-$, $-NR^{19b}S(=O)-$, $-S(=O)NR^{19b}-$, $-C(=O)O-$, or $-OC(=O)-$;

15

Z is H;

C_1-C_8 alkyl substituted with 0-3 R^{12a} ;

C_2-C_6 alkenyl substituted with 0-3 R^{12a} ;

C_2-C_6 alkynyl substituted with 0-3 R^{12a} ;

20

C_6-C_{10} aryl substituted with 0-4 R^{12b} ;

C_3-C_{10} carbocycle substituted with 0-4 R^{12b} ; or

5 to 10 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 10 membered heterocycle

25

is substituted with 0-3 R^{12b} ;

R^{12a} , at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO_2 , $NR^{15}R^{16}$, $-C(=O)NR^{15}R^{16}$, CF_3 , acetyl, SCH_3 , $S(=O)CH_3$, $S(=O)_2CH_3$,

30

C_1-C_6 alkyl, C_1-C_4 alkoxy, C_1-C_4 haloalkyl,

C_1-C_4 haloalkoxy, C_1-C_4 haloalkyl-S-,

C_6-C_{10} aryl substituted with 0-4 R^{12b} ;

C_3-C_{10} carbocycle substituted with 0-4 R^{12b} ; or

5 to 10 membered heterocycle containing 1 to 4

35

heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 10 membered heterocycle

is substituted with 0-3 R^{12b} ;

R^{12b}, at each occurrence, is independently selected from
H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl,
SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₃-C₆ cycloalkyl,
C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,
5 C₁-C₄ haloalkoxy, C₁-C₄ haloalkyl-S, and
aryl substituted with 0-3 R^{12c};

R^{12c}, at each occurrence, is independently selected from
H, methyl, ethyl, propyl, methoxy, ethoxy, amino,
10 hydroxy, Cl, F, Br, I, CF₃, SCH₃, S(O)CH₃, SO₂CH₃,
-N(CH₃)₂, N(CH₃)H, CN, NO₂, OCF₃, C(=O)CH₃, CO₂H,
CO₂CH₃, and C₁-C₃ haloalkyl;

R¹³, at each occurrence, is independently selected from
15 H, OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN,
NO₂, NR¹⁵R¹⁶, and CF₃;

R¹⁴ is H, phenyl, benzyl, C₁-C₆ alkyl, C₂-C₆ alkoxyalkyl, or
C₃-C₆ cycloalkyl;

20 R^{14a} is H, phenyl, benzyl, or C₁-C₄ alkyl;

R¹⁵, at each occurrence, is independently selected from H,
C₁-C₆ alkyl, benzyl, phenethyl, (C₁-C₆ alkyl)-C(=O)-,
25 and (C₁-C₆ alkyl)-S(=O)₂-;

R¹⁶, at each occurrence, is independently selected from
H, C₁-C₆ alkyl, benzyl, phenethyl,
(C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;

30 R¹⁷ is H, C₁-C₆ alkyl, C₂-C₆ alkoxyalkyl,
aryl substituted by 0-4 R^{17a}, or
-CH₂-aryl substituted by 0-4 R^{17a};

35 R^{17a} is H, methyl, ethyl, propyl, butyl, methoxy, ethoxy,
propoxy, butoxy, -OH, F, Cl, Br, I, CF₃, OCF₃, SCH₃,
S(O)CH₃, SO₂CH₃, -NH₂, -N(CH₃)₂, or C₁-C₄ haloalkyl;

R¹⁸, at each occurrence, is independently selected from
H, C₁-C₆ alkyl, phenyl, benzyl, phenethyl,
(C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;

5 R¹⁹, at each occurrence, is independently selected from
H, OH, C₁-C₆ alkyl, phenyl, benzyl, phenethyl,
(C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;

10 additionally, R¹⁸ and R¹⁹, when substituents on the same
atom, may be combined to form a 3 to 6 membered
heterocyclic ring;

R^{19b}, at each occurrence, is independently is H or C₁-C₄
alkyl;

15 R²⁰ is H, C(=O)R¹⁷, C(=O)OR¹⁷, C(=O)NR¹⁸R¹⁹,
S(=O)₂NR¹⁸R¹⁹, S(=O)₂R¹⁷;
C₁-C₆ alkyl optionally substituted with 0-3 R^{20a}; or
C₆-C₁₀ aryl substituted with 0-4 R^{20b};


20 R^{20a}, at each occurrence, is independently selected from H,
C₁-C₄ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶,
CF₃, or aryl substituted with 0-4 R^{20b};

25 R^{20b}, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
S(=O)CH₃, S(=O)₂CH₃, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₁-C₄
haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

30 R²³, at each occurrence, is independently selected from
H, OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN,
NO₂, NR¹⁵R¹⁶, and CF₃.

5. A compound, according to Claim 4, of Formula (Ia):

35



or a pharmaceutically acceptable salt or prodrug thereof,
 5 wherein:

R^3 and R^{3a} are combined to form a 3-8 membered carbocyclic moiety;

10 wherein said 3-8 membered carbocyclic moiety is saturated or partially unsaturated;

wherein said 3-8 membered carbocyclic moiety is substituted with 0-4 R^4 ;

15 additionally, two R^4 substituents on adjacent atoms may be combined to form a benzo fused radical; wherein said benzo fused radical is substituted with 0-4 R^{23} ;

20 additionally, two R^4 substituents on adjacent atoms may be combined to form a 5 to 6 membered heteroaryl fused radical, wherein said 5 to 6 membered heteroaryl fused radical comprises 1 or 2 heteroatoms selected from N, O, and S; wherein said 5 to 6 membered heteroaryl fused radical is substituted with 0-3 R^{23} ;

25 additionally, two R^4 substituents on the same or adjacent carbon atoms may be combined to form a C_3 - C_6 carbocycle substituted with 0-3 R^{23} ;

30 R^4 , at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO_2 , CF_3 , acetyl, SCH_3 , $\text{S}(=\text{O})\text{CH}_3$, $\text{S}(=\text{O})_2\text{CH}_3$, $\text{NR}^{15}\text{R}^{16}$, OR^{14a} , C_1 - C_4 alkyl, C_2 - C_6 alkenyl, alkynyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl, C_1 - C_4 haloalkoxy, and C_1 - C_4 haloalkyl-S-, C_3 - C_6 carbocycle, aryl, and a

5 to 6 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur; and

5 R⁵ is H, C₁-C₄ alkoxy;

C₁-C₆ alkyl substituted with 0-3 R^{5b};

C₂-C₆ alkenyl substituted with 0-3 R^{5b};

C₂-C₆ alkynyl substituted with 0-3 R^{5b};

C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};

10 C₆-C₁₀ aryl substituted with 0-3 R^{5c}; or

5 to 10 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 10 membered heterocycle

is substituted with 0-3 R^{5c};

15

R^{5b}, at each occurrence, is independently selected from:

H, C₁-C₆ alkyl, CF₃, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂,

NR¹⁵R¹⁶;

C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};

20

C₆-C₁₀ aryl substituted with 0-3 R^{5c}; or

5 to 10 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 10 membered heterocycle

is substituted with 0-3 R^{5c};

25

R^{5c}, at each occurrence, is independently selected from H,

OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,

S(=O)CH₃, S(=O)₂CH₃,

C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,


30

C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

R⁶ is H, methyl, or ethyl;

Ring B is selected from:

35



R^{10} is H, $C(=O)R^{17}$, $C(=O)OR^{17}$, $C(=O)NR^{18}R^{19}$,
 $S(=O)_2NR^{18}R^{19}$, $S(=O)_2R^{17}$;

C_1 - C_6 alkyl optionally substituted with 0-3 R^{10a} ;

10 C_6 - C_{10} aryl substituted with 0-4 R^{10b} ;

C_3 - C_{10} carbocycle substituted with 0-3 R^{10b} ; or

5 to 10 membered heterocycle containing 1 to 4
 heteroatoms selected from nitrogen, oxygen, and
 sulphur, wherein said 5 to 10 membered heterocycle
 15 is substituted with 0-3 R^{10b} ;

R^{10a} , at each occurrence, is independently selected from H,
 C_1 - C_6 alkyl, OR^{14} , Cl, F, Br, I, =O, CN, NO_2 , $NR^{15}R^{16}$,
 CF_3 , aryl substituted with 0-4 R^{10b} ; or a 5-6 membered
 20 heterocycle substituted with 0-4 R^{10b} ;

R^{10b} , at each occurrence, is independently selected from H,
 OH, Cl, F, Br, I, CN, NO_2 , $NR^{15}R^{16}$, CF_3 , acetyl, SCH_3 ,
 $S(=O)CH_3$, $S(=O)_2CH_3$, C_1 - C_6 alkyl, C_1 - C_4 alkoxy, C_1 - C_4
 25 haloalkyl, C_1 - C_4 haloalkoxy, and C_1 - C_4 haloalkyl-S-;

R¹¹, at each occurrence, is independently selected from
H, C₁-C₄ alkoxy, Cl, F, Br, I, =O, CN, NO₂, NR¹⁸R¹⁹,
C(=O)R¹⁷, C(=O)OR¹⁷, C(=O)NR¹⁸R¹⁹, S(=O)₂NR¹⁸R¹⁹, CF₃;
5 C₁-C₆ alkyl optionally substituted with 0-3 R^{11a};
C₆-C₁₀ aryl substituted with 0-3 R^{11b};
C₃-C₁₀ carbocycle substituted with 0-3 R^{11b}; or
5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
10 sulphur, wherein said 5 to 10 membered heterocycle
is substituted with 0-3 R^{11b};

R^{11a}, at each occurrence, is independently selected from
H, C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂,
15 NR¹⁵R¹⁶, CF₃;
phenyl substituted with 0-3 R^{11b};
C₃-C₆ cycloalkyl substituted with 0-3 R^{11b}; and
5 to 6 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
20 sulphur, wherein said 5 to 6 membered heterocycle
is substituted with 0-3 R^{11b};

R^{11b}, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
25 S(=O)CH₃, S(=O)₂CH₃,
C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,
C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

W is a bond or -(CH₂)_p-;

30

p is 1 or 2;

X is a bond;

phenyl substituted with 0-2 R^{Xb};
35 C₃-C₆ carbocycle substituted with 0-2 R^{Xb}; or
5 to 6 membered heterocycle substituted with 0-2 R^{Xb};

R^{Xb} , at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₄ alkyl, C₁-C₃ alkoxy, C₁-C₃ haloalkyl, C₁-C₃ haloalkoxy, and C₁-C₃ halothioalkoxy;

5

Y is a bond, -C(=O)-, -O-, -S-, -S(=O)-, -S(=O)₂-, -N(R¹⁹)-, -C(=O)NR^{19b}-, -NR^{19b}C(=O)-, -NR^{19b}S(=O)₂-, -S(=O)₂NR^{19b}-, -NR^{19b}S(=O)-, -S(=O)NR^{19b}-, -C(=O)O-, or -OC(=O)-;

10

Z is H;

C₁-C₈ alkyl substituted with 0-3 R^{12a};

C₂-C₆ alkenyl substituted with 0-3 R^{12a};

C₂-C₆ alkynyl substituted with 0-3 R^{12a};

15

C₆-C₁₀ aryl substituted with 0-4 R^{12b};

C₃-C₁₀ carbocycle substituted with 0-4 R^{12b}; or

5 to 10 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 10 membered heterocycle

20

is substituted with 0-3 R^{12b};

R^{12a}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, -C(=O)NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃,

25

C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,

C₁-C₄ haloalkoxy, C₁-C₄ haloalkyl-S-,

C₆-C₁₀ aryl substituted with 0-4 R^{12b};

C₃-C₁₀ carbocycle substituted with 0-4 R^{12b}; or

5 to 10 membered heterocycle containing 1 to 4

30

heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 10 membered heterocycle

is substituted with 0-3 R^{12b};

R^{12b}, at each occurrence, is independently selected from

35

H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl,

SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₃-C₆ cycloalkyl,

C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,

C₁-C₄ haloalkoxy, C₁-C₄ haloalkyl-S, and

aryl substituted with 0-3 R^{12c};

5 R^{12c}, at each occurrence, is independently selected from
H, methyl, ethyl, propyl, methoxy, ethoxy, amino,
hydroxy, Cl, F, Br, I, CF₃, SCH₃, S(O)CH₃, SO₂CH₃,
-N(CH₃)₂, N(CH₃)H, CN, NO₂, OCF₃, C(=O)CH₃, CO₂H,
CO₂CH₃, and C₁-C₃ haloalkyl;

10 R¹³, at each occurrence, is independently selected from
H, OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN,
NO₂, NR¹⁵R¹⁶, and CF₃;

R¹⁴ is H, phenyl, benzyl, C₁-C₆ alkyl, C₂-C₆ alkoxyalkyl, or
C₃-C₆ cycloalkyl;

15 R^{14a} is H, phenyl, benzyl, or C₁-C₄ alkyl;

R¹⁵, at each occurrence, is independently selected from H,
C₁-C₆ alkyl, benzyl, phenethyl, (C₁-C₆ alkyl)-C(=O)-,
20 and (C₁-C₆ alkyl)-S(=O)₂-;

R¹⁶, at each occurrence, is independently selected from
H, C₁-C₆ alkyl, benzyl, phenethyl,
(C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;

25 R¹⁷ is H, C₁-C₆ alkyl, C₂-C₆ alkoxyalkyl,
aryl substituted by 0-4 R^{17a}, or
-CH₂-aryl substituted by 0-4 R^{17a};

30 R^{17a} is H, methyl, ethyl, propyl, butyl, methoxy, ethoxy,
propoxy, butoxy, -OH, F, Cl, Br, I, CF₃, OCF₃, SCH₃,
S(O)CH₃, SO₂CH₃, -NH₂, -N(CH₃)₂, or C₁-C₄ haloalkyl;

35 R¹⁸, at each occurrence, is independently selected from
H, C₁-C₆ alkyl, phenyl, benzyl, phenethyl,
(C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;

R¹⁹, at each occurrence, is independently selected from


H, OH, methyl, ethyl, propyl, butyl, phenyl, benzyl, phenethyl;

5 additionally, R^{18} and R^{19} , when substituents on the same atom, may be combined to form a 3 to 6 membered heterocyclic ring selected from pyrrolyl, imidazolyl, imidazolidinyl, pyrrolidinyl, piperidinyl, piperazinyl, and morpholinyl;

10 R^{19b} , at each occurrence, is independently is H or C_1-C_4 alkyl; and

R^{23} , at each occurrence, is independently selected from H, OH, C_1-C_6 alkyl, C_1-C_4 alkoxy, Cl, F, Br, I, CN, (15 NO_2 , $NR^{15}R^{16}$, and CF_3 .

6. A compound, according to Claim 5, of Formula (Ib) :



or a pharmaceutically acceptable salt or prodrug thereof, wherein:

25 R^3 and R^{3a} are combined to form a 3-8 membered carbocyclic moiety;

wherein said 3-8 membered carbocyclic moiety is saturated or partially unsaturated;

30 wherein said 3-8 membered carbocyclic moiety is substituted with 0-4 R^4 ;

additionally, two R^4 substituents on adjacent atoms may be combined to form a benzo fused radical; wherein said benzo fused radical is substituted with 0-4 R^{23} ;

35

additionally, two R^4 substituents on adjacent atoms may be combined to form a 5 to 6 membered heteroaryl fused radical, wherein said 5 to 6 membered heteroaryl fused radical comprises 1 or 2 heteroatoms selected from N, O, and S; wherein said 5 to 6 membered heteroaryl fused radical is substituted with 0-3 R^{23} ;

additionally, two R^4 substituents on the same or adjacent carbon atoms may be combined to form a C_3 - C_6 carbocycle substituted with 0-3 R^{23} ;

R^4 , at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO_2 , CF_3 , acetyl, SCH_3 , $S(=O)CH_3$, $S(=O)_2CH_3$, $NR^{15}R^{16}$, OR^{14a} , C_1 - C_4 alkyl, C_2 - C_6 alkenyl, alkynyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl, C_1 - C_4 haloalkoxy, and C_1 - C_4 haloalkyl-S-, C_3 - C_6 carbocycle, aryl, and a 5 to 6 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur; and


R^5 is H;
 C_1 - C_6 alkyl substituted with 0-3 R^{5b} ;
 C_2 - C_6 alkenyl substituted with 0-3 R^{5b} ;
 C_2 - C_6 alkynyl substituted with 0-3 R^{5b} ;
 C_3 - C_{10} carbocycle substituted with 0-3 R^{5c} ;
 C_6 - C_{10} aryl substituted with 0-3 R^{5c} ; or
5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{5c} ;

R^{5b} , at each occurrence, is independently selected from:
H, C_1 - C_6 alkyl, CF_3 , OR^{14} , Cl, F, Br, I, =O, CN, NO_2 ,
 $NR^{15}R^{16}$;
 C_3 - C_{10} carbocycle substituted with 0-3 R^{5c} ;
 C_6 - C_{10} aryl substituted with 0-3 R^{5c} ; or

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{5c} ;

R^{5c} , at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO_2 , $NR^{15}R^{16}$, CF_3 , acetyl, SCH_3 , $S(=O)CH_3$, $S(=O)_2CH_3$, C_1 - C_6 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl, C_1 - C_4 haloalkoxy, and C_1 - C_4 haloalkyl-S-;

Ring B is selected from:



R^{11} , at each occurrence, is independently selected from H, C_1 - C_4 alkoxy, Cl, F, Br, I, $=O$, CN, NO_2 , $NR^{18}R^{19}$, $C(=O)R^{17}$, $C(=O)OR^{17}$, $C(=O)NR^{18}R^{19}$, $S(=O)_2NR^{18}R^{19}$, CF_3 ; C_1 - C_6 alkyl optionally substituted with 0-3 R^{11a} ; C_6 - C_{10} aryl substituted with 0-3 R^{11b} ; C_3 - C_{10} carbocycle substituted with 0-3 R^{11b} ; or 5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{11b} ;

R^{11a} , at each occurrence, is independently selected from

- H, C₁-C₆ alkyl, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂, NR¹⁵R¹⁶, CF₃;
phenyl substituted with 0-3 R^{11b};
C₃-C₆ cycloalkyl substituted with 0-3 R^{11b}; and
5 5 to 6 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R^{11b};
- 10 R^{11b}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;
- 15 W is a bond;
X is a bond;
Y is a bond;
- 20 Z is H;
C₁-C₈ alkyl substituted with 0-3 R^{12a};
C₂-C₆ alkenyl substituted with 0-3 R^{12a};
C₂-C₆ alkynyl substituted with 0-3 R^{12a};
C₆-C₁₀ aryl substituted with 0-4 R^{12b};
25 C₃-C₁₀ carbocycle substituted with 0-4 R^{12b}; or
5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{12b};
- 30 R^{12a}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, -C(=O)NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,
35 C₁-C₄ haloalkoxy, C₁-C₄ haloalkyl-S-, C₆-C₁₀ aryl substituted with 0-4 R^{12b};
C₃-C₁₀ carbocycle substituted with 0-4 R^{12b}; or

5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered heterocycle
is substituted with 0-3 R^{12b};

5

R^{12b}, at each occurrence, is independently selected from
H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl,
SCH₃, S(=O)CH₃, S(=O)₂CH₃,
C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,
10 C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

R¹³, at each occurrence, is independently selected from
H, OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN,
NO₂, NR¹⁵R¹⁶, and CF₃;

15

R¹⁴ is H, phenyl, benzyl, C₁-C₆ alkyl, C₂-C₆ alkoxyalkyl, or
C₃-C₆ cycloalkyl;

R^{14a} is H, phenyl, benzyl, or C₁-C₄ alkyl;

20

R¹⁵, at each occurrence, is independently selected from H,
C₁-C₆ alkyl, benzyl, phenethyl, (C₁-C₆ alkyl)-C(=O)-,
and (C₁-C₆ alkyl)-S(=O)₂-;

25 R¹⁶, at each occurrence, is independently selected from
H, C₁-C₆ alkyl, benzyl, phenethyl,
(C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;

R¹⁷ is H, C₁-C₆ alkyl, C₂-C₆ alkoxyalkyl,

30

aryl substituted by 0-4 R^{17a}, or
-CH₂-aryl substituted by 0-4 R^{17a};

R^{17a} is H, methyl, ethyl, propyl, butyl, methoxy, ethoxy,
propoxy, butoxy, -OH, F, Cl, Br, I, CF₃, OCF₃, SCH₃,
35 S(O)CH₃, SO₂CH₃, -NH₂, -N(CH₃)₂, or C₁-C₄ haloalkyl;

R¹⁸, at each occurrence, is independently selected from
H, C₁-C₆ alkyl, phenyl, benzyl, phenethyl,

(C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;

R¹⁹, at each occurrence, is independently selected from
H, OH, methyl, ethyl, propyl, butyl, phenyl, benzyl,
5 phenethyl; and

R²³, at each occurrence, is independently selected from
H, OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN,
NO₂, NR¹⁵R¹⁶, and CF₃.

10

7. A compound of Formula (Ib) according to Claim 6, or a
pharmaceutically acceptable salt or prodrug thereof,
wherein:

15 R³ and R^{3a} are combined to form a 3-8 membered carbocyclic
moiety;

wherein said 3-8 membered carbocyclic moiety is
saturated or partially unsaturated;

wherein said 3-8 membered carbocyclic moiety is
20 substituted with 0-3 R⁴;

R⁴, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, CF₃, acetyl, SCH₃, S(=O)CH₃,
S(=O)₂CH₃, NR¹⁵R¹⁶, OR^{14a}, C₁-C₄ alkyl, C₂-C₄ alkenyl,
25 C₂-C₄ alkynyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,
C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

R⁵ is H;

C₁-C₆ alkyl substituted with 0-3 R^{5b};

30

C₂-C₆ alkenyl substituted with 0-3 R^{5b};

C₂-C₆ alkynyl substituted with 0-3 R^{5b};

C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};

C₆-C₁₀ aryl substituted with 0-3 R^{5c}; or

5 to 10 membered heterocycle containing 1 to 4

35

heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered heterocycle
is substituted with 0-3 R^{5c};

R^{5b}, at each occurrence, is independently selected from:

H, C₁-C₆ alkyl, CF₃, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂,
NR¹⁵R¹⁶;

C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};

5 C₆-C₁₀ aryl substituted with 0-3 R^{5c}; or


5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered heterocycle
is substituted with 0-3 R^{5c};

10

R^{5c}, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄
haloalkyl, and C₁-C₄ haloalkoxy;

15

Ring B is selected from:



20

R¹¹, at each occurrence, is independently selected from
H, =O, NR¹⁸R¹⁹, CF₃;

C₁-C₄ alkyl optionally substituted with 0-1 R^{11a};

25 phenyl substituted with 0-3 R^{11b};

C₃-C₇ carbocycle substituted with 0-3 R^{11b}; and

5 to 7 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and

sulphur, wherein said 5 to 7 membered heterocycle is
substituted with 0-3 R^{11b}; wherein said 5 to 7

30

5 membered heterocycle is selected from pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, pyrrolyl, piperazinyl, piperidinyl, homopiperidinyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

R^{11a}, at each occurrence, is independently selected from H, C₁-C₄ alkyl, OR¹⁴, F, Cl, =O, NR¹⁵R¹⁶, CF₃, or phenyl substituted with 0-3 R^{11b};

10 R^{11b}, at each occurrence, is independently selected from H, OH, Cl, F, NR¹⁵R¹⁶, CF₃, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, C₁-C₂ haloalkyl, and C₁-C₂ haloalkoxy;

15 W is a bond;
X is a bond;
Y is a bond;

20 Z is H;
C₁-C₈ alkyl substituted with 0-3 R^{12a};
C₂-C₆ alkenyl substituted with 0-3 R^{12a}; or
C₂-C₆ alkynyl substituted with 0-3 R^{12a};

25 R^{12a}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, -C(=O)NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, C₁-C₄ haloalkyl-S-,
30 C₆-C₁₀ aryl substituted with 0-4 R^{12b};
C₃-C₁₀ carbocycle substituted with 0-4 R^{12b}; or
5 to 10 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{12b}; and wherein said 5 to
35 10 membered heterocycle is selected from pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, pyrrolyl, pyrazolyl, imidazolyl,

oxazolyl, isoxazolyl, tetrazolyl, benzofuranyl,
benzothiofuranyl, indolyl, benzimidazolyl,
1H-indazolyl, oxazolidinyl, isoxazolidinyl,
benzotriazolyl, benzisoxazolyl, oxindolyl,
5 benzoxazolinyl, quinolinyl, and isoquinolinyl;

R^{12b}, at each occurrence, is independently selected from
H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl,
SCH₃, S(=O)CH₃, S(=O)₂CH₃,
10 C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,
C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

R¹³, at each occurrence, is independently selected from
H, OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN,
15 NO₂, NR¹⁵R¹⁶, and CF₃;

R¹⁴ is H, phenyl, benzyl, C₁-C₆ alkyl, C₂-C₆ alkoxyalkyl, or
C₃-C₆ cycloalkyl;

20 R^{14a} is H, phenyl, benzyl, or C₁-C₄ alkyl;

R¹⁵, at each occurrence, is independently selected from H,
C₁-C₆ alkyl, benzyl, phenethyl, (C₁-C₄ alkyl)-C(=O)-,
and (C₁-C₄ alkyl)-S(=O)₂-;

25

R¹⁶, at each occurrence, is independently selected from
H, C₁-C₆ alkyl, benzyl, phenethyl,
(C₁-C₄ alkyl)-C(=O)-, and (C₁-C₄ alkyl)-S(=O)₂-;

30 R¹⁸, at each occurrence, is independently selected from
H, C₁-C₆ alkyl, phenyl, benzyl, phenethyl,
(C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-; and

R¹⁹, at each occurrence, is independently selected from
35 H, OH, methyl, ethyl, propyl, butyl, phenyl, benzyl,
and phenethyl.

8. A compound of Formula (Ib) according to Claim 7 or a pharmaceutically acceptable salt or prodrug thereof wherein:

5 R³ and R^{3a} are combined to form a 3-6 membered carbocyclic moiety;
wherein said 3-6 membered carbocyclic moiety is saturated or partially unsaturated;
wherein said 3-6 membered carbocyclic moiety is
10 substituted with 0-2 R⁴;

R⁴, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, CF₃, acetyl, SCH₃, methyl, ethyl, methoxy, ethoxy, allyl, -OCF₃, and -SCF₃;

15

R⁵ is H;

C₁-C₄ alkyl substituted with 0-3 R^{5b};
C₂-C₄ alkenyl substituted with 0-3 R^{5b}; or
C₂-C₄ alkynyl substituted with 0-3 R^{5b};

20

R^{5b}, at each occurrence, is independently selected from:
H, methyl, ethyl, propyl, butyl, CF₃, OR¹⁴, Cl, F, Br, I, =O;

C₃-C₆ carbocycle substituted with 0-3 R^{5c};

25

phenyl substituted with 0-3 R^{5c}; or


5 to 6 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R^{5c};

30

R^{5c}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₄ alkyl, C₁-C₃ alkoxy, C₁-C₂ haloalkyl, and C₁-C₂ haloalkoxy;

35

Ring B is selected from:



- 5 R^{11} , at each occurrence, is independently selected from H, =O, $NR^{18}R^{19}$, CF_3 ;
 C_1 - C_4 alkyl optionally substituted with 0-1 R^{11a} ;
 phenyl substituted with 0-3 R^{11b} ;
 C_3 - C_7 carbocycle substituted with 0-3 R^{11b} ; and
- 10 5 to 7 membered heterocycle containing 1 to 4
 heteroatoms selected from nitrogen, oxygen, and
 sulphur, wherein said 5 to 7 membered heterocycle is
 substituted with 0-3 R^{11b} ; wherein said 5 to 7
 membered heterocycle is selected from pyridinyl,
- 15 pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,
 pyrrolyl, piperazinyl, piperidinyl, homopiperidinyl,
 pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and
 tetrazolyl;
- 20 R^{11a} , at each occurrence, is independently selected from H,
 methyl, ethyl, propyl, butyl, methoxy, ethoxy,
 propoxy, phenoxy, F, Cl, =O, $NR^{15}R^{16}$, CF_3 , or phenyl
 substituted with 0-3 R^{11b} ;
- 25 R^{11b} , at each occurrence, is independently selected from H,
 OH, Cl, F, $NR^{15}R^{16}$, CF_3 , methyl, ethyl, propyl, butyl,
 methoxy, ethoxy, propoxy, C_1 - C_2 haloalkyl, and C_1 - C_2
 haloalkoxy;
- 30 W is a bond;

X is a bond;

Y is a bond;

Z is H;

- 5 C₁-C₄ alkyl substituted with 0-3 R^{12a};
 C₂-C₄ alkenyl substituted with 0-3 R^{12a}; or
 C₂-C₄ alkynyl substituted with 0-3 R^{12a};

- 10 R^{12a}, at each occurrence, is independently selected from
 H, OH, Cl, F, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃,
 S(=O)₂CH₃, C₁-C₄ alkyl, C₁-C₃ alkoxy, C₁-C₂ haloalkyl,
 and C₁-C₂ haloalkoxy;

- 15 R¹³, at each occurrence, is independently selected from
 H, OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN,
 NO₂, NR¹⁵R¹⁶, and CF₃;

 R¹⁴ is H, phenyl, benzyl, C₁-C₄ alkyl, or C₂-C₄ alkoxyalkyl;

- 20 R¹⁵, at each occurrence, is independently selected from H,
 C₁-C₄ alkyl, and benzyl;

- R¹⁶, at each occurrence, is independently selected from
 H, methyl, ethyl, propyl, butyl, benzyl, phenethyl,
25 methyl-C(=O)-, ethyl-C(=O)-,
 methyl-S(=O)₂-, and ethyl-S(=O)₂-;

- R¹⁸, at each occurrence, is independently selected from
 H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and
30 phenethyl; and

 R¹⁹, at each occurrence, is independently selected from
 H, methyl, ethyl, propyl, and butyl.

- 35 9. A compound of Formula (Ib) according to Claim 8 or a
 pharmaceutically acceptable salt or prodrug thereof
 wherein:

R³ and R^{3a} are combined to form a 3-6 membered carbocyclic moiety selected from cyclopropyl, cyclobutyl, cyclopentyl, cyclopentenyl, cyclohexyl, and cyclohexenyl; wherein said 3-6 membered carbocyclic moiety is substituted with 0-1 R⁴;

R⁴ is selected from H, OH, Cl, F, CN, CF₃, methyl, ethyl, methoxy, ethoxy, allyl, and -OCF₃;


R⁵ is C₁-C₄ alkyl substituted with 0-1 R^{5b};
C₂-C₄ alkenyl substituted with 0-1 R^{5b};
C₂-C₄ alkynyl substituted with 0-1 R^{5b};

R^{5b}, at each occurrence, is independently selected from:

H, methyl, ethyl, propyl, butyl, CF₃, OR¹⁴, =O;
C₃-C₆ carbocycle substituted with 0-2 R^{5c};
phenyl substituted with 0-3 R^{5c}; or
5 to 6 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 6 membered heterocycle is substituted with 0-3 R^{5c}; wherein said 5 to 6 membered heterocycle is selected from pyridinyl, pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl, pyrrolyl, piperazinyl, piperidinyl, pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and tetrazolyl;

R^{5c}, at each occurrence, is independently selected from H, OH, Cl, F, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, methyl, ethyl, propyl, butyl, methoxy, ethoxy, propoxy, C₁-C₂ haloalkyl, and C₁-C₂ haloalkoxy;

Ring B is selected from:



- 5 R¹¹, at each occurrence, is independently selected from
 H, =O, NR¹⁸R¹⁹;
 C₁-C₄ alkyl optionally substituted with 0-1 R^{11a};
 phenyl substituted with 0-3 R^{11b};
 cyclohexyl substituted with 0-3 R^{11b};
 10 cycloheptyl substituted with 0-3 R^{11b};
 5 to 7 membered heterocycle containing 1 to 4
 heteroatoms selected from nitrogen, oxygen, and
 sulphur, wherein said 5 to 7 membered heterocycle is
 substituted with 0-3 R^{11b}; wherein said 5 to 7
 15 membered heterocycle is selected from pyridinyl,
 pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,
 pyrrolyl, piperazinyl, piperidinyl, homopiperidinyl,
 pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and
 tetrazolyl;
 20 R^{11a}, at each occurrence, is independently selected from H,
 methyl, ethyl, propyl, methoxy, ethoxy, propoxy,
 phenoxy, F, Cl, =O, NR¹⁵R¹⁶, CF₃, or phenyl substituted
 with 0-3 R^{11b};
 25 R^{11b}, at each occurrence, is independently selected from H,
 OH, Cl, F, NR¹⁵R¹⁶, CF₃, methyl, ethyl, propyl, butyl,
 methoxy, ethoxy, propoxy, C₁-C₂ haloalkyl, and C₁-C₂
 haloalkoxy;
 30 W is a bond;

X is a bond;

Y is a bond;

Z is H;

- 5 C₁-C₄ alkyl substituted with 0-1 R^{12a};
 C₂-C₄ alkenyl substituted with 0-1 R^{12a}; or
 C₂-C₄ alkynyl substituted with 0-1 R^{12a};

- 10 R^{12a}, at each occurrence, is independently selected from
 H, OH, Cl, F, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃,
 S(=O)₂CH₃, methyl, ethyl, propyl, butyl, methoxy,
 ethoxy, propoxy, C₁-C₂ haloalkyl, and C₁-C₂ haloalkoxy;

- 15 R¹³, at each occurrence, is independently selected from
 H, OH, methyl, ethyl, propyl, butyl, methoxy, ethoxy,
 Cl, F, Br, CN, NR¹⁵R¹⁶, and CF₃;

R¹⁴ is H, phenyl, benzyl, methyl, ethyl, propyl, or butyl;

- 20 R¹⁵, at each occurrence, is independently selected from H,
 methyl, ethyl, propyl, and butyl;

- 25 R¹⁶, at each occurrence, is independently selected from
 H, methyl, ethyl, propyl, butyl, benzyl, and
 phenethyl;

- 30 R¹⁸, at each occurrence, is independently selected from
 H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and
 phenethyl; and

- R¹⁹, at each occurrence, is independently selected from
 H, methyl, ethyl, propyl, and butyl.

- 35 10. A compound of Formula (Ib) according to Claim 9
 wherein:

R³ and R^{3a} are combined to form cyclobutyl, cyclopentyl,
 cyclopentenyl, or cyclohexyl;

R^5 is $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2CH_3$, $-CH(CH_3)_2$, $-CH_2CH_2CH_2CH_3$,
 $-CH(CH_3)CH_2CH_3$, $-CH_2CH(CH_3)_2$, $-CH_2C(CH_3)_3$,
 $-CH_2CH_2CH_2CH_2CH_3$, $-CH(CH_3)CH_2CH_2CH_3$, $-CH_2CH(CH_3)CH_2CH_3$,
5 $-CH_2CH_2CH(CH_3)_2$, $-CH(CH_2CH_3)_2$, $-CF_3$, $-CH_2CF_3$, $-CH_2CH_2CF_3$,
 $-CH_2CH_2CH_2CF_3$, $-CH_2CH_2CH_2CH_2CF_3$, $-CH=CH_2$, $-CH_2CH=CH_2$,
 $-CH=CHCH_3$, *cis*- $CH_2CH=CH(CH_3)$, *trans*- $CH_2CH=CH(CH_3)$,
trans- $CH_2CH=CH(C_6H_5)$, $-CH_2CH=C(CH_3)_2$, *cis*- $CH_2CH=CHCH_2CH_3$,
trans- $CH_2CH=CHCH_2CH_3$, *cis*- $CH_2CH_2CH=CH(CH_3)$,
10 *trans*- $CH_2CH_2CH=CH(CH_3)$, *trans*- $CH_2CH=CHCH_2(C_6H_5)$,
 $-C\equiv CH$, $-CH_2C\equiv CH$, $-CH_2C\equiv C(CH_3)$, $-CH_2C\equiv C(C_6H_5)$
 $-CH_2CH_2C\equiv CH$, $-CH_2CH_2C\equiv C(CH_3)$, $-CH_2CH_2C\equiv C(C_6H_5)$
 $-CH_2CH_2CH_2C\equiv CH$, $-CH_2CH_2CH_2C\equiv C(CH_3)$, $-CH_2CH_2CH_2C\equiv C(C_6H_5)$
cyclopropyl- CH_2- , cyclobutyl- CH_2- , cyclopentyl- CH_2- ,
15 cyclohexyl- CH_2- , (2- CH_3 -cyclopropyl) CH_2- ,
(3- CH_3 -cyclobutyl) CH_2- ,
cyclopropyl- CH_2CH_2- , cyclobutyl- CH_2CH_2- ,
cyclopentyl- CH_2CH_2- , cyclohexyl- CH_2CH_2- ,
(2- CH_3 -cyclopropyl) CH_2CH_2- , (3- CH_3 -cyclobutyl) CH_2CH_2- ,
20 phenyl- CH_2- , (2-F-phenyl) CH_2- , (3-F-phenyl) CH_2- ,
(4-F-phenyl) CH_2- , furanyl- CH_2- , thienyl- CH_2- ,
pyridyl- CH_2- , 1-imidazolyl- CH_2- , oxazolyl- CH_2- ,
isoxazolyl- CH_2- , phenyl- CH_2CH_2- , (2-F-phenyl) CH_2CH_2- ,
(3-F-phenyl) CH_2CH_2- , (4-F-phenyl) CH_2CH_2- ,
25 furanyl- CH_2CH_2- , thienyl- CH_2CH_2- , pyridyl- CH_2CH_2- ,
1-imidazolyl- CH_2CH_2- , oxazolyl- CH_2CH_2- ,
isoxazolyl- CH_2CH_2- ,

W is a bond;

30 X is a bond;

Y is a bond;

Z is methyl, ethyl, i-propyl, n-propyl, n-butyl, i-butyl,
s-butyl, t-butyl, or allyl;

35

R^{11} , at each occurrence, is independently selected from
H, =O, methyl, ethyl, phenyl, benzyl, phenethyl,
4-F-phenyl, (4-F-phenyl) CH_2- , (4-F-phenyl) CH_2CH_2- ,

3-F-phenyl, (3-F-phenyl)CH₂-, (3-F-phenyl)CH₂CH₂-,
2-F-phenyl, (2-F-phenyl)CH₂-, (2-F-phenyl)CH₂CH₂-,
4-Cl-phenyl, (4-Cl-phenyl)CH₂-, (4-Cl-phenyl)CH₂CH₂-,
3-Cl-phenyl, (3-Cl-phenyl)CH₂-, (3-Cl-phenyl)CH₂CH₂-,
5 4-CH₃-phenyl, (4-CH₃-phenyl)CH₂-, (4-CH₃-phenyl)CH₂CH₂-,
3-CH₃-phenyl, (3-CH₃-phenyl)CH₂-, (3-CH₃-phenyl)CH₂CH₂-,
4-CF₃-phenyl, (4-CF₃-phenyl)CH₂-, (4-CF₃-phenyl)CH₂CH₂-,
pyrid-2-yl, pyrid-3-yl, pyrid-4-yl, cyclohexyl,
cycloheptyl, piperidinyl, or homopiperidinyl; and

10 R¹³, at each occurrence, is independently selected from
H, F, Cl, OH, -CH₃, -CH₂CH₃, -OCH₃, or -CF₃.

11. A compound of Formula (Ib) according to Claim 9
15 wherein:

R³ and R^{3a} are combined to form cyclobutyl, cyclopentyl,
cyclopentenyl, or cyclohexyl;

20 R⁵ is -CH₃, -CH₂CH₃, -CH₂CH₂CH₃, -CH(CH₃)₂, -CH₂CH₂CH₂CH₃,
-CH(CH₃)CH₂CH₃, -CH₂CH(CH₃)₂, -CH₂C(CH₃)₃,
-CH₂CH₂CH₂CH₂CH₃, -CH(CH₃)CH₂CH₂CH₃, -CH₂CH(CH₃)CH₂CH₃,
-CH₂CH₂CH(CH₃)₂, -CH(CH₂CH₃)₂, cyclopropyl-CH₂-,
cyclobutyl-CH₂-, (2-CH₃-cyclopropyl)CH₂-, or
25 (3-CH₃-cyclobutyl)CH₂-;

W is a bond;

X is a bond;

Y is a bond;

30 Z is methyl, ethyl, i-propyl, n-propyl, n-butyl, i-butyl,
s-butyl, t-butyl, or allyl;


R¹¹, at each occurrence, is independently selected from

35 H, =O, methyl, ethyl, phenyl, benzyl, phenethyl,
4-F-phenyl, (4-F-phenyl)CH₂-, (4-F-phenyl)CH₂CH₂-,
3-F-phenyl, (3-F-phenyl)CH₂-, (3-F-phenyl)CH₂CH₂-,
2-F-phenyl, (2-F-phenyl)CH₂-, (2-F-phenyl)CH₂CH₂-,

4-Cl-phenyl, (4-Cl-phenyl)CH₂-, (4-Cl-phenyl)CH₂CH₂-,
 3-Cl-phenyl, (3-Cl-phenyl)CH₂-, (3-Cl-phenyl)CH₂CH₂-,
 4-CH₃-phenyl, (4-CH₃-phenyl)CH₂-, (4-CH₃-phenyl)CH₂CH₂-,
 3-CH₃-phenyl, (3-CH₃-phenyl)CH₂-, (3-CH₃-phenyl)CH₂CH₂-,
 5 4-CF₃-phenyl, (4-CF₃-phenyl)CH₂-, (4-CF₃-phenyl)CH₂CH₂-,
 pyrid-2-yl, pyrid-3-yl, pyrid-4-yl, cyclohexyl,
 cycloheptyl, piperidinyl, or homopiperidinyl; and

R¹³, at each occurrence, is independently selected from
 10 H, F, Cl, OH, -CH₃, -CH₂CH₃, -OCH₃, or -CF₃.

12. A compound according to Claim 4 of Formula (Ib):



15

or a pharmaceutically acceptable salt or prodrug thereof,
 wherein:

20 R³ and R^{3a} are combined to form a 3-8 membered carbocyclic
 moiety;

wherein said 3-8 membered carbocyclic moiety is
 saturated or partially unsaturated;

wherein said 3-8 membered carbocyclic moiety is
 25 substituted with 0-3 R⁴;

R⁴, at each occurrence, is independently selected from H,
 OH, Cl, F, Br, I, CN, NO₂, CF₃, acetyl, SCH₃, S(=O)CH₃,
 S(=O)₂CH₃, NR¹⁵R¹⁶, OR^{14a}, C₁-C₄ alkyl, C₂-C₄ alkenyl,
 30 C₂-C₄ alkynyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl,
 C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

R⁵ is H;

C₁-C₆ alkyl substituted with 0-3 R^{5b};
 35 C₂-C₆ alkenyl substituted with 0-3 R^{5b};

C₂-C₆ alkynyl substituted with 0-3 R^{5b};
 C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};
 C₆-C₁₀ aryl substituted with 0-3 R^{5c}; or
 5 to 10 membered heterocycle containing 1 to 4
 5 heteroatoms selected from nitrogen, oxygen, and
 sulphur, wherein said 5 to 10 membered heterocycle
 is substituted with 0-3 R^{5c};

R^{5b}, at each occurrence, is independently selected from:

10 H, C₁-C₆ alkyl, CF₃, OR¹⁴, Cl, F, Br, I, =O, CN, NO₂,
 NR¹⁵R¹⁶;

C₃-C₁₀ carbocycle substituted with 0-3 R^{5c};

C₆-C₁₀ aryl substituted with 0-3 R^{5c}; or

5 to 10 membered heterocycle containing 1 to 4


15 heteroatoms selected from nitrogen, oxygen, and
 sulphur, wherein said 5 to 10 membered heterocycle
 is substituted with 0-3 R^{5c};

R^{5c}, at each occurrence, is independently selected from H,

20 OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
 S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄
 haloalkyl, and C₁-C₄ haloalkoxy;

Ring B is selected from:

25



30 R¹¹, at each occurrence, is independently selected from

H, =O, NR¹⁸R¹⁹, CF₃;

C₁-C₄ alkyl optionally substituted with 0-1 R^{11a};

phenyl substituted with 0-3 R^{11b};

C₃-C₇ carbocycle substituted with 0-3 R^{11b}; and

5 5 to 7 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 7 membered heterocycle is
substituted with 0-3 R^{11b}; wherein said 5 to 7
membered heterocycle is selected from pyridinyl,
10 pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,
pyrrolyl, piperazinyl, piperidinyl, homopiperidinyl,
pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and
tetrazolyl;

15 R^{11a}, at each occurrence, is independently selected from H,
C₁-C₄ alkyl, OR¹⁴, F, Cl, =O, NR¹⁵R¹⁶, CF₃, or phenyl
substituted with 0-3 R^{11b};

20 R^{11b}, at each occurrence, is independently selected from H,
OH, Cl, F, NR¹⁵R¹⁶, CF₃, methyl, ethyl, propyl, butyl,
methoxy, ethoxy, propoxy, C₁-C₂ haloalkyl, and C₁-C₂
haloalkoxy;

W is a bond, -CH₂-, -CH₂CH₂-;

25

X is a bond;

phenyl substituted with 0-2 R^{Xb};

C₃-C₆ cycloalkyl substituted with 0-2 R^{Xb}; or

5 to 6 membered heterocycle substituted with 0-2 R^{Xb};

30

R^{Xb}, at each occurrence, is independently selected from H,
OH, Cl, F, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃,
S(=O)₂CH₃, C₁-C₄ alkyl, C₁-C₃ alkoxy, C₁-C₂ haloalkyl,
and C₁-C₂ haloalkoxy;

35

Y is a bond, -C(=O)-, -O-, -S-, -S(=O)-, -S(=O)₂-,
-N(R¹⁹)-, -C(=O)NR^{19b}-, -NR^{19b}C(=O)-, -NR^{19b}S(=O)₂-,
-S(=O)₂NR^{19b}-, -NR^{19b}S(=O)-, -S(=O)NR^{19b}-, -C(=O)O-,

or $-\text{OC}(=\text{O})-$;

Z is C_1 - C_3 alkyl substituted with 1-2 $\text{R}^{12\text{a}}$;

C_6 - C_{10} aryl substituted with 0-4 $\text{R}^{12\text{b}}$;

5 C_3 - C_{10} carbocycle substituted with 0-3 $\text{R}^{12\text{b}}$; or

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 $\text{R}^{12\text{b}}$;

10

$\text{R}^{12\text{a}}$, at each occurrence, is independently selected from C_6 - C_{10} aryl substituted with 0-4 $\text{R}^{12\text{b}}$;

C_3 - C_{10} carbocycle substituted with 0-4 $\text{R}^{12\text{b}}$; and

5 to 10 membered heterocycle containing 1 to 4

15 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 $\text{R}^{12\text{b}}$;

$\text{R}^{12\text{b}}$, at each occurrence, is independently selected from

20 H, OH, Cl, F, Br, I, CN, NO_2 , $\text{NR}^{15}\text{R}^{16}$, CF_3 , acetyl, SCH_3 , $\text{S}(=\text{O})\text{CH}_3$, $\text{S}(=\text{O})_2\text{CH}_3$, C_1 - C_6 alkyl, C_1 - C_4 alkoxy, C_1 - C_4 haloalkyl, C_1 - C_4 haloalkoxy, and C_1 - C_4 haloalkyl-S-;

25 R^{13} , at each occurrence, is independently selected from H, OH, C_1 - C_6 alkyl, C_1 - C_4 alkoxy, Cl, F, Br, I, CN, NO_2 , $\text{NR}^{15}\text{R}^{16}$, and CF_3 ;

R^{14} is H, phenyl, benzyl, C_1 - C_6 alkyl, C_2 - C_6 alkoxyalkyl, or
30 C_3 - C_6 cycloalkyl;

$\text{R}^{14\text{a}}$ is H, phenyl, benzyl, or C_1 - C_4 alkyl;

R^{15} , at each occurrence, is independently selected from H,
35 C_1 - C_6 alkyl, benzyl, phenethyl, $(\text{C}_1$ - C_4 alkyl)- $\text{C}(=\text{O})-$, and $(\text{C}_1$ - C_4 alkyl)- $\text{S}(=\text{O})_2-$;

R^{16} , at each occurrence, is independently selected from

H, OH, C₁-C₆ alkyl, benzyl, phenethyl,
(C₁-C₄ alkyl)-C(=O)-, and (C₁-C₄ alkyl)-S(=O)₂-;

5 R¹⁸, at each occurrence, is independently selected from
H, C₁-C₆ alkyl, phenyl, benzyl, phenethyl,
(C₁-C₆ alkyl)-C(=O)-, and (C₁-C₆ alkyl)-S(=O)₂-;

10 R¹⁹, at each occurrence, is independently selected from
H, OH, methyl, ethyl, propyl, butyl, phenyl, benzyl,
and phenethyl; and

15 R^{19b}, at each occurrence, is independently selected from
H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and
phenethyl.

13. A compound of Formula (Ib) according to Claim 12 or a
pharmaceutically acceptable salt or prodrug thereof
wherein:

20 R³ and R^{3a} are combined to form a 3-6 membered carbocyclic
moiety;
wherein said 3-6 membered carbocyclic moiety is
saturated or partially unsaturated;
wherein said 3-6 membered carbocyclic moiety is
25 substituted with 0-2 R⁴;

30 R⁴, at each occurrence, is independently selected from H,
OH, Cl, F, Br, I, CN, NO₂, CF₃, acetyl, SCH₃, methyl,
ethyl, methoxy, ethoxy, allyl, -OCF₃, and -SCF₃;

R⁵ is H;

C₁-C₄ alkyl substituted with 0-3 R^{5b};
C₂-C₄ alkenyl substituted with 0-3 R^{5b}; or
C₂-C₄ alkynyl substituted with 0-3 R^{5b};


35

R^{5b}, at each occurrence, is independently selected from:
H, methyl, ethyl, propyl, butyl, CF₃, OR¹⁴, Cl, F, Br,
I, =O;

C₃-C₆ carbocycle substituted with 0-3 R^{5c};
 phenyl substituted with 0-3 R^{5c}; or
 5 to 6 membered heterocycle containing 1 to 4
 heteroatoms selected from nitrogen, oxygen, and
 sulphur, wherein said 5 to 6 membered heterocycle
 is substituted with 0-3 R^{5c};

R^{5c}, at each occurrence, is independently selected from H,
 OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃,
 S(=O)CH₃, S(=O)₂CH₃, C₁-C₄ alkyl, C₁-C₃ alkoxy, C₁-C₂
 haloalkyl, and C₁-C₂ haloalkoxy;

Ring B is selected from:



R¹¹, at each occurrence, is independently selected from
 H, =O, NR¹⁸R¹⁹, CF₃;

C₁-C₄ alkyl optionally substituted with 0-1 R^{11a};

phenyl substituted with 0-3 R^{11b};

C₃-C₇ carbocycle substituted with 0-3 R^{11b}; and

5 to 7 membered heterocycle containing 1 to 4

heteroatoms selected from nitrogen, oxygen, and
 sulphur, wherein said 5 to 7 membered heterocycle is
 substituted with 0-3 R^{11b}; wherein said 5 to 7
 membered heterocycle is selected from pyridinyl,
 pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,
 pyrrolyl, piperazinyl, piperidinyl, homopiperidinyl,

pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and
tetrazolyl;

5 R^{11a}, at each occurrence, is independently selected from H,
methyl, ethyl, propyl, butyl, methoxy, ethoxy,
propoxy, phenoxy, F, Cl, =O, NR¹⁵R¹⁶, CF₃, or phenyl
substituted with 0-3 R^{11b};

10 R^{11b}, at each occurrence, is independently selected from H,
OH, Cl, F, NR¹⁵R¹⁶, CF₃, methyl, ethyl, propyl, butyl,
methoxy, ethoxy, propoxy, C₁-C₂ haloalkyl, and C₁-C₂
haloalkoxy;

15 W is a bond, -CH₂-, -CH₂CH₂-;

X is a bond;

phenyl substituted with 0-1 R^{Xb};

C₃-C₆ cycloalkyl substituted with 0-1 R^{Xb}; or

5 to 6 membered heterocycle substituted with 0-1 R^{Xb};

20 R^{Xb} is selected from H, OH, Cl, F, NR¹⁵R¹⁶, CF₃, acetyl,
SCH₃, S(=O)CH₃, S(=O)₂CH₃, methyl, ethyl, propyl,
methoxy, ethoxy, propoxy, and -OCF₃;

25 Y is a bond, -C(=O)-, -O-, -S-, -S(=O)-, -S(=O)₂-, -NH-,
-N(CH₃)-, or -N(CH₂CH₃)-;

Z is C₁-C₂ alkyl substituted with 1-2 R^{12a};

C₆-C₁₀ aryl substituted with 0-4 R^{12b};

30 C₃-C₁₀ carbocycle substituted with 0-3 R^{12b}; or

5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered heterocycle
is substituted with 0-3 R^{12b};

35

R^{12a}, at each occurrence, is independently selected from

C₆-C₁₀ aryl substituted with 0-4 R^{12b};

C₃-C₁₀ carbocycle substituted with 0-4 R^{12b}; and

5 to 10 membered heterocycle containing 1 to 4 heteroatoms selected from nitrogen, oxygen, and sulphur, wherein said 5 to 10 membered heterocycle is substituted with 0-3 R^{12b};

5

R^{12b}, at each occurrence, is independently selected from H, OH, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃, S(=O)₂CH₃, C₁-C₆ alkyl, C₁-C₄ alkoxy, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, and C₁-C₄ haloalkyl-S-;

10

R¹³, at each occurrence, is independently selected from H, OH, C₁-C₆ alkyl, C₁-C₄ alkoxy, Cl, F, Br, I, CN, NO₂, NR¹⁵R¹⁶, and CF₃;

15

R¹⁴ is H, phenyl, benzyl, C₁-C₄ alkyl, or C₂-C₄ alkoxyalkyl;

R¹⁵, at each occurrence, is independently selected from H, C₁-C₄ alkyl, and benzyl;

20

R¹⁶, at each occurrence, is independently selected from H, OH, methyl, ethyl, propyl, butyl, benzyl, phenethyl, methyl-C(=O)-, ethyl-C(=O)-, methyl-S(=O)₂-, and ethyl-S(=O)₂-;

25

R¹⁸, at each occurrence, is independently selected from H, methyl, ethyl, propyl, butyl, phenyl, benzyl, and phenethyl; and

30 R¹⁹, at each occurrence, is independently selected from H, methyl, ethyl, propyl, and butyl.

14. A compound of Formula (Ib) according to Claim 13 or a pharmaceutically acceptable salt or prodrug thereof wherein:

35

R³ and R^{3a} are combined to form a 3-6 membered carbocyclic moiety selected from cyclopropyl, cyclobutyl,

cyclopentyl, cyclopentenyl, cyclohexyl, and cyclohexenyl; wherein said 3-6 membered carbocyclic moiety is substituted with 0-1 R⁴;

5 R⁴ is selected from H, OH, Cl, F, CN, CF₃, methyl, ethyl, methoxy, ethoxy, allyl, and -OCF₃;


R⁵ is C₁-C₄ alkyl substituted with 0-1 R^{5b};
 C₂-C₄ alkenyl substituted with 0-1 R^{5b};
 10 C₂-C₄ alkynyl substituted with 0-1 R^{5b};


R^{5b}, at each occurrence, is independently selected from:
 H, methyl, ethyl, propyl, butyl, CF₃, OR¹⁴, =O;
 C₃-C₆ carbocycle substituted with 0-2 R^{5c};
 15 phenyl substituted with 0-3 R^{5c}; or
 5 to 6 membered heterocycle containing 1 to 4
 heteroatoms selected from nitrogen, oxygen, and
 sulphur, wherein said 5 to 6 membered heterocycle
 is substituted with 0-3 R^{5c}; wherein said 5 to 6
 20 membered heterocycle is selected from pyridinyl,
 pyrimidinyl, triazinyl, furanyl, thienyl,
 thiazolyl, pyrrolyl, piperazinyl, piperidinyl,
 pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and
 tetrazolyl;

25 R^{5c}, at each occurrence, is independently selected from H,
 OH, Cl, F, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃,
 S(=O)₂CH₃, methyl, ethyl, propyl, butyl, methoxy,
 ethoxy, propoxy, C₁-C₂ haloalkyl, and C₁-C₂ haloalkoxy;

30

Ring B is selected from:





R^{11} , at each occurrence, is independently selected from
H, =O, $NR^{18}R^{19}$;

5 C_1 - C_4 alkyl optionally substituted with 0-1 R^{11a} ;

phenyl substituted with 0-3 R^{11b} ;

cyclohexyl substituted with 0-3 R^{11b} ;

cycloheptyl substituted with 0-3 R^{11b} ;

5 to 7 membered heterocycle containing 1 to 4

10 heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 7 membered heterocycle is
substituted with 0-3 R^{11b} ; wherein said 5 to 7

membered heterocycle is selected from pyridinyl,

pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,

15 pyrrolyl, piperazinyl, piperidinyl, homopiperidinyl,

pyrazolyl, imidazolyl, oxazolyl, isoxazolyl, and

tetrazolyl;

R^{11a} , at each occurrence, is independently selected from H,

20 methyl, ethyl, propyl, methoxy, ethoxy, propoxy,

phenoxy, F, Cl, =O, $NR^{15}R^{16}$, CF_3 , or phenyl substituted
with 0-3 R^{11b} ;

R^{11b} , at each occurrence, is independently selected from H,

25 OH, Cl, F, $NR^{15}R^{16}$, CF_3 , methyl, ethyl, propyl, butyl,

methoxy, ethoxy, propoxy, C_1 - C_2 haloalkyl, and C_1 - C_2

haloalkoxy;

W is a bond or $-CH_2-$;

30

X is a bond;

phenyl substituted with 0-1 R^{Xb} ;

C_3 - C_6 cycloalkyl substituted with 0-1 R^{Xb} ; or

5 to 6 membered heterocycle substituted with 0-1 R^{Xb} ;

R^{Xb} is selected from H, OH, Cl, F, NR¹⁵R¹⁶, CF₃, acetyl, methyl, ethyl, methoxy, ethoxy, and -OCF₃;

5 Y is a bond, -C(=O)-, -O-, -S-, -S(=O)-, -S(=O)₂-, -NH-, -N(CH₃)-, or -N(CH₂CH₃)-;

Z is C₁-C₂ alkyl substituted with 1-2 R^{12a};
C₆-C₁₀ aryl substituted with 0-4 R^{12b};
10 C₃-C₁₀ carbocycle substituted with 0-3 R^{12b}; or
5 to 10 membered heterocycle containing 1 to 4
heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered heterocycle
is substituted with 0-3 R^{12b};

15 R^{12a}, at each occurrence, is independently selected from
C₆-C₁₀ aryl substituted with 0-4 R^{12b};
C₃-C₁₀ carbocycle substituted with 0-4 R^{12b}; and
5 to 10 membered heterocycle containing 1 to 4
20 heteroatoms selected from nitrogen, oxygen, and
sulphur, wherein said 5 to 10 membered heterocycle
is substituted with 0-3 R^{12b}; and wherein said 5 to
10 membered heterocycle is selected from pyridinyl,
pyrimidinyl, triazinyl, furanyl, thienyl, thiazolyl,
25 pyrrolyl, pyrazolyl, imidazolyl, oxazolyl,
isoxazolyl, tetrazolyl, benzofuranyl,
benzothiofuranyl, indolyl, benzimidazolyl,
1H-indazolyl, oxazolidinyl, isoxazolidinyl,
benzotriazolyl, benzisoxazolyl, oxindolyl,
30 benzoxazoliny, quinolinyl, and isoquinolinyl;

R^{12b}, at each occurrence, is independently selected from
H, OH, Cl, F, NR¹⁵R¹⁶, CF₃, acetyl, SCH₃, S(=O)CH₃,
S(=O)₂CH₃, methyl, ethyl, propyl, butyl, methoxy,
35 ethoxy, propoxy, and -OCF₃;

R¹³, at each occurrence, is independently selected from

- 1-[3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-(4-chlorophenyl)-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclohexanecarboxylic amide;
- 5 1-[3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclohexanecarboxylic amide;
- 10 1-[3-Methyl-1-(5-methyl-6-oxo-6,7-dihydro-5H-dibenzo[b,d]-azepin-7-ylcarbamoyl)-butyl]-cyclohexanecarboxylic acid amide;
- 15 1-[3-Methyl-1-(5-methyl-6-oxo-6,7-dihydro-5H-dibenzo[b,d]-azepin-7-ylcarbamoyl)-butyl]-cyclohexanecarboxylic acid amide;
- 20 1-(1-{1-[3-(2-Fluoro-phenoxy)-benzyl]-2-oxo-2,3,4,5-tetrahydro-1H-benzo[b]azepin-3-ylcarbamoyl}-3-methyl-butyl)-cyclopent-3-enecarboxylic acid amide;
- 25 1-{3-Methyl-1-[2-oxo-1-(3-phenylamino-benzyl)-2,3,4,5-tetrahydro-1H-benzo[b]azepin-3-ylcarbamoyl]-butyl}-cyclopentanecarboxylic acid amide;
- 30 1-[2-Cyclopropyl-1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-ylcarbamoyl]-ethyl]-cyclopent-3-enecarboxylic amide;
- 35 1-[2-Cyclopropyl-1-[1,3-dihydro-1-methyl-2-oxo-5-(2-trifluoromethylphenyl)-2H-1,4-benzodiazepin-3-ylcarbamoyl]-ethyl]-cyclopent-3-enecarboxylic amide;

1-[2-Cyclopropyl-1-(5-methyl-6-oxo-6,7-dihydro-5H-dibenzo[b,d]azepin-7-ylcarbamoyl)-ethyl]-cyclopent-3-enecarboxylic acid amide;

- 5 1-{3-Methyl-1-[2-oxo-1-(3-o-tolylamino-benzyl)-azepan-3-ylcarbamoyl]-butyl}-cyclopent-3-enecarboxylic acid amide;

- 1-[3-Methyl-1-(5-methyl-6-oxo-6,7-dihydro-5H-dibenzo[b,d]azepin-7-ylcarbamoyl)-butyl]-cyclopent-3-enecarboxylic acid amide;
- 10

1-[3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-(4-chlorophenyl)-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclopent-3-enecarboxylic amide;

15

1-[3-Methyl-1-(5-methyl-6-oxo-6,7-dihydro-5H-dibenzo[b,d]azepin-7-ylcarbamoyl)-butyl]-cyclopentanecarboxylic acid amide;

- 20 1-[3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-(4-chlorophenyl)-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclopentanecarboxylic amide;

- 1-[3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-(4-trifluoromethylphenyl)-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclopent-3-enecarboxylic amide;
- 25

- 1-[3-methyl-1-[1,3-dihydro-1-(i-propyl)-2-oxo-5-(2-fluorophenyl)-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclopent-3-enecarboxylic amide;
- 30

- 1-[3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-(4-trifluoromethylphenyl)-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclopentanecarboxylic amide;
- 35

1-[3-methyl-1-[1,3-dihydro-1-(2-cyclopropylethyl)-2-oxo-5-(2-fluorophenyl)-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclopent-3-enecarboxylic amide;

1-[3-methyl-1-[1,3-dihydro-1-(2-methylpropyl)-2-oxo-5-(2-fluorophenyl)-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclopent-3-enecarboxylic amide;

5

1-[3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-(4-chlorophenyl)-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclobutanecarboxylic amide;

10 1-[3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-phenyl-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclobutanecarboxylic amide; and

15 1-[3-methyl-1-[1,3-dihydro-1-methyl-2-oxo-5-cycloheptyl-2H-1,4-benzodiazepin-3-ylcarbamoyl]-butyl]-cyclopent-3-enecarboxylic amide.

22. A pharmaceutical composition comprising a compound according to one of Claims 1-21 and a pharmaceutically acceptable carrier.

23. A method for the treatment of neurological disorders associated with β -amyloid production comprising administering to a host in need of such treatment a therapeutically effective amount of a compound according to one of Claims 1-21.

24. A method for inhibiting γ -secretase activity comprising administering to a host in need of such inhibition a therapeutically effective amount of a compound according to one of Claims 1-21 that inhibits γ -secretase activity.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/17865

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D243/24 C07D223/18 C07D223/16 C07D223/12 C07D243/14
 C07D243/12 A61K31/55 A61K31/5513 A61P25/28

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 00 07995 A (DU PONT PHARM CO) 17 February 2000 (2000-02-17) cited in the application table 1 claims 1,19-25	1,22,23
A	WO 00 28331 A (DU PONT PHARM CO) 18 May 2000 (2000-05-18) claims 6,11,12	1,22,23
A	WO 99 66934 A (AUDIA JAMES E ; ELAN PHARM INC (US); LILLY CO ELI (US); SHI QING (U) 29 December 1999 (1999-12-29) page 25 -page 30 claims 41-43	1,22,23

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
 "E" earlier document but published on or after the international filing date
 "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
 "O" document referring to an oral disclosure, use, exhibition or other means
 "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
 "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
 "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
 "&" document member of the same patent family

Date of the actual completion of the international search

5 October 2001

Date of mailing of the international search report

16/10/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel (+31-70) 340-2040, Tx. 31 651 epo nl
 Fax (+31-70) 340-3016

Authorized officer

Seitner, I

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/17865

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
E	WO 01 60826 A (DU PONT PHARM CO) 23 August 2001 (2001-08-23) scheme 5-10 claims 1,11-13 -----	1, 22, 23

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/US 01/17865

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0007995	A	17-02-2000	AU 5337899 A BR 9912969 A CN 1311779 T EP 1102752 A1 HR 990246 A1 WO 0007995 A1	28-02-2000 25-09-2001 05-09-2001 30-05-2001 30-06-2000 17-02-2000
WO 0028331	A	18-05-2000	AU 1618000 A EP 1129355 A1 NO 20011891 A WO 0028331 A1	29-05-2000 05-09-2001 02-07-2001 18-05-2000
WO 9966934	A	29-12-1999	AU 4707999 A AU 4710199 A AU 4710499 A EP 1089977 A1 EP 1089980 A1 EP 1093372 A1 WO 9967219 A1 WO 9967221 A1 WO 9966934 A1	10-01-2000 10-01-2000 10-01-2000 11-04-2001 11-04-2001 25-04-2001 29-12-1999 29-12-1999 29-12-1999
WO 0160826	A	23-08-2001	WO 0140843 A1 WO 0160826 A2	07-06-2001 23-08-2001